

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
7 June 2001 (07.06.2001)

PCT

(10) International Publication Number  
WO 01/40269 A2

(51) International Patent Classification: C07K 14/00

Jennifer, L. [US/US]; 16677 NE 88th Street, Redmond, WA 98052 (US). WANG, Aijun [CN/US]; 3106 213th Place SE, Issaquah, WA 98029 (US).

(21) International Application Number: PCT/US00/32520

(22) International Filing Date:  
29 November 2000 (29.11.2000)

(74) Agents: POTTER, Jane, E., R.; Seed Intellectual Property Law Group PLLC, Suite 6300, 701 Fifth Avenue, Seattle, WA 98104-7092 et al. (US).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
09/451,651 30 November 1999 (30.11.1999) US  
09/510,662 22 February 2000 (22.02.2000) US  
09/523,586 10 March 2000 (10.03.2000) US  
09/545,068 7 April 2000 (07.04.2000) US  
09/571,025 15 May 2000 (15.05.2000) US

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (for all designated States except US): CORIXA CORPORATION [US/US]; Suite 200, 1124 Columbia Street, Seattle, WA 98104 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): DILLON, Davin, C. [US/US]; 18112 NW Montreux Drive, Issaquah, WA 98027 (US). DAY, Craig, H. [US/US]; 11501 Stone Ave. N., C122, Seattle, WA 98133-8317 (US). JIANG, Yuqiu [CN/US]; 5001 South 232nd Street, Kent, WA 98032 (US). HOUGHTON, Raymond, L. [US/US]; 2636 - 242nd Place SE, Bothell, WA 98021 (US). MITCHAM,

Published:

— Without international search report and to be republished upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 01/40269 A2

(54) Title: COMPOSITIONS AND METHODS FOR THERAPY AND DIAGNOSIS OF BREAST CANCER

(57) Abstract: Compositions and methods for the therapy and diagnosis of cancer, such as breast cancer, are disclosed. Compositions may comprise one or more breast tumor proteins, immunogenic portions thereof, or polynucleotides that encode such portions. Alternatively, a therapeutic composition may comprise an antigen presenting cell that expresses a breast tumor protein, or a T cell that is specific for cells expressing such a protein. Such compositions may be used, for example, for the prevention and treatment of diseases such as breast cancer. Diagnostic methods based on detecting a breast tumor protein, or mRNA encoding such a protein, in a sample are also provided.

RTS-0343

## COMPOSITIONS AND METHODS FOR THERAPY AND DIAGNOSIS OF BREAST CANCER

### TECHNICAL FIELD

The present invention relates generally to therapy and diagnosis of cancer, such as breast cancer. The invention is more specifically related to polypeptides comprising at least a portion of a breast tumor protein, and to polynucleotides encoding such polypeptides. Such polypeptides and polynucleotides may be used in vaccines and pharmaceutical compositions for prevention and treatment of breast cancer, and for the diagnosis and monitoring of such cancers.

### 10 BACKGROUND OF THE INVENTION

Breast cancer is a significant health problem for women in the United States and throughout the world. Although advances have been made in detection and treatment of the disease, breast cancer remains the second leading cause of cancer-related deaths in women, affecting more than 180,000 women in the United States each year. For women in North America, the life-time odds of getting breast cancer are now one in eight.

No vaccine or other universally successful method for the prevention or treatment of breast cancer is currently available. Management of the disease currently relies on a combination of early diagnosis (through routine breast screening procedures) and aggressive treatment, which may include one or more of a variety of treatments such as surgery, radiotherapy, chemotherapy and hormone therapy. The course of treatment for a particular breast cancer is often selected based on a variety of prognostic parameters, including an analysis of specific tumor markers. *See, e.g., Porter-Jordan and Lippman, Breast Cancer 8:73-100 (1994).* However, the use of established markers often leads to a result that is difficult to interpret, and the high mortality observed in breast cancer patients indicates that improvements are needed in the treatment, diagnosis and prevention of the disease.

Accordingly, there is a need in the art for improved methods for therapy and diagnosis of breast cancer. The present invention fulfills these needs and further provides other related advantages.

#### SUMMARY OF THE INVENTION

5 Briefly stated, the present invention provides compositions and methods for the diagnosis and therapy of cancer, such as breast cancer. In one aspect, the present invention provides polypeptides comprising at least a portion of a breast tumor protein, or a variant thereof. Certain portions and other variants are immunogenic, such that the ability of the variant to react with antigen-specific antisera is not substantially  
10 diminished. Within certain embodiments, the polypeptide comprises a sequence that is encoded by a polynucleotide sequence selected from the group consisting of: (a) sequences recited in SEQ ID NO: 1-38, 42-204, 205, 207 and 210-290; (b) variants of a sequence recited in SEQ ID NO: 1-38, 42-204, 205, 207 and 210-290; and (c) complements of a sequence of (a) or (b).

15 The present invention further provides polynucleotides that encode a polypeptide as described above, or a portion thereof (such as a portion encoding at least 15 amino acid residues of a breast tumor protein), expression vectors comprising such polynucleotides and host cells transformed or transfected with such expression vectors.

Within other aspects, the present invention provides pharmaceutical  
20 compositions comprising a polypeptide or polynucleotide as described above and a physiologically acceptable carrier.

Within a related aspect of the present invention, vaccines for prophylactic or therapeutic use are provided. Such vaccines comprise a polypeptide or polynucleotide as described above and an immunostimulant.

25 The present invention further provides pharmaceutical compositions that comprise: (a) an antibody or antigen-binding fragment thereof that specifically binds to a breast tumor protein; and (b) a physiologically acceptable carrier.

Within further aspects, the present invention provides pharmaceutical compositions comprising: (a) an antigen presenting cell that expresses a polypeptide as  
30 described above and (b) a pharmaceutically acceptable carrier or excipient. Antigen

presenting cells include dendritic cells, macrophages, monocytes, fibroblasts and B cells.

Within related aspects, vaccines are provided that comprise: (a) an antigen presenting cell that expresses a polypeptide as described above and (b) an immunostimulant.

The present invention further provides, in other aspects, fusion proteins that comprise at least one polypeptide as described above, as well as polynucleotides encoding such fusion proteins.

Within related aspects, pharmaceutical compositions comprising a fusion protein, or a polynucleotide encoding a fusion protein, in combination with a physiologically acceptable carrier are provided.

Vaccines are further provided, within other aspects, that comprise a fusion protein, or a polynucleotide encoding a fusion protein, in combination with an immunostimulant.

Within further aspects, the present invention provides methods for inhibiting the development of a cancer in a patient, comprising administering to a patient a pharmaceutical composition or vaccine as recited above.

The present invention further provides, within other aspects, methods for removing tumor cells from a biological sample, comprising contacting a biological sample with T cells that specifically react with a breast tumor protein, wherein the step of contacting is performed under conditions and for a time sufficient to permit the removal of cells expressing the protein from the sample.

Within related aspects, methods are provided for inhibiting the development of a cancer in a patient, comprising administering to a patient a biological sample treated as described above.

Methods are further provided, within other aspects, for stimulating and/or expanding T cells specific for a breast tumor protein, comprising contacting T cells with one or more of: (i) a polypeptide as described above; (ii) a polynucleotide encoding such a polypeptide; and/or (iii) an antigen presenting cell that expresses such a polypeptide; under conditions and for a time sufficient to permit the stimulation and/or



expansion of T cells. Isolated T cell populations comprising T cells prepared as described above are also provided.

Within further aspects, the present invention provides methods for inhibiting the development of a cancer in a patient, comprising administering to a  
5 patient an effective amount of a T cell population as described above.

The present invention further provides methods for inhibiting the development of a cancer in a patient, comprising the steps of: (a) incubating CD4<sup>+</sup> and/or CD8<sup>+</sup> T cells isolated from a patient with one or more of: (i) a polypeptide comprising at least an immunogenic portion of a breast tumor protein; (ii) a  
10 polynucleotide encoding such a polypeptide; and (iii) an antigen-presenting cell that expresses such a polypeptide; and (b) administering to the patient an effective amount of the proliferated T cells, and thereby inhibiting the development of a cancer in the patient. Proliferated cells may, but need not, be cloned prior to administration to the patient.

15 Within further aspects, the present invention provides methods for determining the presence or absence of a cancer in a patient, comprising: (a) contacting a biological sample obtained from a patient with a binding agent that binds to a polypeptide as recited above; (b) detecting in the sample an amount of polypeptide that binds to the binding agent; and (c) comparing the amount of polypeptide with a  
20 predetermined cut-off value, and therefrom determining the presence or absence of a cancer in the patient. Within preferred embodiments, the binding agent is an antibody, more preferably a monoclonal antibody. The cancer may be breast cancer.

The present invention also provides, within other aspects, methods for monitoring the progression of a cancer in a patient. Such methods comprise the steps  
25 of: (a) contacting a biological sample obtained from a patient at a first point in time with a binding agent that binds to a polypeptide as recited above; (b) detecting in the sample an amount of polypeptide that binds to the binding agent; (c) repeating steps (a) and (b) using a biological sample obtained from the patient at a subsequent point in time; and (d) comparing the amount of polypeptide detected in step (c) with the amount  
30 detected in step (b) and therefrom monitoring the progression of the cancer in the patient.

The present invention further provides, within other aspects, methods for determining the presence or absence of a cancer in a patient, comprising the steps of: (a) contacting a biological sample obtained from a patient with an oligonucleotide that hybridizes to a polynucleotide that encodes a breast tumor protein; (b) detecting in the sample a level of a polynucleotide, preferably mRNA, that hybridizes to the oligonucleotide; and (c) comparing the level of polynucleotide that hybridizes to the oligonucleotide with a predetermined cut-off value, and therefrom determining the presence or absence of a cancer in the patient. Within certain embodiments, the amount of mRNA is detected via polymerase chain reaction using, for example, at least one oligonucleotide primer that hybridizes to a polynucleotide encoding a polypeptide as recited above, or a complement of such a polynucleotide. Within other embodiments, the amount of mRNA is detected using a hybridization technique, employing an oligonucleotide probe that hybridizes to a polynucleotide that encodes a polypeptide as recited above, or a complement of such a polynucleotide.

In related aspects, methods are provided for monitoring the progression of a cancer in a patient, comprising the steps of: (a) contacting a biological sample obtained from a patient with an oligonucleotide that hybridizes to a polynucleotide that encodes a breast tumor protein; (b) detecting in the sample an amount of a polynucleotide that hybridizes to the oligonucleotide; (c) repeating steps (a) and (b) using a biological sample obtained from the patient at a subsequent point in time; and (d) comparing the amount of polynucleotide detected in step (c) with the amount detected in step (b) and therefrom monitoring the progression of the cancer in the patient.

Within further aspects, the present invention provides antibodies, such as monoclonal antibodies, that bind to a polypeptide as described above, as well as diagnostic kits comprising such antibodies. Diagnostic kits comprising one or more oligonucleotide probes or primers as described above are also provided.

These and other aspects of the present invention will become apparent upon reference to the following detailed description. All references disclosed herein are hereby incorporated by reference in their entirety as if each was incorporated individually.

**SEQUENCE IDENTIFIERS**

- SEQ ID NO: 1 is the determined cDNA sequence for clone 26915.
- SEQ ID NO: 2 is the determined cDNA sequence for clone 26914.
- SEQ ID NO: 3 is the determined cDNA sequence for clone 26673.
- 5 SEQ ID NO: 4 is the determined cDNA sequence for clone 26672.
- SEQ ID NO: 5 is the determined cDNA sequence for clone 26671.
- SEQ ID NO: 6 is the determined cDNA sequence for clone 26670.
- SEQ ID NO: 7 is the determined cDNA sequence for clone 26669.
- SEQ ID NO: 8 is a first determined cDNA sequence for clone 26668.
- 10 SEQ ID NO: 9 is a second determined cDNA sequence for clone 26668.
- SEQ ID NO: 10 is the determined cDNA sequence for clone 26667.
- SEQ ID NO: 11 is the determined cDNA sequence for clone 26666.
- SEQ ID NO: 12 is the determined cDNA sequence for clone 26665.
- SEQ ID NO: 13 is the determined cDNA sequence for clone 26664.
- 15 SEQ ID NO: 14 is the determined cDNA sequence for clone 26662.
- SEQ ID NO: 15 is the determined cDNA sequence for clone 26661.
- SEQ ID NO: 16 is the determined cDNA sequence for clone 26660.
- SEQ ID NO: 17 is the determined cDNA sequence for clone 26603.
- SEQ ID NO: 18 is the determined cDNA sequence for clone 26601.
- 20 SEQ ID NO: 19 is the determined cDNA sequence for clone 26600.
- SEQ ID NO: 20 is the determined cDNA sequence for clone 26587.
- SEQ ID NO: 21 is the determined cDNA sequence for clone 26586.
- SEQ ID NO: 22 is the determined cDNA sequence for clone 26584.
- SEQ ID NO: 23 is the determined cDNA sequence for clone 26583.
- 25 SEQ ID NO: 24 is the determined cDNA sequence for clone 26580.
- SEQ ID NO: 25 is the determined cDNA sequence for clone 26579.
- SEQ ID NO: 26 is the determined cDNA sequence for clone 26577.
- SEQ ID NO: 27 is the determined cDNA sequence for clone 26575.
- SEQ ID NO: 28 is the determined cDNA sequence for clone 26574.
- 30 SEQ ID NO: 29 is the determined cDNA sequence for clone 26573.
- SEQ ID NO: 30 is the determined cDNA sequence for clone 25612.

- SEQ ID NO: 31 is the determined cDNA sequence for clone 22295.
- SEQ ID NO: 32 is the determined cDNA sequence for clone 22301.
- SEQ ID NO: 33 is the determined cDNA sequence for clone 22298.
- SEQ ID NO: 34 is the determined cDNA sequence for clone 22297.
- 5 SEQ ID NO: 35 is the determined cDNA sequence for clone 22303.
- SEQ ID NO: 36 is the determined cDNA sequence for a first GABA<sub>A</sub> receptor clone.
- SEQ ID NO: 37 is the determined cDNA sequence for a second GABA<sub>A</sub> receptor clone.
- 10 SEQ ID NO: 38 is the determined cDNA sequence for a third GABA<sub>A</sub> receptor clone.
- SEQ ID NO: 39 is the amino acid sequence encoded by SEQ ID NO: 36.
- SEQ ID NO: 40 is the amino acid sequence encoded by SEQ ID NO: 37.
- SEQ ID NO: 41 is the amino acid sequence encoded by SEQ ID NO: 38.
- 15 SEQ ID NO: 42 is the determined cDNA sequence for contig 1.
- SEQ ID NO: 43 is the determined cDNA sequence for contig 2.
- SEQ ID NO: 44 is the determined cDNA sequence for contig 3.
- SEQ ID NO: 45 is the determined cDNA sequence for contig 4.
- SEQ ID NO: 46 is the determined cDNA sequence for contig 5.
- 20 SEQ ID NO: 47 is the determined cDNA sequence for contig 6.
- SEQ ID NO: 48 is the determined cDNA sequence for contig 7.
- SEQ ID NO: 49 is the determined cDNA sequence for contig 8.
- SEQ ID NO: 50 is the determined cDNA sequence for contig 9.
- SEQ ID NO: 51 is the determined cDNA sequence for contig 10.
- 25 SEQ ID NO: 52 is the determined cDNA sequence for contig 11.
- SEQ ID NO: 53 is the determined cDNA sequence for contig 12.
- SEQ ID NO: 54 is the determined cDNA sequence for contig 13.
- SEQ ID NO: 55 is the determined cDNA sequence for contig 14.
- SEQ ID NO: 56 is the determined cDNA sequence for contig 15.
- 30 SEQ ID NO: 57 is the determined cDNA sequence for contig 16.
- SEQ ID NO: 58 is the determined cDNA sequence for contig 17.

SEQ ID NO: 59 is the determined cDNA sequence for contig 18.  
SEQ ID NO: 60 is the determined cDNA sequence for contig 19.  
SEQ ID NO: 61 is the determined cDNA sequence for contig 20.  
SEQ ID NO: 62 is the determined cDNA sequence for contig 21.  
5 SEQ ID NO: 63 is the determined cDNA sequence for contig 22.  
SEQ ID NO: 64 is the determined cDNA sequence for contig 23.  
SEQ ID NO: 65 is the determined cDNA sequence for contig 24.  
SEQ ID NO: 66 is the determined cDNA sequence for contig 25.  
SEQ ID NO: 67 is the determined cDNA sequence for contig 26.  
10 SEQ ID NO: 68 is the determined cDNA sequence for contig 27.  
SEQ ID NO: 69 is the determined cDNA sequence for contig 28.  
SEQ ID NO: 70 is the determined cDNA sequence for contig 29.  
SEQ ID NO: 71 is the determined cDNA sequence for contig 30.  
SEQ ID NO: 72 is the determined cDNA sequence for contig 31.  
15 SEQ ID NO: 73 is the determined cDNA sequence for contig 32.  
SEQ ID NO: 74 is the determined cDNA sequence for contig 33.  
SEQ ID NO: 75 is the determined cDNA sequence for contig 34.  
SEQ ID NO: 76 is the determined cDNA sequence for contig 35.  
SEQ ID NO: 77 is the determined cDNA sequence for contig 36.  
20 SEQ ID NO: 78 is the determined cDNA sequence for contig 37.  
SEQ ID NO: 79 is the determined cDNA sequence for contig 38.  
SEQ ID NO: 80 is the determined cDNA sequence for contig 39.  
SEQ ID NO: 81 is the determined cDNA sequence for contig 40.  
SEQ ID NO: 82 is the determined cDNA sequence for contig 41.  
25 SEQ ID NO: 83 is the determined cDNA sequence for contig 42.  
SEQ ID NO: 84 is the determined cDNA sequence for contig 43.  
SEQ ID NO: 85 is the determined cDNA sequence for contig 44.  
SEQ ID NO: 85 is the determined cDNA sequence for contig 45.  
SEQ ID NO: 85 is the determined cDNA sequence for contig 46.  
30 SEQ ID NO: 88 is the determined cDNA sequence for contig 47.  
SEQ ID NO: 89 is the determined cDNA sequence for contig 48.

SEQ ID NO: 90 is the determined cDNA sequence for contig 49.  
SEQ ID NO: 91 is the determined cDNA sequence for contig 50.  
SEQ ID NO: 92 is the determined cDNA sequence for contig 51.  
SEQ ID NO: 93 is the determined cDNA sequence for contig 52.  
5 SEQ ID NO: 94 is the determined cDNA sequence for contig 53.  
SEQ ID NO: 95 is the determined cDNA sequence for contig 54.  
SEQ ID NO: 96 is the determined cDNA sequence for contig 55.  
SEQ ID NO: 97 is the determined cDNA sequence for contig 56.  
SEQ ID NO: 98 is the determined cDNA sequence for contig 57.  
10 SEQ ID NO: 99 is the determined cDNA sequence for contig 58.  
SEQ ID NO: 100 is the determined cDNA sequence for contig 59.  
SEQ ID NO: 101 is the determined cDNA sequence for contig 60.  
SEQ ID NO: 102 is the determined cDNA sequence for contig 61.  
SEQ ID NO: 103 is the determined cDNA sequence for contig 62.  
15 SEQ ID NO: 104 is the determined cDNA sequence for contig 63.  
SEQ ID NO: 105 is the determined cDNA sequence for contig 64.  
SEQ ID NO: 106 is the determined cDNA sequence for contig 65.  
SEQ ID NO: 107 is the determined cDNA sequence for contig 66.  
SEQ ID NO: 108 is the determined cDNA sequence for contig 67.  
20 SEQ ID NO: 109 is the determined cDNA sequence for contig 68.  
SEQ ID NO: 110 is the determined cDNA sequence for contig 69.  
SEQ ID NO: 111 is the determined cDNA sequence for contig 70.  
SEQ ID NO: 112 is the determined cDNA sequence for contig 71.  
SEQ ID NO: 113 is the determined cDNA sequence for contig 72.  
25 SEQ ID NO: 114 is the determined cDNA sequence for contig 73.  
SEQ ID NO: 115 is the determined cDNA sequence for contig 74.  
SEQ ID NO: 116 is the determined cDNA sequence for contig 75.  
SEQ ID NO: 117 is the determined cDNA sequence for contig 76.  
SEQ ID NO: 118 is the determined cDNA sequence for contig 77.  
30 SEQ ID NO: 119 is the determined cDNA sequence for contig 78.  
SEQ ID NO: 120 is the determined cDNA sequence for contig 79.

SEQ ID NO: 121 is the determined cDNA sequence for contig 80.  
SEQ ID NO: 122 is the determined cDNA sequence for contig 81.  
SEQ ID NO: 123 is the determined cDNA sequence for contig 82.  
SEQ ID NO: 124 is the determined cDNA sequence for contig 83.  
5 SEQ ID NO: 125 is the determined cDNA sequence for contig 84.  
SEQ ID NO: 126 is the determined cDNA sequence for contig 85.  
SEQ ID NO: 127 is the determined cDNA sequence for contig 86.  
SEQ ID NO: 128 is the determined cDNA sequence for contig 87.  
SEQ ID NO: 129 is the determined cDNA sequence for contig 88.  
10 SEQ ID NO: 130 is the determined cDNA sequence for contig 89.  
SEQ ID NO: 131 is the determined cDNA sequence for contig 90.  
SEQ ID NO: 132 is the determined cDNA sequence for contig 91.  
SEQ ID NO: 133 is the determined cDNA sequence for contig 92.  
SEQ ID NO: 134 is the determined cDNA sequence for contig 93.  
15 SEQ ID NO: 135 is the determined cDNA sequence for contig 94.  
SEQ ID NO: 136 is the determined cDNA sequence for contig 95.  
SEQ ID NO: 137 is the determined cDNA sequence for contig 96.  
SEQ ID NO: 138 is the determined cDNA sequence for clone 47589.  
SEQ ID NO: 139 is the determined cDNA sequence for clone 47578.  
20 SEQ ID NO: 140 is the determined cDNA sequence for clone 47602.  
SEQ ID NO: 141 is the determined cDNA sequence for clone 47593.  
SEQ ID NO: 142 is the determined cDNA sequence for clone 47583.  
SEQ ID NO: 143 is the determined cDNA sequence for clone 47624.  
SEQ ID NO: 144 is the determined cDNA sequence for clone 47622.  
25 SEQ ID NO: 145 is the determined cDNA sequence for clone 47649.  
SEQ ID NO: 146 is the determined cDNA sequence for clone 48955.  
SEQ ID NO: 147 is the determined cDNA sequence for clone 48962.  
SEQ ID NO: 148 is the determined cDNA sequence for clone 48964.  
SEQ ID NO: 149 is the determined cDNA sequence for clone 48987.  
30 SEQ ID NO: 150 is the determined cDNA sequence for clone 49002.  
SEQ ID NO: 151 is the determined cDNA sequence for clone 48950.

SEQ ID NO: 152 is the determined cDNA sequence for clone 48934.  
SEQ ID NO: 153 is the determined cDNA sequence for clone 48960.  
SEQ ID NO: 154 is the determined cDNA sequence for clone 48931.  
SEQ ID NO: 155 is the determined cDNA sequence for clone 48935.  
5 SEQ ID NO: 156 is the determined cDNA sequence for clone 48940.  
SEQ ID NO: 157 is the determined cDNA sequence for clone 48936.  
SEQ ID NO: 158 is the determined cDNA sequence for clone 48930.  
SEQ ID NO: 159 is the determined cDNA sequence for clone 48956.  
SEQ ID NO: 160 is the determined cDNA sequence for clone 48959.  
10 SEQ ID NO: 161 is the determined cDNA sequence for clone 48949.  
SEQ ID NO: 162 is the determined cDNA sequence for clone 48965.  
SEQ ID NO: 163 is the determined cDNA sequence for clone 48970.  
SEQ ID NO: 164 is the determined cDNA sequence for clone 48984.  
SEQ ID NO: 165 is the determined cDNA sequence for clone 48969.  
15 SEQ ID NO: 166 is the determined cDNA sequence for clone 48978.  
SEQ ID NO: 167 is the determined cDNA sequence for clone 48968.  
SEQ ID NO: 168 is the determined cDNA sequence for clone 48929.  
SEQ ID NO: 169 is the determined cDNA sequence for clone 48937.  
SEQ ID NO: 170 is the determined cDNA sequence for clone 48982.  
20 SEQ ID NO: 171 is the determined cDNA sequence for clone 48983.  
SEQ ID NO: 172 is the determined cDNA sequence for clone 48997.  
SEQ ID NO: 173 is the determined cDNA sequence for clone 48992.  
SEQ ID NO: 174 is the determined cDNA sequence for clone 49006.  
SEQ ID NO: 175 is the determined cDNA sequence for clone 48994.  
25 SEQ ID NO: 176 is the determined cDNA sequence for clone 49013.  
SEQ ID NO: 177 is the determined cDNA sequence for clone 49008.  
SEQ ID NO: 178 is the determined cDNA sequence for clone 48990.  
SEQ ID NO: 179 is the determined cDNA sequence for clone 48989.  
SEQ ID NO: 180 is the determined cDNA sequence for clone 49014.  
30 SEQ ID NO: 181 is the determined cDNA sequence for clone 48988.  
SEQ ID NO: 182 is the determined cDNA sequence for clone 49018.



SEQ ID NO: 183 is the determined cDNA sequence for clone 6921.  
 SEQ ID NO: 184 is the determined cDNA sequence for clone 6837.  
 SEQ ID NO: 185 is the determined cDNA sequence for clone 6840.  
 SEQ ID NO: 186 is the determined cDNA sequence for clone 6844.  
 5 SEQ ID NO: 187 is the determined cDNA sequence for clone 6854.  
 SEQ ID NO: 188 is the determined cDNA sequence for clone 6872.  
 SEQ ID NO: 189 is the determined cDNA sequence for clone 6906.  
 SEQ ID NO: 190 is the determined cDNA sequence for clone 6908.  
 SEQ ID NO: 191 is the determined cDNA sequence for clone 6910.  
 10 SEQ ID NO: 192 is the determined cDNA sequence for clone 6912.  
 SEQ ID NO: 193 is the determined cDNA sequence for clone 6913.  
 SEQ ID NO: 194 is the determined cDNA sequence for clone 6914.  
 SEQ ID NO: 195 is the determined cDNA sequence for clone 6916.  
 SEQ ID NO: 196 is the determined cDNA sequence for clone 6918.  
 15 SEQ ID NO: 197 is the determined cDNA sequence for clone 6924.  
 SEQ ID NO: 198 is the determined cDNA sequence for clone 6928.  
 SEQ ID NO: 199 is the determined cDNA sequence for clone 6978A.  
 SEQ ID NO: 200 is the determined cDNA sequence for clone 6978B.  
 SEQ ID NO: 201 is the determined cDNA sequence for clone 6982A.  
 20 SEQ ID NO: 202 is the determined cDNA sequence for clone 6982B.  
 SEQ ID NO: 203 is the determined cDNA sequence for clone 6850.  
 SEQ ID NO: 204 is the determined cDNA sequence for clone 6860.  
 SEQ ID NO: 205 is the determined cDNA sequence for O772P.  
 SEQ ID NO: 206 is the amino acid sequence encoded by SEQ ID NO:  
 25 205.  
 SEQ ID NO: 207 is the full-length cDNA sequence for O8E.  
 SEQ ID NO: 208 is a first amino acid sequence encoded by SEQ ID NO:  
 207.  
 SEQ ID NO: 209 is a second amino acid sequence encoded by SEQ ID  
 30 NO: 209.

SEQ ID NO: 210-290 are determined cDNA sequence of breast-tumor specific clones.

#### DETAILED DESCRIPTION OF THE INVENTION

As noted above, the present invention is generally directed to compositions and methods for using the compositions, for example in the therapy and diagnosis of cancer, such as breast cancer. Certain illustrative compositions described herein include breast tumor polypeptides, polynucleotides encoding such polypeptides, binding agents such as antibodies, antigen presenting cells (APCs) and/or immune system cells (e.g., T cells). A "breast tumor protein," as the term is used herein, refers generally to a protein that is expressed in breast tumor cells at a level that is at least two fold, and preferably at least five fold, greater than the level of expression in other normal tissues, as determined using a representative assay provided herein. Certain breast tumor proteins are tumor proteins that react detectably (within an immunoassay, such as an ELISA or Western blot) with antisera of a patient afflicted with breast cancer.

Therefore, in accordance with the above, and as described further below, the present invention provides illustrative polynucleotide compositions having sequences set forth in SEQ ID NO:1-38, 42-204, 205, 207 and 210-290, polypeptides encoded by such polynucleotides, antibody compositions capable of binding such polypeptides, and numerous additional embodiments employing such compositions, for example in the detection, diagnosis and/or therapy of human breast cancer.

#### POLYNUCLEOTIDE COMPOSITIONS

As used herein, the terms "DNA segment" and "polynucleotide" refer to a DNA molecule that has been isolated free of total genomic DNA of a particular species. Therefore, a DNA segment encoding a polypeptide refers to a DNA segment that contains one or more coding sequences yet is substantially isolated away from, or purified free from, total genomic DNA of the species from which the DNA segment is obtained. Included within the terms "DNA segment" and "polynucleotide" are DNA

segments and smaller fragments of such segments, and also recombinant vectors, including, for example, plasmids, cosmids, phagemids, phage, viruses, and the like.

As will be understood by those skilled in the art, the DNA segments of this invention can include genomic sequences, extra-genomic and plasmid-encoded sequences and smaller engineered gene segments that express, or may be adapted to express, proteins, polypeptides, peptides and the like. Such segments may be naturally isolated, or modified synthetically by the hand of man.

"Isolated," as used herein, means that a polynucleotide is substantially away from other coding sequences, and that the DNA segment does not contain large portions of unrelated coding DNA, such as large chromosomal fragments or other functional genes or polypeptide coding regions. Of course, this refers to the DNA segment as originally isolated, and does not exclude genes or coding regions later added to the segment by the hand of man.

As will be recognized by the skilled artisan, polynucleotides may be single-stranded (coding or antisense) or double-stranded, and may be DNA (genomic, cDNA or synthetic) or RNA molecules. RNA molecules include HnRNA molecules, which contain introns and correspond to a DNA molecule in a one-to-one manner, and mRNA molecules, which do not contain introns. Additional coding or non-coding sequences may, but need not, be present within a polynucleotide of the present invention, and a polynucleotide may, but need not, be linked to other molecules and/or support materials.

Polynucleotides may comprise a native sequence (*i.e.*, an endogenous sequence that encodes a breast tumor protein or a portion thereof) or may comprise a variant, or a biological or antigenic functional equivalent of such a sequence. Polynucleotide variants may contain one or more substitutions, additions, deletions and/or insertions, as further described below, preferably such that the immunogenicity of the encoded polypeptide is not diminished, relative to a native tumor protein. The effect on the immunogenicity of the encoded polypeptide may generally be assessed as described herein. The term "variants" also encompasses homologous genes of xenogenic origin.

When comparing polynucleotide or polypeptide sequences, two sequences are said to be "identical" if the sequence of nucleotides or amino acids in the two sequences is the same when aligned for maximum correspondence, as described below. Comparisons between two sequences are typically performed by comparing the sequences over a comparison window to identify and compare local regions of sequence similarity. A "comparison window" as used herein, refers to a segment of at least about 20 contiguous positions, usually 30 to about 75, 40 to about 50, in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned.

Optimal alignment of sequences for comparison may be conducted using the Megalign program in the Lasergene suite of bioinformatics software (DNASTAR, Inc., Madison, WI), using default parameters. This program embodies several alignment schemes described in the following references: Dayhoff, M.O. (1978) A model of evolutionary change in proteins – Matrices for detecting distant relationships. In Dayhoff, M.O. (ed.) *Atlas of Protein Sequence and Structure*, National Biomedical Research Foundation, Washington DC Vol. 5, Suppl. 3, pp. 345-358; Hein J. (1990) *Unified Approach to Alignment and Phylogenies* pp. 626-645 *Methods in Enzymology* vol. 183, Academic Press, Inc., San Diego, CA; Higgins, D.G. and Sharp, P.M. (1989) *CABIOS* 5:151-153; Myers, E.W. and Muller W. (1988) *CABIOS* 4:11-17; Robinson, E.D. (1971) *Comb. Theor* 11:105; Santou, N. Nes, M. (1987) *Mol. Biol. Evol.* 4:406-425; Sneath, P.H.A. and Sokal, R.R. (1973) *Numerical Taxonomy – the Principles and Practice of Numerical Taxonomy*, Freeman Press, San Francisco, CA; Wilbur, W.J. and Lipman, D.J. (1983) *Proc. Natl. Acad. Sci. USA* 80:726-730.

Alternatively, optimal alignment of sequences for comparison may be conducted by the local identity algorithm of Smith and Waterman (1981) *Add. APL. Math* 2:482, by the identity alignment algorithm of Needleman and Wunsch (1970) *J. Mol. Biol.* 48:443, by the search for similarity methods of Pearson and Lipman (1988) *Proc. Natl. Acad. Sci. USA* 85: 2444, by computerized implementations of these algorithms (GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), 575 Science Dr., Madison, WI), or by inspection.

One preferred example of algorithms that are suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul et al. (1977) *Nucl. Acids Res.* 25:3389-3402 and Altschul et al. (1990) *J. Mol. Biol.* 215:403-410, respectively. BLAST and BLAST 2.0 can be used, for example with the parameters described herein, to determine percent sequence identity for the polynucleotides and polypeptides of the invention. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information. In one illustrative example, cumulative scores can be calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always >0) and N (penalty score for mismatching residues; always <0). For amino acid sequences, a scoring matrix can be used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, and expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff and Henikoff (1989) *Proc. Natl. Acad. Sci. USA* 89:10915) alignments, (B) of 50, expectation (E) of 10, M=5, N=4 and a comparison of both strands.

Preferably, the "percentage of sequence identity" is determined by comparing two optimally aligned sequences over a window of comparison of at least 20 positions, wherein the portion of the polynucleotide or polypeptide sequence in the comparison window may comprise additions or deletions (*i.e.*, gaps) of 20 percent or less, usually 5 to 15 percent, or 10 to 12 percent, as compared to the reference sequences (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid bases or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the reference sequence (*i.e.*, the window size) and multiplying the results by 100 to yield the percentage of sequence identity.

Therefore, the present invention encompasses polynucleotide and polypeptide sequences having substantial identity to the sequences disclosed herein, for example those comprising at least 50% sequence identity, preferably at least 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% or higher, sequence  
5 identity compared to a polynucleotide or polypeptide sequence of this invention using the methods described herein, (e.g., BLAST analysis using standard parameters, as described below). One skilled in this art will recognize that these values can be appropriately adjusted to determine corresponding identity of proteins encoded by two nucleotide sequences by taking into account codon degeneracy, amino acid similarity,  
10 reading frame positioning and the like.

In additional embodiments, the present invention provides isolated polynucleotides and polypeptides comprising various lengths of contiguous stretches of sequence identical to or complementary to one or more of the sequences disclosed herein. For example, polynucleotides are provided by this invention that comprise at  
15 least about 15, 20, 30, 40, 50, 75, 100, 150, 200, 300, 400, 500 or 1000 or more contiguous nucleotides of one or more of the sequences disclosed herein as well as all intermediate lengths there between. It will be readily understood that "intermediate lengths", in this context, means any length between the quoted values, such as 16, 17, 18, 19, *etc.*; 21, 22, 23, *etc.*; 30, 31, 32, *etc.*; 50, 51, 52, 53, *etc.*; 100, 101, 102, 103,  
20 *etc.*; 150, 151, 152, 153, *etc.*; including all integers through 200-500; 500-1,000, and the like.

The polynucleotides of the present invention, or fragments thereof, regardless of the length of the coding sequence itself, may be combined with other DNA sequences, such as promoters, polyadenylation signals, additional restriction  
25 enzyme sites, multiple cloning sites, other coding segments, and the like, such that their overall length may vary considerably. It is therefore contemplated that a nucleic acid fragment of almost any length may be employed, with the total length preferably being limited by the ease of preparation and use in the intended recombinant DNA protocol. For example, illustrative DNA segments with total lengths of about 10,000, about 5000,  
30 about 3000, about 2,000, about 1,000, about 500, about 200, about 100, about 50 base

pairs in length, and the like, (including all intermediate lengths) are contemplated to be useful in many implementations of this invention.

In other embodiments, the present invention is directed to polynucleotides that are capable of hybridizing under moderately stringent conditions to a polynucleotide sequence provided herein, or a fragment thereof, or a complementary sequence thereof. Hybridization techniques are well known in the art of molecular biology. For purposes of illustration, suitable moderately stringent conditions for testing the hybridization of a polynucleotide of this invention with other polynucleotides include prewashing in a solution of 5 X SSC, 0.5% SDS, 1.0 mM EDTA (pH 8.0); hybridizing at 50°C-65°C, 5 X SSC, overnight; followed by washing twice at 65°C for 20 minutes with each of 2X, 0.5X and 0.2X SSC containing 0.1% SDS.

Moreover, it will be appreciated by those of ordinary skill in the art that, as a result of the degeneracy of the genetic code, there are many nucleotide sequences that encode a polypeptide as described herein. Some of these polynucleotides bear minimal homology to the nucleotide sequence of any native gene. Nonetheless, polynucleotides that vary due to differences in codon usage are specifically contemplated by the present invention. Further, alleles of the genes comprising the polynucleotide sequences provided herein are within the scope of the present invention. Alleles are endogenous genes that are altered as a result of one or more mutations, such as deletions, additions and/or substitutions of nucleotides. The resulting mRNA and protein may, but need not, have an altered structure or function. Alleles may be identified using standard techniques (such as hybridization, amplification and/or database sequence comparison).

## 25 PROBES AND PRIMERS

In other embodiments of the present invention, the polynucleotide sequences provided herein can be advantageously used as probes or primers for nucleic acid hybridization. As such, it is contemplated that nucleic acid segments that comprise a sequence region of at least about 15 nucleotide long contiguous sequence that has the same sequence as, or is complementary to, a 15 nucleotide long contiguous sequence

disclosed herein will find particular utility. Longer contiguous identical or complementary sequences, *e.g.*, those of about 20, 30, 40, 50, 100, 200, 500, 1000 (including all intermediate lengths) and even up to full length sequences will also be of use in certain embodiments.

5           The ability of such nucleic acid probes to specifically hybridize to a sequence of interest will enable them to be of use in detecting the presence of complementary sequences in a given sample. However, other uses are also envisioned, such as the use of the sequence information for the preparation of mutant species primers, or primers for use in preparing other genetic constructions.

10           Polynucleotide molecules having sequence regions consisting of contiguous nucleotide stretches of 10-14, 15-20, 30, 50, or even of 100-200 nucleotides or so (including intermediate lengths as well), identical or complementary to a polynucleotide sequence disclosed herein, are particularly contemplated as hybridization probes for use in, *e.g.*, Southern and Northern blotting. This would allow  
15 a gene product, or fragment thereof, to be analyzed, both in diverse cell types and also in various bacterial cells. The total size of fragment, as well as the size of the complementary stretch(es), will ultimately depend on the intended use or application of the particular nucleic acid segment. Smaller fragments will generally find use in hybridization embodiments, wherein the length of the contiguous complementary  
20 region may be varied, such as between about 15 and about 100 nucleotides, but larger contiguous complementarity stretches may be used, according to the length complementary sequences one wishes to detect.

          The use of a hybridization probe of about 15-25 nucleotides in length allows the formation of a duplex molecule that is both stable and selective. Molecules  
25 having contiguous complementary sequences over stretches greater than 15 bases in length are generally preferred, though, in order to increase stability and selectivity of the hybrid, and thereby improve the quality and degree of specific hybrid molecules obtained. One will generally prefer to design nucleic acid molecules having gene-complementary stretches of 15 to 25 contiguous nucleotides, or even longer where  
30 desired.



Hybridization probes may be selected from any portion of any of the sequences disclosed herein. All that is required is to review the sequence set forth in SEQ ID NO: 1-38, 42-204, 205, 207 and 210-290, or to any continuous portion of the sequence, from about 15-25 nucleotides in length up to and including the full length  
5 sequence, that one wishes to utilize as a probe or primer. The choice of probe and primer sequences may be governed by various factors. For example, one may wish to employ primers from towards the termini of the total sequence.

Small polynucleotide segments or fragments may be readily prepared by, for example, directly synthesizing the fragment by chemical means, as is commonly  
10 practiced using an automated oligonucleotide synthesizer. Also, fragments may be obtained by application of nucleic acid reproduction technology, such as the PCR™ technology of U. S. Patent 4,683,202 (incorporated herein by reference), by introducing selected sequences into recombinant vectors for recombinant production, and by other recombinant DNA techniques generally known to those of skill in the art of molecular  
15 biology.

The nucleotide sequences of the invention may be used for their ability to selectively form duplex molecules with complementary stretches of the entire gene or gene fragments of interest. Depending on the application envisioned, one will typically desire to employ varying conditions of hybridization to achieve varying degrees of  
20 selectivity of probe towards target sequence. For applications requiring high selectivity, one will typically desire to employ relatively stringent conditions to form the hybrids, *e.g.*, one will select relatively low salt and/or high temperature conditions, such as provided by a salt concentration of from about 0.02 M to about 0.15 M salt at temperatures of from about 50°C to about 70°C. Such selective conditions tolerate  
25 little, if any, mismatch between the probe and the template or target strand, and would be particularly suitable for isolating related sequences.

Of course, for some applications, for example, where one desires to prepare mutants employing a mutant primer strand hybridized to an underlying template, less stringent (reduced stringency) hybridization conditions will typically be  
30 needed in order to allow formation of the heteroduplex. In these circumstances, one may desire to employ salt conditions such as those of from about 0.15 M to about 0.9 M

salt, at temperatures ranging from about 20°C to about 55°C. Cross-hybridizing species can thereby be readily identified as positively hybridizing signals with respect to control hybridizations. In any case, it is generally appreciated that conditions can be rendered more stringent by the addition of increasing amounts of formamide, which serves to  
5 destabilize the hybrid duplex in the same manner as increased temperature. Thus, hybridization conditions can be readily manipulated, and thus will generally be a method of choice depending on the desired results.

#### POLYNUCLEOTIDE IDENTIFICATION AND CHARACTERIZATION

Polynucleotides may be identified, prepared and/or manipulated using  
10 any of a variety of well established techniques. For example, a polynucleotide may be identified, as described in more detail below, by screening a microarray of cDNAs for tumor-associated expression (*i.e.*, expression that is at least two fold greater in a tumor than in normal tissue, as determined using a representative assay provided herein). Such screens may be performed, for example, using a Synteni microarray (Palo Alto,  
15 CA) according to the manufacturer's instructions (and essentially as described by Schena et al., *Proc. Natl. Acad. Sci. USA* 93:10614-10619, 1996 and Heller et al., *Proc. Natl. Acad. Sci. USA* 94:2150-2155, 1997). Alternatively, polynucleotides may be amplified from cDNA prepared from cells expressing the proteins described herein, such as breast tumor cells. Such polynucleotides may be amplified via polymerase  
20 chain reaction (PCR). For this approach, sequence-specific primers may be designed based on the sequences provided herein, and may be purchased or synthesized.

An amplified portion of a polynucleotide of the present invention may be used to isolate a full length gene from a suitable library (*e.g.*, a breast tumor cDNA library) using well known techniques. Within such techniques, a library (cDNA or  
25 genomic) is screened using one or more polynucleotide probes or primers suitable for amplification. Preferably, a library is size-selected to include larger molecules. Random primed libraries may also be preferred for identifying 5' and upstream regions of genes. Genomic libraries are preferred for obtaining introns and extending 5' sequences.

For hybridization techniques, a partial sequence may be labeled (e.g., by nick-translation or end-labeling with  $^{32}\text{P}$ ) using well known techniques. A bacterial or bacteriophage library is then generally screened by hybridizing filters containing denatured bacterial colonies (or lawns containing phage plaques) with the labeled probe  
5 (see Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratories, Cold Spring Harbor, NY, 1989). Hybridizing colonies or plaques are selected and expanded, and the DNA is isolated for further analysis. cDNA clones may be analyzed to determine the amount of additional sequence by, for example, PCR using a primer from the partial sequence and a primer from the vector. Restriction maps and  
10 partial sequences may be generated to identify one or more overlapping clones. The complete sequence may then be determined using standard techniques, which may involve generating a series of deletion clones. The resulting overlapping sequences can then be assembled into a single contiguous sequence. A full length cDNA molecule can be generated by ligating suitable fragments, using well known techniques.

15 Alternatively, there are numerous amplification techniques for obtaining a full length coding sequence from a partial cDNA sequence. Within such techniques, amplification is generally performed via PCR. Any of a variety of commercially available kits may be used to perform the amplification step. Primers may be designed using, for example, software well known in the art. Primers are preferably 22-30  
20 nucleotides in length, have a GC content of at least 50% and anneal to the target sequence at temperatures of about 68°C to 72°C. The amplified region may be sequenced as described above, and overlapping sequences assembled into a contiguous sequence.

One such amplification technique is inverse PCR (see Triglia et al., *Nucl. Acids Res.* 16:8186, 1988), which uses restriction enzymes to generate a fragment in the  
25 known region of the gene. The fragment is then circularized by intramolecular ligation and used as a template for PCR with divergent primers derived from the known region. Within an alternative approach, sequences adjacent to a partial sequence may be retrieved by amplification with a primer to a linker sequence and a primer specific to a  
30 known region. The amplified sequences are typically subjected to a second round of amplification with the same linker primer and a second primer specific to the known

region. A variation on this procedure, which employs two primers that initiate extension in opposite directions from the known sequence, is described in WO 96/38591. Another such technique is known as "rapid amplification of cDNA ends" or RACE. This technique involves the use of an internal primer and an external primer, which hybridizes to a polyA region or vector sequence, to identify sequences that are 5' and 3' of a known sequence. Additional techniques include capture PCR (Lagerstrom et al., *PCR Methods Applic. 1*:111-19, 1991) and walking PCR (Parker et al., *Nucl. Acids. Res. 19*:3055-60, 1991). Other methods employing amplification may also be employed to obtain a full length cDNA sequence.

10 In certain instances, it is possible to obtain a full length cDNA sequence by analysis of sequences provided in an expressed sequence tag (EST) database, such as that available from GenBank. Searches for overlapping ESTs may generally be performed using well known programs (*e.g.*, NCBI BLAST searches), and such ESTs may be used to generate a contiguous full length sequence. Full length DNA sequences  
15 may also be obtained by analysis of genomic fragments.

#### POLYNUCLEOTIDE EXPRESSION IN HOST CELLS

In other embodiments of the invention, polynucleotide sequences or fragments thereof which encode polypeptides of the invention, or fusion proteins or functional equivalents thereof, may be used in recombinant DNA molecules to direct  
20 expression of a polypeptide in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences that encode substantially the same or a functionally equivalent amino acid sequence may be produced and these sequences may be used to clone and express a given polypeptide.

As will be understood by those of skill in the art, it may be advantageous  
25 in some instances to produce polypeptide-encoding nucleotide sequences possessing non-naturally occurring codons. For example, codons preferred by a particular prokaryotic or eukaryotic host can be selected to increase the rate of protein expression or to produce a recombinant RNA transcript having desirable properties, such as a half-life which is longer than that of a transcript generated from the naturally occurring  
30 sequence.

Moreover, the polynucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter polypeptide encoding sequences for a variety of reasons, including but not limited to, alterations which modify the cloning, processing, and/or expression of the gene product. For example, DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. In addition, site-directed mutagenesis may be used to insert new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, or introduce mutations, and so forth.

10 In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences may be ligated to a heterologous sequence to encode a fusion protein. For example, to screen peptide libraries for inhibitors of polypeptide activity, it may be useful to encode a chimeric protein that can be recognized by a commercially available antibody. A fusion protein may also be  
15 engineered to contain a cleavage site located between the polypeptide-encoding sequence and the heterologous protein sequence, so that the polypeptide may be cleaved and purified away from the heterologous moiety.

Sequences encoding a desired polypeptide may be synthesized, in whole or in part, using chemical methods well known in the art (see Caruthers, M. H. et al. (1980) *Nucl. Acids Res. Symp. Ser.* 215-223, Horn, T. et al. (1980) *Nucl. Acids Res. Symp. Ser.* 225-232). Alternatively, the protein itself may be produced using chemical methods to synthesize the amino acid sequence of a polypeptide, or a portion thereof. For example, peptide synthesis can be performed using various solid-phase techniques (Roberge, J. Y. et al. (1995) *Science* 269:202-204) and automated synthesis may be  
25 achieved, for example, using the ABI 431A Peptide Synthesizer (Perkin Elmer, Palo Alto, CA).

A newly synthesized peptide may be substantially purified by preparative high performance liquid chromatography (e.g., Creighton, T. (1983) *Proteins, Structures and Molecular Principles*, WH Freeman and Co., New York, N.Y.)  
30 or other comparable techniques available in the art. The composition of the synthetic peptides may be confirmed by amino acid analysis or sequencing (e.g., the Edman

degradation procedure). Additionally, the amino acid sequence of a polypeptide, or any part thereof, may be altered during direct synthesis and/or combined using chemical methods with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

5 In order to express a desired polypeptide, the nucleotide sequences encoding the polypeptide, or functional equivalents, may be inserted into appropriate expression vector, i.e., a vector which contains the necessary elements for the transcription and translation of the inserted coding sequence. Methods which are well known to those skilled in the art may be used to construct expression vectors containing  
10 sequences encoding a polypeptide of interest and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. Such techniques are described in Sambrook, J. et al. (1989) *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Press, Plainview, N.Y., and Ausubel, F. M. et al. (1989) *Current*  
15 *Protocols in Molecular Biology*, John Wiley & Sons, New York, N.Y.

A variety of expression vector/host systems may be utilized to contain and express polynucleotide sequences. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors;  
20 insect cell systems infected with virus expression vectors (e.g., baculovirus); plant cell systems transformed with virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems.

The "control elements" or "regulatory sequences" present in an  
25 expression vector are those non-translated regions of the vector--enhancers, promoters, 5' and 3' untranslated regions--which interact with host cellular proteins to carry out transcription and translation. Such elements may vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, may be used.  
30 For example, when cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the PBLUESCRIPT phagemid (Stratagene, La Jolla, Calif.) or

PSPORT1 plasmid (Gibco BRL, Gaithersburg, MD) and the like may be used. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are generally preferred. If it is necessary to generate a cell line that contains multiple copies of the sequence encoding a polypeptide, vectors based on SV40 or EBV  
5 may be advantageously used with an appropriate selectable marker.

In bacterial systems, a number of expression vectors may be selected depending upon the use intended for the expressed polypeptide. For example, when large quantities are needed, for example for the induction of antibodies, vectors which direct high level expression of fusion proteins that are readily purified may be used.  
10 Such vectors include, but are not limited to, the multifunctional *E. coli* cloning and expression vectors such as BLUESCRIPT (Stratagene), in which the sequence encoding the polypeptide of interest may be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of  $\beta$ -galactosidase so that a hybrid protein is produced; pIN vectors (Van Heeke, G. and S. M. Schuster (1989) *J.*  
15 *Biol. Chem.* 264:5503-5509); and the like. pGEX Vectors (Promega, Madison, Wis.) may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems may be designed to include  
20 heparin, thrombin, or factor XA protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

In the yeast, *Saccharomyces cerevisiae*, a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase, and PGH may be used. For reviews, see Ausubel et al. (supra) and Grant et al. (1987) *Methods*  
25 *Enzymol.* 153:516-544.

In cases where plant expression vectors are used, the expression of sequences encoding polypeptides may be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV may be used alone or in combination with the omega leader sequence from TMV (Takamatsu, N.  
30 (1987) *EMBO J.* 6:307-311. Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used (Coruzzi, G. et al. (1984) *EMBO J.*

3:1671-1680; Broglie, R. et al. (1984) *Science* 224:838-843; and Winter, J. et al. (1991) *Results Probl. Cell Differ.* 17:85-105). These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. Such techniques are described in a number of generally available reviews (see, for example, Hobbs, S. or  
5 Murry, L. E. in McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York, N.Y.; pp. 191-196).

An insect system may also be used to express a polypeptide of interest. For example, in one such system, Autographa californica nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda* cells or  
10 in *Trichoplusia* larvae. The sequences encoding the polypeptide may be cloned into a non-essential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of the polypeptide-encoding sequence will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses may then be used to infect, for example, *S. frugiperda*  
15 cells or *Trichoplusia* larvae in which the polypeptide of interest may be expressed (Engelhard, E. K. et al. (1994) *Proc. Natl. Acad. Sci.* 91 :3224-3227).

In mammalian host cells, a number of viral-based expression systems are generally available. For example, in cases where an adenovirus is used as an expression vector, sequences encoding a polypeptide of interest may be ligated into an adenovirus  
20 transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain a viable virus which is capable of expressing the polypeptide in infected host cells (Logan, J. and Shenk, T. (1984) *Proc. Natl. Acad. Sci.* 81:3655-3659). In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used  
25 to increase expression in mammalian host cells.

Specific initiation signals may also be used to achieve more efficient translation of sequences encoding a polypeptide of interest. Such signals include the ATG initiation codon and adjacent sequences. In cases where sequences encoding the polypeptide, its initiation codon, and upstream sequences are inserted into the  
30 appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a portion



thereof, is inserted, exogenous translational control signals including the ATG initiation codon should be provided. Furthermore, the initiation codon should be in the correct reading frame to ensure translation of the entire insert. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic.

- 5 The efficiency of expression may be enhanced by the inclusion of enhancers which are appropriate for the particular cell system which is used, such as those described in the literature (Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162).

In addition, a host cell strain may be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed protein in the  
10 desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" form of the protein may also be used to facilitate correct insertion, folding and/or function. Different host cells such as CHO, HeLa, MDCK, HEK293, and WI38, which have specific cellular machinery and  
15 characteristic mechanisms for such post-translational activities, may be chosen to ensure the correct modification and processing of the foreign protein.

For long-term, high-yield production of recombinant proteins, stable expression is generally preferred. For example, cell lines which stably express a polynucleotide of interest may be transformed using expression vectors which may  
20 contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for 1-2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which  
25 successfully express the introduced sequences. Resistant clones of stably transformed cells may be proliferated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase (Wigler, M. et al. (1977) *Cell* 11:223-32) and adenine phosphoribosyltransferase  
30 (Lowy, I. et al. (1990) *Cell* 22:817-23) genes which can be employed in tk.sup.- or aprt.sup.- cells, respectively. Also, antimetabolite, antibiotic or herbicide resistance can

be used as the basis for selection; for example, dhfr which confers resistance to methotrexate (Wigler, M. et al. (1980) *Proc. Natl. Acad. Sci.* 77:3567-70); npt, which confers resistance to the aminoglycosides, neomycin and G-418 (Colbere-Garapin, F. et al (1981) *J. Mol. Biol.* 150:1-14); and als or pat, which confer resistance to  
5 chlorsulfuron and phosphinotricin acetyltransferase, respectively (Murry, *supra*). Additional selectable genes have been described, for example, trpB, which allows cells to utilize indole in place of tryptophan, or hisD, which allows cells to utilize histinol in place of histidine (Hartman, S. C. and R. C. Mulligan (1988) *Proc. Natl. Acad. Sci.* 85:8047-51). Recently, the use of visible markers has gained popularity with such  
10 markers as anthocyanins, beta-glucuronidase and its substrate GUS, and luciferase and its substrate luciferin, being widely used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system (Rhodes, C. A. et al. (1995) *Methods Mol. Biol.* 55:121-131).

Although the presence/absence of marker gene expression suggests that  
15 the gene of interest is also present, its presence and expression may need to be confirmed. For example, if the sequence encoding a polypeptide is inserted within a marker gene sequence, recombinant cells containing sequences can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a polypeptide-encoding sequence under the control of a single promoter.  
20 Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

Alternatively, host cells which contain and express a desired polynucleotide sequence may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-  
25 RNA hybridizations and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein.

A variety of protocols for detecting and measuring the expression of polynucleotide-encoded products, using either polyclonal or monoclonal antibodies  
30 specific for the product are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), and fluorescence activated

cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on a given polypeptide may be preferred for some applications, but a competitive binding assay may also be employed. These and other assays are described, among other places, in Hampton, R. et al. (1990; 5 Serological Methods, a Laboratory Manual, APS Press, St Paul, Minn.) and Maddox, D. E. et al. (1983; *J. Exp. Med.* 158:1211-1216).

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to 10 polynucleotides include oligolabeling, nick translation, end-labeling or PCR amplification using a labeled nucleotide. Alternatively, the sequences, or any portions thereof may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 15 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits. Suitable reporter molecules or labels, which may be used include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with a polynucleotide sequence of interest may be 20 cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a recombinant cell may be secreted or contained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides of the invention may be designed to contain signal sequences which direct secretion of the 25 encoded polypeptide through a prokaryotic or eukaryotic cell membrane. Other recombinant constructions may be used to join sequences encoding a polypeptide of interest to nucleotide sequence encoding a polypeptide domain which will facilitate purification of soluble proteins. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow 30 purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity

purification system (Immunex Corp., Seattle, Wash.). The inclusion of cleavable linker sequences such as those specific for Factor XA or enterokinase (Invitrogen, San Diego, Calif.) between the purification domain and the encoded polypeptide may be used to facilitate purification. One such expression vector provides for expression of a fusion protein containing a polypeptide of interest and a nucleic acid encoding 6 histidine residues preceding a thioredoxin or an enterokinase cleavage site. The histidine residues facilitate purification on IMIAC (immobilized metal ion affinity chromatography) as described in Porath, J. et al. (1992, *Prot. Exp. Purif.* 3:263-281) while the enterokinase cleavage site provides a means for purifying the desired polypeptide from the fusion protein. A discussion of vectors which contain fusion proteins is provided in Kroll, D. J. et al. (1993; *DNA Cell Biol.* 12:441-453).

In addition to recombinant production methods, polypeptides of the invention, and fragments thereof, may be produced by direct peptide synthesis using solid-phase techniques (Merrifield J. (1963) *J. Am. Chem. Soc.* 85:2149-2154). Protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be achieved, for example, using Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer). Alternatively, various fragments may be chemically synthesized separately and combined using chemical methods to produce the full length molecule.

## 20 SITE-SPECIFIC MUTAGENESIS

Site-specific mutagenesis is a technique useful in the preparation of individual peptides, or biologically functional equivalent polypeptides, through specific mutagenesis of the underlying polynucleotides that encode them. The technique, well-known to those of skill in the art, further provides a ready ability to prepare and test sequence variants, for example, incorporating one or more of the foregoing considerations, by introducing one or more nucleotide sequence changes into the DNA. Site-specific mutagenesis allows the production of mutants through the use of specific oligonucleotide sequences which encode the DNA sequence of the desired mutation, as well as a sufficient number of adjacent nucleotides, to provide a primer sequence of sufficient size and sequence complexity to form a stable duplex on both sides of the

deletion junction being traversed. Mutations may be employed in a selected polynucleotide sequence to improve, alter, decrease, modify, or otherwise change the properties of the polynucleotide itself, and/or alter the properties, activity, composition, stability, or primary sequence of the encoded polypeptide.

5 In certain embodiments of the present invention, the inventors contemplate the mutagenesis of the disclosed polynucleotide sequences to alter one or more properties of the encoded polypeptide, such as the antigenicity of a polypeptide vaccine. The techniques of site-specific mutagenesis are well-known in the art, and are widely used to create variants of both polypeptides and polynucleotides. For example,  
10 site-specific mutagenesis is often used to alter a specific portion of a DNA molecule. In such embodiments, a primer comprising typically about 14 to about 25 nucleotides or so in length is employed, with about 5 to about 10 residues on both sides of the junction of the sequence being altered.

As will be appreciated by those of skill in the art, site-specific  
15 mutagenesis techniques have often employed a phage vector that exists in both a single stranded and double stranded form. Typical vectors useful in site-directed mutagenesis include vectors such as the M13 phage. These phage are readily commercially-available and their use is generally well-known to those skilled in the art. Double-stranded plasmids are also routinely employed in site directed mutagenesis that  
20 eliminates the step of transferring the gene of interest from a plasmid to a phage.

In general, site-directed mutagenesis in accordance herewith is performed by first obtaining a single-stranded vector or melting apart of two strands of a double-stranded vector that includes within its sequence a DNA sequence that encodes the desired peptide. An oligonucleotide primer bearing the desired mutated sequence is  
25 prepared, generally synthetically. This primer is then annealed with the single-stranded vector, and subjected to DNA polymerizing enzymes such as *E. coli* polymerase I Klenow fragment, in order to complete the synthesis of the mutation-bearing strand. Thus, a heteroduplex is formed wherein one strand encodes the original non-mutated sequence and the second strand bears the desired mutation. This heteroduplex vector is  
30 then used to transform appropriate cells, such as *E. coli* cells, and clones are selected which include recombinant vectors bearing the mutated sequence arrangement.

The preparation of sequence variants of the selected peptide-encoding DNA segments using site-directed mutagenesis provides a means of producing potentially useful species and is not meant to be limiting as there are other ways in which sequence variants of peptides and the DNA sequences encoding them may be  
5 obtained. For example, recombinant vectors encoding the desired peptide sequence may be treated with mutagenic agents, such as hydroxylamine, to obtain sequence variants. Specific details regarding these methods and protocols are found in the teachings of Maloy *et al.*, 1994; Segal, 1976; Prokop and Bajpai, 1991; Kuby, 1994; and Maniatis *et al.*, 1982, each incorporated herein by reference, for that purpose.

10 As used herein, the term "oligonucleotide directed mutagenesis procedure" refers to template-dependent processes and vector-mediated propagation which result in an increase in the concentration of a specific nucleic acid molecule relative to its initial concentration, or in an increase in the concentration of a detectable signal, such as amplification. As used herein, the term "oligonucleotide directed  
15 mutagenesis procedure" is intended to refer to a process that involves the template-dependent extension of a primer molecule. The term template dependent process refers to nucleic acid synthesis of an RNA or a DNA molecule wherein the sequence of the newly synthesized strand of nucleic acid is dictated by the well-known rules of complementary base pairing (see, for example, Watson, 1987). Typically,  
20 vector mediated methodologies involve the introduction of the nucleic acid fragment into a DNA or RNA vector, the clonal amplification of the vector, and the recovery of the amplified nucleic acid fragment. Examples of such methodologies are provided by U. S. Patent No. 4,237,224, specifically incorporated herein by reference in its entirety.

#### POLYNUCLEOTIDE AMPLIFICATION TECHNIQUES

25 A number of template dependent processes are available to amplify the target sequences of interest present in a sample. One of the best known amplification methods is the polymerase chain reaction (PCR™) which is described in detail in U.S. Patent Nos. 4,683,195, 4,683,202 and 4,800,159, each of which is incorporated herein by reference in its entirety. Briefly, in PCR™, two primer sequences are prepared  
30 which are complementary to regions on opposite complementary strands of the target

sequence. An excess of deoxynucleoside triphosphates is added to a reaction mixture along with a DNA polymerase (e.g., *Taq* polymerase). If the target sequence is present in a sample, the primers will bind to the target and the polymerase will cause the primers to be extended along the target sequence by adding on nucleotides. By raising  
5 and lowering the temperature of the reaction mixture, the extended primers will dissociate from the target to form reaction products, excess primers will bind to the target and to the reaction product and the process is repeated. Preferably reverse transcription and PCR™ amplification procedure may be performed in order to quantify the amount of mRNA amplified. Polymerase chain reaction methodologies are well  
10 known in the art.

Another method for amplification is the ligase chain reaction (referred to as LCR), disclosed in Eur. Pat. Appl. Publ. No. 320,308 (specifically incorporated herein by reference in its entirety). In LCR, two complementary probe pairs are prepared, and in the presence of the target sequence, each pair will bind to opposite  
15 complementary strands of the target such that they abut. In the presence of a ligase, the two probe pairs will link to form a single unit. By temperature cycling, as in PCR™, bound ligated units dissociate from the target and then serve as "target sequences" for ligation of excess probe pairs. U.S. Patent No. 4,883,750, incorporated herein by reference in its entirety, describes an alternative method of amplification similar to LCR  
20 for binding probe pairs to a target sequence.

Qbeta Replicase, described in PCT Intl. Pat. Appl. Publ. No. PCT/US87/00880, incorporated herein by reference in its entirety, may also be used as still another amplification method in the present invention. In this method, a replicative sequence of RNA that has a region complementary to that of a target is added to a  
25 sample in the presence of an RNA polymerase. The polymerase will copy the replicative sequence that can then be detected.

An isothermal amplification method, in which restriction endonucleases and ligases are used to achieve the amplification of target molecules that contain nucleotide 5'-[α-thio]triphosphates in one strand of a restriction site (Walker *et al.*,  
30 1992, incorporated herein by reference in its entirety), may also be useful in the amplification of nucleic acids in the present invention.

Strand Displacement Amplification (SDA) is another method of carrying out isothermal amplification of nucleic acids which involves multiple rounds of strand displacement and synthesis, *i.e.* nick translation. A similar method, called Repair Chain Reaction (RCR) is another method of amplification which may be useful in the present invention and is involves annealing several probes throughout a region targeted for amplification, followed by a repair reaction in which only two of the four bases are present. The other two bases can be added as biotinylated derivatives for easy detection. A similar approach is used in SDA.

Sequences can also be detected using a cyclic probe reaction (CPR). In CPR, a probe having a 3' and 5' sequences of non-target DNA and an internal or "middle" sequence of the target protein specific RNA is hybridized to DNA which is present in a sample. Upon hybridization, the reaction is treated with RNaseH, and the products of the probe are identified as distinctive products by generating a signal that is released after digestion. The original template is annealed to another cycling probe and the reaction is repeated. Thus, CPR involves amplifying a signal generated by hybridization of a probe to a target gene specific expressed nucleic acid.

Still other amplification methods described in Great Britain Pat. Appl. No. 2 202 328, and in PCT Intl. Pat. Appl. Publ. No. PCT/US89/01025, each of which is incorporated herein by reference in its entirety, may be used in accordance with the present invention. In the former application, "modified" primers are used in a PCR-like, template and enzyme dependent synthesis. The primers may be modified by labeling with a capture moiety (*e.g.*, biotin) and/or a detector moiety (*e.g.*, enzyme). In the latter application, an excess of labeled probes is added to a sample. In the presence of the target sequence, the probe binds and is cleaved catalytically. After cleavage, the target sequence is released intact to be bound by excess probe. Cleavage of the labeled probe signals the presence of the target sequence.

Other nucleic acid amplification procedures include transcription-based amplification systems (TAS) (Kwoh *et al.*, 1989; PCT Intl. Pat. Appl. Publ. No. WO 88/10315, incorporated herein by reference in its entirety), including nucleic acid sequence based amplification (NASBA) and 3SR. In NASBA, the nucleic acids can be prepared for amplification by standard phenol/chloroform extraction, heat denaturation



of a sample, treatment with lysis buffer and minispin columns for isolation of DNA and RNA or guanidinium chloride extraction of RNA. These amplification techniques involve annealing a primer that has sequences specific to the target sequence. Following polymerization, DNA/RNA hybrids are digested with RNase H while double stranded DNA molecules are heat-denatured again. In either case the single stranded DNA is made fully double stranded by addition of second target-specific primer, followed by polymerization. The double stranded DNA molecules are then multiply transcribed by a polymerase such as T7 or SP6. In an isothermal cyclic reaction, the RNAs are reverse transcribed into DNA, and transcribed once again with a polymerase such as T7 or SP6. The resulting products, whether truncated or complete, indicate target-specific sequences.

Eur. Pat. Appl. Publ. No. 329,822, incorporated herein by reference in its entirety, disclose a nucleic acid amplification process involving cyclically synthesizing single-stranded RNA ("ssRNA"), ssDNA, and double-stranded DNA (dsDNA), which may be used in accordance with the present invention. The ssRNA is a first template for a first primer oligonucleotide, which is elongated by reverse transcriptase (RNA-dependent DNA polymerase). The RNA is then removed from resulting DNA:RNA duplex by the action of ribonuclease H (RNase H, an RNase specific for RNA in a duplex with either DNA or RNA). The resultant ssDNA is a second template for a second primer, which also includes the sequences of an RNA polymerase promoter (exemplified by T7 RNA polymerase) 5' to its homology to its template. This primer is then extended by DNA polymerase (exemplified by the large "Klenow" fragment of *E. coli* DNA polymerase I), resulting as a double-stranded DNA ("dsDNA") molecule, having a sequence identical to that of the original RNA between the primers and having additionally, at one end, a promoter sequence. This promoter sequence can be used by the appropriate RNA polymerase to make many RNA copies of the DNA. These copies can then re-enter the cycle leading to very swift amplification. With proper choice of enzymes, this amplification can be done isothermally without addition of enzymes at each cycle. Because of the cyclical nature of this process, the starting sequence can be chosen to be in the form of either DNA or RNA.

PCT Intl. Pat. Appl. Publ. No. WO 89/06700, incorporated herein by reference in its entirety, disclose a nucleic acid sequence amplification scheme based on the hybridization of a promoter/primer sequence to a target single-stranded DNA ("ssDNA") followed by transcription of many RNA copies of the sequence. This  
5 scheme is not cyclic; i.e. new templates are not produced from the resultant RNA transcripts. Other amplification methods include "RACE" (Frohman, 1990), and "one-sided PCR" (Ohara, 1989) which are well-known to those of skill in the art.

Methods based on ligation of two (or more) oligonucleotides in the presence of nucleic acid having the sequence of the resulting "di-oligonucleotide",  
10 thereby amplifying the di-oligonucleotide (Wu and Dean, 1996, incorporated herein by reference in its entirety), may also be used in the amplification of DNA sequences of the present invention.

#### BIOLOGICAL FUNCTIONAL EQUIVALENTS

Modification and changes may be made in the structure of the  
15 polynucleotides and polypeptides of the present invention and still obtain a functional molecule that encodes a polypeptide with desirable characteristics. As mentioned above, it is often desirable to introduce one or more mutations into a specific polynucleotide sequence. In certain circumstances, the resulting encoded polypeptide sequence is altered by this mutation, or in other cases, the sequence of the polypeptide  
20 is unchanged by one or more mutations in the encoding polynucleotide.

When it is desirable to alter the amino acid sequence of a polypeptide to create an equivalent, or even an improved, second-generation molecule, the amino acid changes may be achieved by changing one or more of the codons of the encoding DNA sequence, according to Table 1.

25 For example, certain amino acids may be substituted for other amino acids in a protein structure without appreciable loss of interactive binding capacity with structures such as, for example, antigen-binding regions of antibodies or binding sites on substrate molecules. Since it is the interactive capacity and nature of a protein that defines that protein's biological functional activity, certain amino acid sequence  
30 substitutions can be made in a protein sequence, and, of course, its underlying DNA

coding sequence, and nevertheless obtain a protein with like properties. It is thus contemplated by the inventors that various changes may be made in the peptide sequences of the disclosed compositions, or corresponding DNA sequences which encode said peptides without appreciable loss of their biological utility or activity.

5

TABLE I

Amino Acids			Codons						
Alanine	Ala	A	GCA	GCC	GCG	GCU			
Cysteine	Cys	C	UGC	UGU					
Aspartic acid	Asp	D	GAC	GAU					
Glutamic acid	Glu	E	GAA	GAG					
Phenylalanine	Phe	F	UUC	UUU					
Glycine	Gly	G	GGA	GGC	GGG	GGU			
Histidine	His	H	CAC	CAU					
Isoleucine	Ile	I	AUA	AUC	AUU				
Lysine	Lys	K	AAA	AAG					
Leucine	Leu	L	UUA	UUG	CUA	CUC	CUG	CUU	
Methionine	Met	M	AUG						
Asparagine	Asn	N	AAC	AAU					
Proline	Pro	P	CCA	CCC	CCG	CCU			
Glutamine	Gln	Q	CAA	CAG					
Arginine	Arg	R	AGA	AGG	CGA	CGC	CGG	CGU	
Serine	Ser	S	AGC	AGU	UCA	UCC	UCG	UCU	
Threonine	Thr	T	ACA	ACC	ACG	ACU			
Valine	Val	V	GUA	GUC	GUG	GUU			
Tryptophan	Trp	W	UGG						
Tyrosine	Tyr	Y	UAC	UAU					

In making such changes, the hydropathic index of amino acids may be considered. The importance of the hydropathic amino acid index in conferring interactive biologic function on a protein is generally understood in the art (Kyte and  
 10 Doolittle, 1982, incorporated herein by reference). It is accepted that the relative

Doolittle, 1982, incorporated herein by reference). It is accepted that the relative hydrophobic character of the amino acid contributes to the secondary structure of the resultant protein, which in turn defines the interaction of the protein with other molecules, for example, enzymes, substrates, receptors, DNA, antibodies, antigens, and the like. Each amino acid has been assigned a hydrophobic index on the basis of its hydrophobicity and charge characteristics (Kyte and Doolittle, 1982). These values are: isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (-0.4); threonine (-0.7); serine (-0.8); tryptophan (-0.9); tyrosine (-1.3); proline (-1.6); histidine (-3.2); glutamate (-3.5); glutamine (-3.5); aspartate (-3.5); asparagine (-3.5); lysine (-3.9); and arginine (-4.5).

It is known in the art that certain amino acids may be substituted by other amino acids having a similar hydrophobic index or score and still result in a protein with similar biological activity, *i.e.* still obtain a biological functionally equivalent protein. In making such changes, the substitution of amino acids whose hydrophobic indices are within  $\pm 2$  is preferred, those within  $\pm 1$  are particularly preferred, and those within  $\pm 0.5$  are even more particularly preferred. It is also understood in the art that the substitution of like amino acids can be made effectively on the basis of hydrophobicity. U. S. Patent 4,554,101 (specifically incorporated herein by reference in its entirety), states that the greatest local average hydrophobicity of a protein, as governed by the hydrophobicity of its adjacent amino acids, correlates with a biological property of the protein.

As detailed in U. S. Patent 4,554,101, the following hydrophobicity values have been assigned to amino acid residues: arginine (+3.0); lysine (+3.0); aspartate (+3.0  $\pm$  1); glutamate (+3.0  $\pm$  1); serine (+0.3); asparagine (+0.2); glutamine (+0.2); glycine (0); threonine (-0.4); proline (-0.5  $\pm$  1); alanine (-0.5); histidine (-0.5); cysteine (-1.0); methionine (-1.3); valine (-1.5); leucine (-1.8); isoleucine (-1.8); tyrosine (-2.3); phenylalanine (-2.5); tryptophan (-3.4). It is understood that an amino acid can be substituted for another having a similar hydrophobicity value and still obtain a biologically equivalent, and in particular, an immunologically equivalent protein. In such changes, the substitution of amino acids whose hydrophobicity values are within  $\pm 2$

is preferred, those within  $\pm 1$  are particularly preferred, and those within  $\pm 0.5$  are even more particularly preferred.

As outlined above, amino acid substitutions are generally therefore based on the relative similarity of the amino acid side-chain substituents, for example, their hydrophobicity, hydrophilicity, charge, size, and the like. Exemplary substitutions that take various of the foregoing characteristics into consideration are well known to those of skill in the art and include: arginine and lysine; glutamate and aspartate; serine and threonine; glutamine and asparagine; and valine, leucine and isoleucine.

In addition, any polynucleotide may be further modified to increase stability *in vivo*. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends; the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages in the backbone; and/or the inclusion of nontraditional bases such as inosine, queosine and wybutosine, as well as acetyl-methyl-, thio- and other modified forms of adenine, cytidine, guanine, thymine and uridine.

#### IN VIVO POLYNUCLEOTIDE DELIVERY TECHNIQUES

In additional embodiments, genetic constructs comprising one or more of the polynucleotides of the invention are introduced into cells *in vivo*. This may be achieved using any of a variety of well known approaches, several of which are outlined below for the purpose of illustration.

##### 1. ADENOVIRUS

One of the preferred methods for *in vivo* delivery of one or more nucleic acid sequences involves the use of an adenovirus expression vector. "Adenovirus expression vector" is meant to include those constructs containing adenovirus sequences sufficient to (a) support packaging of the construct and (b) to express a polynucleotide that has been cloned therein in a sense or antisense orientation. Of course, in the context of an antisense construct, expression does not require that the gene product be synthesized.

The expression vector comprises a genetically engineered form of an adenovirus. Knowledge of the genetic organization of adenovirus, a 36 kb, linear, double-stranded DNA virus, allows substitution of large pieces of adenoviral DNA with foreign sequences up to 7 kb (Grunhaus and Horwitz, 1992). In contrast to retrovirus, the adenoviral infection of host cells does not result in chromosomal integration because adenoviral DNA can replicate in an episomal manner without potential genotoxicity. Also, adenoviruses are structurally stable, and no genome rearrangement has been detected after extensive amplification. Adenovirus can infect virtually all epithelial cells regardless of their cell cycle stage. So far, adenoviral infection appears to be linked only to mild disease such as acute respiratory disease in humans.

Adenovirus is particularly suitable for use as a gene transfer vector because of its mid-sized genome, ease of manipulation, high titer, wide target-cell range and high infectivity. Both ends of the viral genome contain 100-200 base pair inverted repeats (ITRs), which are *cis* elements necessary for viral DNA replication and packaging. The early (E) and late (L) regions of the genome contain different transcription units that are divided by the onset of viral DNA replication. The E1 region (E1A and E1B) encodes proteins responsible for the regulation of transcription of the viral genome and a few cellular genes. The expression of the E2 region (E2A and E2B) results in the synthesis of the proteins for viral DNA replication. These proteins are involved in DNA replication, late gene expression and host cell shut-off (Renan, 1990). The products of the late genes, including the majority of the viral capsid proteins, are expressed only after significant processing of a single primary transcript issued by the major late promoter (MLP). The MLP, (located at 16.8 m.u.) is particularly efficient during the late phase of infection, and all the mRNA's issued from this promoter possess a 5'-tripartite leader (TPL) sequence which makes them preferred mRNA's for translation.

In a current system, recombinant adenovirus is generated from homologous recombination between shuttle vector and provirus vector. Due to the possible recombination between two proviral vectors, wild-type adenovirus may be generated from this process. Therefore, it is critical to isolate a single clone of virus from an individual plaque and examine its genomic structure.

Generation and propagation of the current adenovirus vectors, which are replication deficient, depend on a unique helper cell line, designated 293, which was transformed from human embryonic kidney cells by Ad5 DNA fragments and constitutively expresses E1 proteins (Graham *et al.*, 1977). Since the E3 region is dispensable from the adenovirus genome (Jones and Shenk, 1978), the current adenovirus vectors, with the help of 293 cells, carry foreign DNA in either the E1, the D3 or both regions (Graham and Prevec, 1991). In nature, adenovirus can package approximately 105% of the wild-type genome (Ghosh-Choudhury *et al.*, 1987), providing capacity for about 2 extra kB of DNA. Combined with the approximately 5.5 kB of DNA that is replaceable in the E1 and E3 regions, the maximum capacity of the current adenovirus vector is under 7.5 kB, or about 15% of the total length of the vector. More than 80% of the adenovirus viral genome remains in the vector backbone and is the source of vector-borne cytotoxicity. Also, the replication deficiency of the E1-deleted virus is incomplete. For example, leakage of viral gene expression has been observed with the currently available vectors at high multiplicities of infection (MOI) (Mulligan, 1993).

Helper cell lines may be derived from human cells such as human embryonic kidney cells, muscle cells, hematopoietic cells or other human embryonic mesenchymal or epithelial cells. Alternatively, the helper cells may be derived from the cells of other mammalian species that are permissive for human adenovirus. Such cells include, *e.g.*, Vero cells or other monkey embryonic mesenchymal or epithelial cells. As stated above, the currently preferred helper cell line is 293.

Recently, Racher *et al.* (1995) disclosed improved methods for culturing 293 cells and propagating adenovirus. In one format, natural cell aggregates are grown by inoculating individual cells into 1 liter siliconized spinner flasks (Techne, Cambridge, UK) containing 100-200 ml of medium. Following stirring at 40 rpm, the cell viability is estimated with trypan blue. In another format, Fibra-Cel microcarriers (Bibby Sterlin, Stone, UK) (5 g/l) is employed as follows. A cell inoculum, resuspended in 5 ml of medium, is added to the carrier (50 ml) in a 250 ml Erlenmeyer flask and left stationary, with occasional agitation, for 1 to 4 h. The medium is then replaced with 50 ml of fresh medium and shaking initiated. For virus production, cells

are allowed to grow to about 80% confluence, after which time the medium is replaced (to 25% of the final volume) and adenovirus added at an MOI of 0.05. Cultures are left stationary overnight, following which the volume is increased to 100% and shaking commenced for another 72 h.

5                   Other than the requirement that the adenovirus vector be replication defective, or at least conditionally defective, the nature of the adenovirus vector is not believed to be crucial to the successful practice of the invention. The adenovirus may be of any of the 42 different known serotypes or subgroups A-F. Adenovirus type 5 of subgroup C is the preferred starting material in order to obtain a conditional replication-  
10   defective adenovirus vector for use in the present invention, since Adenovirus type 5 is a human adenovirus about which a great deal of biochemical and genetic information is known, and it has historically been used for most constructions employing adenovirus as a vector.

As stated above, the typical vector according to the present invention is  
15   replication defective and will not have an adenovirus E1 region. Thus, it will be most convenient to introduce the polynucleotide encoding the gene of interest at the position from which the E1-coding sequences have been removed. However, the position of insertion of the construct within the adenovirus sequences is not critical to the invention. The polynucleotide encoding the gene of interest may also be inserted in lieu  
20   of the deleted E3 region in E3 replacement vectors as described by Karlsson *et al.* (1986) or in the E4 region where a helper cell line or helper virus complements the E4 defect.

Adenovirus is easy to grow and manipulate and exhibits broad host range *in vitro* and *in vivo*. This group of viruses can be obtained in high titers, *e.g.*,  $10^9$ - $10^{11}$   
25   plaque-forming units per ml, and they are highly infective. The life cycle of adenovirus does not require integration into the host cell genome. The foreign genes delivered by adenovirus vectors are episomal and, therefore, have low genotoxicity to host cells. No side effects have been reported in studies of vaccination with wild-type adenovirus (Couch *et al.*, 1963; Top *et al.*, 1971), demonstrating their safety and therapeutic  
30   potential as *in vivo* gene transfer vectors.



Adenovirus vectors have been used in eukaryotic gene expression (Levrero *et al.*, 1991; Gomez-Foix *et al.*, 1992) and vaccine development (Grunhaus and Horwitz, 1992; Graham and Prevec, 1992). Recently, animal studies suggested that recombinant adenovirus could be used for gene therapy (Stratford-Perricaudet and Perricaudet, 1991; Stratford-Perricaudet *et al.*, 1990; Rich *et al.*, 1993). Studies in administering recombinant adenovirus to different tissues include trachea instillation (Rosenfeld *et al.*, 1991; Rosenfeld *et al.*, 1992), muscle injection (Ragot *et al.*, 1993), peripheral intravenous injections (Herz and Gerard, 1993) and stereotactic inoculation into the brain (Le Gal La Salle *et al.*, 1993).

## 10 2. RETROVIRUSES

The retroviruses are a group of single-stranded RNA viruses characterized by an ability to convert their RNA to double-stranded DNA in infected cells by a process of reverse-transcription (Coffin, 1990). The resulting DNA then stably integrates into cellular chromosomes as a provirus and directs synthesis of viral proteins. The integration results in the retention of the viral gene sequences in the recipient cell and its descendants. The retroviral genome contains three genes, gag, pol, and env that code for capsid proteins, polymerase enzyme, and envelope components, respectively. A sequence found upstream from the gag gene contains a signal for packaging of the genome into virions. Two long terminal repeat (LTR) sequences are present at the 5' and 3' ends of the viral genome. These contain strong promoter and enhancer sequences and are also required for integration in the host cell genome (Coffin, 1990).

In order to construct a retroviral vector, a nucleic acid encoding one or more oligonucleotide or polynucleotide sequences of interest is inserted into the viral genome in the place of certain viral sequences to produce a virus that is replication-defective. In order to produce virions, a packaging cell line containing the gag, pol, and env genes but without the LTR and packaging components is constructed (Mann *et al.*, 1983). When a recombinant plasmid containing a cDNA, together with the retroviral LTR and packaging sequences is introduced into this cell line (by calcium phosphate precipitation for example), the packaging sequence allows the RNA transcript of the

recombinant plasmid to be packaged into viral particles, which are then secreted into the culture media (Nicolas and Rubenstein, 1988; Temin, 1986; Mann *et al.*, 1983). The media containing the recombinant retroviruses is then collected, optionally concentrated, and used for gene transfer. Retroviral vectors are able to infect a broad  
5 variety of cell types. However, integration and stable expression require the division of host cells (Paskind *et al.*, 1975).

A novel approach designed to allow specific targeting of retrovirus vectors was recently developed based on the chemical modification of a retrovirus by the chemical addition of lactose residues to the viral envelope. This modification could  
10 permit the specific infection of hepatocytes *via* sialoglycoprotein receptors.

A different approach to targeting of recombinant retroviruses was designed in which biotinylated antibodies against a retroviral envelope protein and against a specific cell receptor were used. The antibodies were coupled *via* the biotin components by using streptavidin (Roux *et al.*, 1989). Using antibodies against major  
15 histocompatibility complex class I and class II antigens, they demonstrated the infection of a variety of human cells that bore those surface antigens with an ecotropic virus *in vitro* (Roux *et al.*, 1989).

### 3. ADENO-ASSOCIATED VIRUSES

AAV (Ridgeway, 1988; Hermonat and Muzyczka, 1984) is a parovirus,  
20 discovered as a contamination of adenoviral stocks. It is a ubiquitous virus (antibodies are present in 85% of the US human population) that has not been linked to any disease. It is also classified as a dependovirus, because its replications is dependent on the presence of a helper virus, such as adenovirus. Five serotypes have been isolated, of which AAV-2 is the best characterized. AAV has a single-stranded linear DNA that is  
25 encapsidated into capsid proteins VP1, VP2 and VP3 to form an icosahedral virion of 20 to 24 nm in diameter (Muzyczka and McLaughlin, 1988).

The AAV DNA is approximately 4.7 kilobases long. It contains two open reading frames and is flanked by two ITRs. There are two major genes in the AAV genome: *rep* and *cap*. The *rep* gene codes for proteins responsible for viral  
30 replications, whereas *cap* codes for capsid protein VP1-3. Each ITR forms a T-shaped

hairpin structure. These terminal repeats are the only essential *cis* components of the AAV for chromosomal integration. Therefore, the AAV can be used as a vector with all viral coding sequences removed and replaced by the cassette of genes for delivery. Three viral promoters have been identified and named p5, p19, and p40, according to their map position. Transcription from p5 and p19 results in production of rep proteins, and transcription from p40 produces the capsid proteins (Hermonat and Muzyczka, 1984).

There are several factors that prompted researchers to study the possibility of using rAAV as an expression vector. One is that the requirements for delivering a gene to integrate into the host chromosome are surprisingly few. It is necessary to have the 145-bp ITRs, which are only 6% of the AAV genome. This leaves room in the vector to assemble a 4.5-kb DNA insertion. While this carrying capacity may prevent the AAV from delivering large genes, it is amply suited for delivering the antisense constructs of the present invention.

AAV is also a good choice of delivery vehicles due to its safety. There is a relatively complicated rescue mechanism: not only wild type adenovirus but also AAV genes are required to mobilize rAAV. Likewise, AAV is not pathogenic and not associated with any disease. The removal of viral coding sequences minimizes immune reactions to viral gene expression, and therefore, rAAV does not evoke an inflammatory response.

#### 4. OTHER VIRAL VECTORS AS EXPRESSION CONSTRUCTS

Other viral vectors may be employed as expression constructs in the present invention for the delivery of oligonucleotide or polynucleotide sequences to a host cell. Vectors derived from viruses such as vaccinia virus (Ridgeway, 1988; Coupar *et al.*, 1988), lentiviruses, polio viruses and herpes viruses may be employed. They offer several attractive features for various mammalian cells (Friedmann, 1989; Ridgeway, 1988; Coupar *et al.*, 1988; Horwich *et al.*, 1990).

With the recent recognition of defective hepatitis B viruses, new insight was gained into the structure-function relationship of different viral sequences. *In vitro* studies showed that the virus could retain the ability for helper-dependent packaging

and reverse transcription despite the deletion of up to 80% of its genome (Horwich *et al.*, 1990). This suggested that large portions of the genome could be replaced with foreign genetic material. The hepatotropism and persistence (integration) were particularly attractive properties for liver-directed gene transfer. Chang *et al.* (1991) introduced the chloramphenicol acetyltransferase (CAT) gene into duck hepatitis B virus genome in the place of the polymerase, surface, and pre-surface coding sequences. It was cotransfected with wild-type virus into an avian hepatoma cell line. Culture media containing high titers of the recombinant virus were used to infect primary duckling hepatocytes. Stable CAT gene expression was detected for at least 24 days after transfection (Chang *et al.*, 1991).

#### 5. NON-VIRAL VECTORS

In order to effect expression of the oligonucleotide or polynucleotide sequences of the present invention, the expression construct must be delivered into a cell. This delivery may be accomplished *in vitro*, as in laboratory procedures for transforming cells lines, or *in vivo* or *ex vivo*, as in the treatment of certain disease states. As described above, one preferred mechanism for delivery is *via* viral infection where the expression construct is encapsulated in an infectious viral particle.

Once the expression construct has been delivered into the cell the nucleic acid encoding the desired oligonucleotide or polynucleotide sequences may be positioned and expressed at different sites. In certain embodiments, the nucleic acid encoding the construct may be stably integrated into the genome of the cell. This integration may be in the specific location and orientation *via* homologous recombination (gene replacement) or it may be integrated in a random, non-specific location (gene augmentation). In yet further embodiments, the nucleic acid may be stably maintained in the cell as a separate, episomal segment of DNA. Such nucleic acid segments or "episomes" encode sequences sufficient to permit maintenance and replication independent of or in synchronization with the host cell cycle. How the expression construct is delivered to a cell and where in the cell the nucleic acid remains is dependent on the type of expression construct employed.

In certain embodiments of the invention, the expression construct comprising one or more oligonucleotide or polynucleotide sequences may simply consist of naked recombinant DNA or plasmids. Transfer of the construct may be performed by any of the methods mentioned above which physically or chemically permeabilize the cell membrane. This is particularly applicable for transfer *in vitro* but it may be applied to *in vivo* use as well. Dubensky *et al.* (1984) successfully injected polyomavirus DNA in the form of calcium phosphate precipitates into liver and spleen of adult and newborn mice demonstrating active viral replication and acute infection. Benvenisty and Reshef (1986) also demonstrated that direct intraperitoneal injection of calcium phosphate-precipitated plasmids results in expression of the transfected genes. It is envisioned that DNA encoding a gene of interest may also be transferred in a similar manner *in vivo* and express the gene product.

Another embodiment of the invention for transferring a naked DNA expression construct into cells may involve particle bombardment. This method depends on the ability to accelerate DNA-coated microprojectiles to a high velocity allowing them to pierce cell membranes and enter cells without killing them (Klein *et al.*, 1987). Several devices for accelerating small particles have been developed. One such device relies on a high voltage discharge to generate an electrical current, which in turn provides the motive force (Yang *et al.*, 1990). The microprojectiles used have consisted of biologically inert substances such as tungsten or gold beads.

Selected organs including the liver, skin, and muscle tissue of rats and mice have been bombarded *in vivo* (Yang *et al.*, 1990; Zelenin *et al.*, 1991). This may require surgical exposure of the tissue or cells, to eliminate any intervening tissue between the gun and the target organ, *i.e. ex vivo* treatment. Again, DNA encoding a particular gene may be delivered *via* this method and still be incorporated by the present invention.

#### ANTISENSE OLIGONUCLEOTIDES

The end result of the flow of genetic information is the synthesis of protein. DNA is transcribed by polymerases into messenger RNA and translated on the ribosome to yield a folded, functional protein. Thus there are several steps along the

route where protein synthesis can be inhibited. The native DNA segment coding for a polypeptide described herein, as all such mammalian DNA strands, has two strands: a sense strand and an antisense strand held together by hydrogen bonding. The messenger RNA coding for polypeptide has the same nucleotide sequence as the sense  
5 DNA strand except that the DNA thymidine is replaced by uridine. Thus, synthetic antisense nucleotide sequences will bind to a mRNA and inhibit expression of the protein encoded by that mRNA.

The targeting of antisense oligonucleotides to mRNA is thus one mechanism to shut down protein synthesis, and, consequently, represents a powerful  
10 and targeted therapeutic approach. For example, the synthesis of polygalacturonase and the muscarine type 2 acetylcholine receptor are inhibited by antisense oligonucleotides directed to their respective mRNA sequences (U. S. Patent 5,739,119 and U. S. Patent 5,759,829, each specifically incorporated herein by reference in its entirety). Further, examples of antisense inhibition have been demonstrated with the  
15 nuclear protein cyclin, the multiple drug resistance gene (MDG1), ICAM-1, E-selectin, STK-1, striatal GABA<sub>A</sub> receptor and human EGF (Jaskulski *et al.*, 1988; Vasanthakumar and Ahmed, 1989; Peris *et al.*, 1998; U. S. Patent 5,801,154; U. S. Patent 5,789,573; U. S. Patent 5,718,709 and U. S. Patent 5,610,288, each specifically incorporated herein by reference in its entirety). Antisense constructs have also been  
20 described that inhibit and can be used to treat a variety of abnormal cellular proliferations, *e.g.* cancer (U. S. Patent 5,747,470; U. S. Patent 5,591,317 and U. S. Patent 5,783,683, each specifically incorporated herein by reference in its entirety).

Therefore, in exemplary embodiments, the invention provides oligonucleotide sequences that comprise all, or a portion of, any sequence that is  
25 capable of specifically binding to polynucleotide sequence described herein, or a complement thereof. In one embodiment, the antisense oligonucleotides comprise DNA or derivatives thereof. In another embodiment, the oligonucleotides comprise RNA or derivatives thereof. In a third embodiment, the oligonucleotides are modified DNAs comprising a phosphorothioated modified backbone. In a fourth embodiment, the  
30 oligonucleotide sequences comprise peptide nucleic acids or derivatives thereof. In each case, preferred compositions comprise a sequence region that is complementary,

and more preferably substantially-complementary, and even more preferably, completely complementary to one or more portions of polynucleotides disclosed herein.

Selection of antisense compositions specific for a given gene sequence is based upon analysis of the chosen target sequence (*i.e.* in these illustrative examples the  
5 rat and human sequences) and determination of secondary structure,  $T_m$ , binding energy, relative stability, and antisense compositions were selected based upon their relative inability to form dimers, hairpins, or other secondary structures that would reduce or prohibit specific binding to the target mRNA in a host cell.

Highly preferred target regions of the mRNA, are those which are at or  
10 near the AUG translation initiation codon, and those sequences which were substantially complementary to 5' regions of the mRNA. These secondary structure analyses and target site selection considerations were performed using v.4 of the OLIGO primer analysis software (Rychlik, 1997) and the BLASTN 2.0.5 algorithm software (Altschul *et al.*, 1997).

15 The use of an antisense delivery method employing a short peptide vector, termed MPG (27 residues), is also contemplated. The MPG peptide contains a hydrophobic domain derived from the fusion sequence of HIV gp41 and a hydrophilic domain from the nuclear localization sequence of SV40 T-antigen (Morris *et al.*, 1997). It has been demonstrated that several molecules of the MPG peptide coat the antisense  
20 oligonucleotides and can be delivered into cultured mammalian cells in less than 1 hour with relatively high efficiency (90%). Further, the interaction with MPG strongly increases both the stability of the oligonucleotide to nuclease and the ability to cross the plasma membrane (Morris *et al.*, 1997).

#### RIBOZYMES

25 Although proteins traditionally have been used for catalysis of nucleic acids, another class of macromolecules has emerged as useful in this endeavor. Ribozymes are RNA-protein complexes that cleave nucleic acids in a site-specific fashion. Ribozymes have specific catalytic domains that possess endonuclease activity (Kim and Cech, 1987; Gerlach *et al.*, 1987; Forster and Symons, 1987). For example, a  
30 large number of ribozymes accelerate phosphoester transfer reactions with a high degree

of specificity, often cleaving only one of several phosphoesters in an oligonucleotide substrate (Cech *et al.*, 1981; Michel and Westhof, 1990; Reinhold-Hurek and Shub, 1992). This specificity has been attributed to the requirement that the substrate bind via specific base-pairing interactions to the internal guide sequence ("IGS") of the ribozyme  
5 prior to chemical reaction.

Ribozyme catalysis has primarily been observed as part of sequence-specific cleavage/ligation reactions involving nucleic acids (Joyce, 1989; Cech *et al.*, 1981). For example, U. S. Patent No. 5,354,855 (specifically incorporated herein by reference) reports that certain ribozymes can act as endonucleases with a sequence  
10 specificity greater than that of known ribonucleases and approaching that of the DNA restriction enzymes. Thus, sequence-specific ribozyme-mediated inhibition of gene expression may be particularly suited to therapeutic applications (Scanlon *et al.*, 1991; Sarver *et al.*, 1990). Recently, it was reported that ribozymes elicited genetic changes in some cells lines to which they were applied; the altered genes included the oncogenes  
15 H-*ras*, c-*fos* and genes of HIV. Most of this work involved the modification of a target mRNA, based on a specific mutant codon that is cleaved by a specific ribozyme.

Six basic varieties of naturally-occurring enzymatic RNAs are known presently. Each can catalyze the hydrolysis of RNA phosphodiester bonds *in trans* (and thus can cleave other RNA molecules) under physiological conditions. In general,  
20 enzymatic nucleic acids act by first binding to a target RNA. Such binding occurs through the target binding portion of a enzymatic nucleic acid which is held in close proximity to an enzymatic portion of the molecule that acts to cleave the target RNA. Thus, the enzymatic nucleic acid first recognizes and then binds a target RNA through complementary base-pairing, and once bound to the correct site, acts enzymatically to  
25 cut the target RNA. Strategic cleavage of such a target RNA will destroy its ability to direct synthesis of an encoded protein. After an enzymatic nucleic acid has bound and cleaved its RNA target, it is released from that RNA to search for another target and can repeatedly bind and cleave new targets.

The enzymatic nature of a ribozyme is advantageous over many  
30 technologies, such as antisense technology (where a nucleic acid molecule simply binds to a nucleic acid target to block its translation) since the concentration of ribozyme



necessary to affect a therapeutic treatment is lower than that of an antisense oligonucleotide. This advantage reflects the ability of the ribozyme to act enzymatically. Thus, a single ribozyme molecule is able to cleave many molecules of target RNA. In addition, the ribozyme is a highly specific inhibitor, with the specificity  
5 of inhibition depending not only on the base pairing mechanism of binding to the target RNA, but also on the mechanism of target RNA cleavage. Single mismatches, or base-substitutions, near the site of cleavage can completely eliminate catalytic activity of a ribozyme. Similar mismatches in antisense molecules do not prevent their action (Woolf *et al.*, 1992). Thus, the specificity of action of a ribozyme is greater than that of  
10 an antisense oligonucleotide binding the same RNA site.

The enzymatic nucleic acid molecule may be formed in a hammerhead, hairpin, a hepatitis  $\delta$  virus, group I intron or RNaseP RNA (in association with an RNA guide sequence) or Neurospora VS RNA motif. Examples of hammerhead motifs are described by Rossi *et al.* (1992). Examples of hairpin motifs are described by Hampel  
15 *et al.* (Eur. Pat. Appl. Publ. No. EP 0360257), Hampel and Tritz (1989), Hampel *et al.* (1990) and U. S. Patent 5,631,359 (specifically incorporated herein by reference). An example of the hepatitis  $\delta$  virus motif is described by Perrotta and Been (1992); an example of the RNaseP motif is described by Guerrier-Takada *et al.* (1983); Neurospora VS RNA ribozyme motif is described by Collins (Saville and Collins, 1990; Saville and  
20 Collins, 1991; Collins and Olive, 1993); and an example of the Group I intron is described in (U. S. Patent 4,987,071, specifically incorporated herein by reference). All that is important in an enzymatic nucleic acid molecule of this invention is that it has a specific substrate binding site which is complementary to one or more of the target gene RNA regions, and that it have nucleotide sequences within or surrounding that substrate  
25 binding site which impart an RNA cleaving activity to the molecule. Thus the ribozyme constructs need not be limited to specific motifs mentioned herein.

In certain embodiments, it may be important to produce enzymatic cleaving agents which exhibit a high degree of specificity for the RNA of a desired target, such as one of the sequences disclosed herein. The enzymatic nucleic acid  
30 molecule is preferably targeted to a highly conserved sequence region of a target mRNA. Such enzymatic nucleic acid molecules can be delivered exogenously to

specific cells as required. Alternatively, the ribozymes can be expressed from DNA or RNA vectors that are delivered to specific cells.

Small enzymatic nucleic acid motifs (*e.g.*, of the hammerhead or the hairpin structure) may also be used for exogenous delivery. The simple structure of these molecules increases the ability of the enzymatic nucleic acid to invade targeted regions of the mRNA structure. Alternatively, catalytic RNA molecules can be expressed within cells from eukaryotic promoters (*e.g.*, Scanlon *et al.*, 1991; Kashani-Sabet *et al.*, 1992; Dropulic *et al.*, 1992; Weerasinghe *et al.*, 1991; Ojwang *et al.*, 1992; Chen *et al.*, 1992; Sarver *et al.*, 1990). Those skilled in the art realize that any ribozyme can be expressed in eukaryotic cells from the appropriate DNA vector. The activity of such ribozymes can be augmented by their release from the primary transcript by a second ribozyme (Int. Pat. Appl. Publ. No. WO 93/23569, and Int. Pat. Appl. Publ. No. WO 94/02595, both hereby incorporated by reference; Ohkawa *et al.*, 1992; Taira *et al.*, 1991; and Ventura *et al.*, 1993).

Ribozymes may be added directly, or can be complexed with cationic lipids, lipid complexes, packaged within liposomes, or otherwise delivered to target cells. The RNA or RNA complexes can be locally administered to relevant tissues *ex vivo*, or *in vivo* through injection, aerosol inhalation, infusion pump or stent, with or without their incorporation in biopolymers.

Ribozymes may be designed as described in Int. Pat. Appl. Publ. No. WO 93/23569 and Int. Pat. Appl. Publ. No. WO 94/02595, each specifically incorporated herein by reference) and synthesized to be tested *in vitro* and *in vivo*, as described. Such ribozymes can also be optimized for delivery. While specific examples are provided, those in the art will recognize that equivalent RNA targets in other species can be utilized when necessary.

Hammerhead or hairpin ribozymes may be individually analyzed by computer folding (Jaeger *et al.*, 1989) to assess whether the ribozyme sequences fold into the appropriate secondary structure. Those ribozymes with unfavorable intramolecular interactions between the binding arms and the catalytic core are eliminated from consideration. Varying binding arm lengths can be chosen to optimize

activity. Generally, at least 5 or so bases on each arm are able to bind to, or otherwise interact with, the target RNA.

Ribozymes of the hammerhead or hairpin motif may be designed to anneal to various sites in the mRNA message, and can be chemically synthesized. The method of synthesis used follows the procedure for normal RNA synthesis as described in Usman *et al.* (1987) and in Scaringe *et al.* (1990) and makes use of common nucleic acid protecting and coupling groups, such as dimethoxytrityl at the 5'-end, and phosphoramidites at the 3'-end. Average stepwise coupling yields are typically >98%. Hairpin ribozymes may be synthesized in two parts and annealed to reconstruct an active ribozyme (Chowrira and Burke, 1992). Ribozymes may be modified extensively to enhance stability by modification with nuclease resistant groups, for example, 2'-amino, 2'-C-allyl, 2'-fluoro, 2'-o-methyl, 2'-H (for a review see *e.g.*, Usman and Cedergren, 1992). Ribozymes may be purified by gel electrophoresis using general methods or by high pressure liquid chromatography and resuspended in water.

Ribozyme activity can be optimized by altering the length of the ribozyme binding arms, or chemically synthesizing ribozymes with modifications that prevent their degradation by serum ribonucleases (see *e.g.*, Int. Pat. Appl. Publ. No. WO 92/07065; Perrault *et al.*, 1990; Pieken *et al.*, 1991; Usman and Cedergren, 1992; Int. Pat. Appl. Publ. No. WO 93/15187; Int. Pat. Appl. Publ. No. WO 91/03162; Eur. Pat. Appl. Publ. No. 92110298.4; U. S. Patent 5,334,711; and Int. Pat. Appl. Publ. No. WO 94/13688, which describe various chemical modifications that can be made to the sugar moieties of enzymatic RNA molecules), modifications which enhance their efficacy in cells, and removal of stem II bases to shorten RNA synthesis times and reduce chemical requirements.

Sullivan *et al.* (Int. Pat. Appl. Publ. No. WO 94/02595) describes the general methods for delivery of enzymatic RNA molecules. Ribozymes may be administered to cells by a variety of methods known to those familiar to the art, including, but not restricted to, encapsulation in liposomes, by iontophoresis, or by incorporation into other vehicles, such as hydrogels, cyclodextrins, biodegradable nanocapsules, and bioadhesive microspheres. For some indications, ribozymes may be directly delivered *ex vivo* to cells or tissues with or without the aforementioned vehicles.

Alternatively, the RNA/vehicle combination may be locally delivered by direct inhalation, by direct injection or by use of a catheter, infusion pump or stent. Other routes of delivery include, but are not limited to, intravascular, intramuscular, subcutaneous or joint injection, aerosol inhalation, oral (tablet or pill form), topical, systemic, ocular, intraperitoneal and/or intrathecal delivery. More detailed descriptions of ribozyme delivery and administration are provided in Int. Pat. Appl. Publ. No. WO 94/02595 and Int. Pat. Appl. Publ. No. WO 93/23569, each specifically incorporated herein by reference.

Another means of accumulating high concentrations of a ribozyme(s) within cells is to incorporate the ribozyme-encoding sequences into a DNA expression vector. Transcription of the ribozyme sequences are driven from a promoter for eukaryotic RNA polymerase I (pol I), RNA polymerase II (pol II), or RNA polymerase III (pol III). Transcripts from pol II or pol III promoters will be expressed at high levels in all cells; the levels of a given pol II promoter in a given cell type will depend on the nature of the gene regulatory sequences (enhancers, silencers, *etc.*) present nearby. Prokaryotic RNA polymerase promoters may also be used, providing that the prokaryotic RNA polymerase enzyme is expressed in the appropriate cells (Elroy-Stein and Moss, 1990; Gao and Huang, 1993; Lieber *et al.*, 1993; Zhou *et al.*, 1990). Ribozymes expressed from such promoters can function in mammalian cells (*e.g.* Kashani-Saber *et al.*, 1992; Ojwang *et al.*, 1992; Chen *et al.*, 1992; Yu *et al.*, 1993; L'Huillier *et al.*, 1992; Lisziewicz *et al.*, 1993). Such transcription units can be incorporated into a variety of vectors for introduction into mammalian cells, including but not restricted to, plasmid DNA vectors, viral DNA vectors (such as adenovirus or adeno-associated vectors), or viral RNA vectors (such as retroviral, semliki forest virus, sindbis virus vectors).

Ribozymes may be used as diagnostic tools to examine genetic drift and mutations within diseased cells. They can also be used to assess levels of the target RNA molecule. The close relationship between ribozyme activity and the structure of the target RNA allows the detection of mutations in any region of the molecule which alters the base-pairing and three-dimensional structure of the target RNA. By using multiple ribozymes, one may map nucleotide changes which are important to RNA

structure and function *in vitro*, as well as in cells and tissues. Cleavage of target RNAs with ribozymes may be used to inhibit gene expression and define the role (essentially) of specified gene products in the progression of disease. In this manner, other genetic targets may be defined as important mediators of the disease. These studies will lead to better treatment of the disease progression by affording the possibility of combinational therapies (e.g., multiple ribozymes targeted to different genes, ribozymes coupled with known small molecule inhibitors, or intermittent treatment with combinations of ribozymes and/or other chemical or biological molecules). Other *in vitro* uses of ribozymes are well known in the art, and include detection of the presence of mRNA associated with an IL-5 related condition. Such RNA is detected by determining the presence of a cleavage product after treatment with a ribozyme using standard methodology.

#### PEPTIDE NUCLEIC ACIDS

In certain embodiments, the inventors contemplate the use of peptide nucleic acids (PNAs) in the practice of the methods of the invention. PNA is a DNA mimic in which the nucleobases are attached to a pseudopeptide backbone (Good and Nielsen, 1997). PNA is able to be utilized in a number methods that traditionally have used RNA or DNA. Often PNA sequences perform better in techniques than the corresponding RNA or DNA sequences and have utilities that are not inherent to RNA or DNA. A review of PNA including methods of making, characteristics of, and methods of using, is provided by Corey (1997) and is incorporated herein by reference. As such, in certain embodiments, one may prepare PNA sequences that are complementary to one or more portions of the ACE mRNA sequence, and such PNA compositions may be used to regulate, alter, decrease, or reduce the translation of ACE-specific mRNA, and thereby alter the level of ACE activity in a host cell to which such PNA compositions have been administered.

PNAs have 2-aminoethyl-glycine linkages replacing the normal phosphodiester backbone of DNA (Nielsen *et al.*, 1991; Hanvey *et al.*, 1992; Hyrup and Nielsen, 1996; Neilsen, 1996). This chemistry has three important consequences: firstly, in contrast to DNA or phosphorothioate oligonucleotides, PNAs are neutral

molecules; secondly, PNAs are achiral, which avoids the need to develop a stereoselective synthesis; and thirdly, PNA synthesis uses standard Boc (Dueholm *et al.*, 1994) or Fmoc (Thomson *et al.*, 1995) protocols for solid-phase peptide synthesis, although other methods, including a modified Merrifield method, have been used  
5 (Christensen *et al.*, 1995).

PNA monomers or ready-made oligomers are commercially available from PerSeptive Biosystems (Framingham, MA). PNA syntheses by either Boc or Fmoc protocols are straightforward using manual or automated protocols (Norton *et al.*, 1995). The manual protocol lends itself to the production of chemically modified PNAs  
10 or the simultaneous synthesis of families of closely related PNAs.

As with peptide synthesis, the success of a particular PNA synthesis will depend on the properties of the chosen sequence. For example, while in theory PNAs can incorporate any combination of nucleotide bases, the presence of adjacent purines can lead to deletions of one or more residues in the product. In expectation of this  
15 difficulty, it is suggested that, in producing PNAs with adjacent purines, one should repeat the coupling of residues likely to be added inefficiently. This should be followed by the purification of PNAs by reverse-phase high-pressure liquid chromatography (Norton *et al.*, 1995) providing yields and purity of product similar to those observed during the synthesis of peptides.

20 Modifications of PNAs for a given application may be accomplished by coupling amino acids during solid-phase synthesis or by attaching compounds that contain a carboxylic acid group to the exposed N-terminal amine. Alternatively, PNAs can be modified after synthesis by coupling to an introduced lysine or cysteine. The ease with which PNAs can be modified facilitates optimization for better solubility or  
25 for specific functional requirements. Once synthesized, the identity of PNAs and their derivatives can be confirmed by mass spectrometry. Several studies have made and utilized modifications of PNAs (Norton *et al.*, 1995; Haaima *et al.*, 1996; Stetsenko *et al.*, 1996; Petersen *et al.*, 1995; Ulmann *et al.*, 1996; Koch *et al.*, 1995; Orum *et al.*, 1995; Footer *et al.*, 1996; Griffith *et al.*, 1995; Kremsky *et al.*, 1996; Pardridge *et al.*,  
30 1995; Boffa *et al.*, 1995; Landsdorp *et al.*, 1996; Gambacorti-Passerini *et al.*, 1996; Armitage *et al.*, 1997; Seeger *et al.*, 1997; Ruskowski *et al.*, 1997). U.S. Patent No.

5,700,922 discusses PNA-DNA-PNA chimeric molecules and their uses in diagnostics, modulating protein in organisms, and treatment of conditions susceptible to therapeutics.

In contrast to DNA and RNA, which contain negatively charged linkages, the PNA backbone is neutral. In spite of this dramatic alteration, PNAs recognize complementary DNA and RNA by Watson-Crick pairing (Egholm *et al.*, 1993), validating the initial modeling by Nielsen *et al.* (1991). PNAs lack 3' to 5' polarity and can bind in either parallel or antiparallel fashion, with the antiparallel mode being preferred (Egholm *et al.*, 1993).

Hybridization of DNA oligonucleotides to DNA and RNA is destabilized by electrostatic repulsion between the negatively charged phosphate backbones of the complementary strands. By contrast, the absence of charge repulsion in PNA-DNA or PNA-RNA duplexes increases the melting temperature ( $T_m$ ) and reduces the dependence of  $T_m$  on the concentration of mono- or divalent cations (Nielsen *et al.*, 1991). The enhanced rate and affinity of hybridization are significant because they are responsible for the surprising ability of PNAs to perform strand invasion of complementary sequences within relaxed double-stranded DNA. In addition, the efficient hybridization at inverted repeats suggests that PNAs can recognize secondary structure effectively within double-stranded DNA. Enhanced recognition also occurs with PNAs immobilized on surfaces, and Wang *et al.* have shown that support-bound PNAs can be used to detect hybridization events (Wang *et al.*, 1996).

One might expect that tight binding of PNAs to complementary sequences would also increase binding to similar (but not identical) sequences, reducing the sequence specificity of PNA recognition. As with DNA hybridization, however, selective recognition can be achieved by balancing oligomer length and incubation temperature. Moreover, selective hybridization of PNAs is encouraged by PNA-DNA hybridization being less tolerant of base mismatches than DNA-DNA hybridization. For example, a single mismatch within a 16 bp PNA-DNA duplex can reduce the  $T_m$  by up to 15°C (Egholm *et al.*, 1993). This high level of discrimination has allowed the development of several PNA-based strategies for the analysis of point mutations (Wang

*et al.*, 1996; Carlsson *et al.*, 1996; Thiede *et al.*, 1996; Webb and Hurskainen, 1996; Perry-O'Keefe *et al.*, 1996).

High-affinity binding provides clear advantages for molecular recognition and the development of new applications for PNAs. For example, 11-13 nucleotide PNAs inhibit the activity of telomerase, a ribonucleo-protein that extends telomere ends using an essential RNA template, while the analogous DNA oligomers do not (Norton *et al.*, 1996).

Neutral PNAs are more hydrophobic than analogous DNA oligomers, and this can lead to difficulty solubilizing them at neutral pH, especially if the PNAs have a high purine content or if they have the potential to form secondary structures. Their solubility can be enhanced by attaching one or more positive charges to the PNA termini (Nielsen *et al.*, 1991).

Findings by Allfrey and colleagues suggest that strand invasion will occur spontaneously at sequences within chromosomal DNA (Boffa *et al.*, 1995; Boffa *et al.*, 1996). These studies targeted PNAs to triplet repeats of the nucleotides CAG and used this recognition to purify transcriptionally active DNA (Boffa *et al.*, 1995) and to inhibit transcription (Boffa *et al.*, 1996). This result suggests that if PNAs can be delivered within cells then they will have the potential to be general sequence-specific regulators of gene expression. Studies and reviews concerning the use of PNAs as antisense and anti-gene agents include Nielsen *et al.* (1993b), Hanvey *et al.* (1992), and Good and Nielsen (1997). Koppelhus *et al.* (1997) have used PNAs to inhibit HIV-1 inverse transcription, showing that PNAs may be used for antiviral therapies.

Methods of characterizing the antisense binding properties of PNAs are discussed in Rose (1993) and Jensen *et al.* (1997). Rose uses capillary gel electrophoresis to determine binding of PNAs to their complementary oligonucleotide, measuring the relative binding kinetics and stoichiometry. Similar types of measurements were made by Jensen *et al.* using BIAcore™ technology.

Other applications of PNAs include use in DNA strand invasion (Nielsen *et al.*, 1991), antisense inhibition (Hanvey *et al.*, 1992), mutational analysis (Orum *et al.*, 1993), enhancers of transcription (Mollegaard *et al.*, 1994), nucleic acid purification (Orum *et al.*, 1995), isolation of transcriptionally active genes (Boffa *et al.*, 1995),



blocking of transcription factor binding (Vickers *et al.*, 1995), genome cleavage (Veselkov *et al.*, 1996), biosensors (Wang *et al.*, 1996), *in situ* hybridization (Thisted *et al.*, 1996), and in an alternative to Southern blotting (Perry-O'Keefe, 1996).

#### POLYPEPTIDE COMPOSITIONS

5           The present invention, in other aspects, provides polypeptide compositions. Generally, a polypeptide of the invention will be an isolated polypeptide (or an epitope, variant, or active fragment thereof) derived from a mammalian species. Preferably, the polypeptide is encoded by a polynucleotide sequence disclosed herein or a sequence which hybridizes under moderately stringent conditions to a polynucleotide  
10           sequence disclosed herein. Alternatively, the polypeptide may be defined as a polypeptide which comprises a contiguous amino acid sequence from an amino acid sequence disclosed herein, or which polypeptide comprises an entire amino acid sequence disclosed herein.

          In the present invention, a polypeptide composition is also understood to  
15           comprise one or more polypeptides that are immunologically reactive with antibodies generated against a polypeptide of the invention, or to active fragments, or to variants or biological functional equivalents thereof.

          Likewise, a polypeptide composition of the present invention is understood to comprise one or more polypeptides that are capable of eliciting antibodies  
20           that are immunologically reactive with one or more polypeptides encoded by one or more contiguous nucleic acid sequences contained in SEQ ID NO: 1-38, 42-204, 205, 207 and 210-290, or to active fragments, or to variants thereof, or to one or more nucleic acid sequences which hybridize to one or more of these sequences under conditions of moderate to high stringency.

25           As used herein, an active fragment of a polypeptide includes a whole or a portion of a polypeptide which is modified by conventional techniques, *e.g.*, mutagenesis, or by addition, deletion, or substitution, but which active fragment exhibits substantially the same structure function, antigenicity, etc., as a polypeptide as described herein.

In certain illustrative embodiments, the polypeptides of the invention will comprise at least an immunogenic portion of a breast tumor protein or a variant thereof, as described herein. As noted above, a "breast tumor protein" is a protein that is expressed by breast tumor cells. Proteins that are breast tumor proteins react  
5 detectably within an immunoassay (such as an ELISA) with antisera from a patient with breast cancer. Polypeptides as described herein may be of any length. Additional sequences derived from the native protein and/or heterologous sequences may be present, and such sequences may (but need not) possess further immunogenic or antigenic properties.

10 An "immunogenic portion," as used herein is a portion of a protein that is recognized (*i.e.*, specifically bound) by a B-cell and/or T-cell surface antigen receptor. Such immunogenic portions generally comprise at least 5 amino acid residues, more preferably at least 10, and still more preferably at least 20 amino acid residues of a breast tumor protein or a variant thereof. Certain preferred immunogenic  
15 portions include peptides in which an N-terminal leader sequence and/or transmembrane domain have been deleted. Other preferred immunogenic portions may contain a small N- and/or C-terminal deletion (*e.g.*, 1-30 amino acids, preferably 5-15 amino acids), relative to the mature protein.

Immunogenic portions may generally be identified using well known  
20 techniques, such as those summarized in Paul, *Fundamental Immunology*, 3rd ed., 243-247 (Raven Press, 1993) and references cited therein. Such techniques include screening polypeptides for the ability to react with antigen-specific antibodies, antisera and/or T-cell lines or clones. As used herein, antisera and antibodies are "antigen-specific" if they specifically bind to an antigen (*i.e.*, they react with the protein in an  
25 ELISA or other immunoassay, and do not react detectably with unrelated proteins). Such antisera and antibodies may be prepared as described herein, and using well known techniques. An immunogenic portion of a native breast tumor protein is a portion that reacts with such antisera and/or T-cells at a level that is not substantially less than the reactivity of the full length polypeptide (*e.g.*, in an ELISA and/or T-cell  
30 reactivity assay). Such immunogenic portions may react within such assays at a level that is similar to or greater than the reactivity of the full length polypeptide. Such

5 screens may generally be performed using methods well known to those of ordinary skill in the art, such as those described in Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. For example, a polypeptide may be immobilized on a solid support and contacted with patient sera to allow binding of antibodies within the sera to the immobilized polypeptide. Unbound sera may then be removed and bound antibodies detected using, for example, <sup>125</sup>I-labeled Protein A.

As noted above, a composition may comprise a variant of a native breast tumor protein. A polypeptide "variant," as used herein, is a polypeptide that differs from a native breast tumor protein in one or more substitutions, deletions, additions and/or insertions, such that the immunogenicity of the polypeptide is not substantially diminished. In other words, the ability of a variant to react with antigen-specific antisera may be enhanced or unchanged, relative to the native protein, or may be diminished by less than 50%, and preferably less than 20%, relative to the native protein. Such variants may generally be identified by modifying one of the above polypeptide sequences and evaluating the reactivity of the modified polypeptide with antigen-specific antibodies or antisera as described herein. Preferred variants include those in which one or more portions, such as an N-terminal leader sequence or transmembrane domain, have been removed. Other preferred variants include variants in which a small portion (e.g., 1-30 amino acids, preferably 5-15 amino acids) has been removed from the N- and/or C-terminal of the mature protein.

Polypeptide variants encompassed by the present invention include those exhibiting at least about 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% or more identity (determined as described above) to the polypeptides disclosed herein.

25 Preferably, a variant contains conservative substitutions. A "conservative substitution" is one in which an amino acid is substituted for another amino acid that has similar properties, such that one skilled in the art of peptide chemistry would expect the secondary structure and hydropathic nature of the polypeptide to be substantially unchanged. Amino acid substitutions may generally be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity and/or the amphipathic nature of the residues. For example, negatively

charged amino acids include aspartic acid and glutamic acid; positively charged amino acids include lysine and arginine; and amino acids with uncharged polar head groups having similar hydrophilicity values include leucine, isoleucine and valine; glycine and alanine; asparagine and glutamine; and serine, threonine, phenylalanine and tyrosine.

- 5 Other groups of amino acids that may represent conservative changes include: (1) ala, pro, gly, glu, asp, gln, asn, ser, thr; (2) cys, ser, tyr, thr; (3) val, ile, leu, met, ala, phe; (4) lys, arg, his; and (5) phe, tyr, trp, his. A variant may also, or alternatively, contain nonconservative changes. In a preferred embodiment, variant polypeptides differ from a native sequence by substitution, deletion or addition of five amino acids or fewer.
- 10 Variants may also (or alternatively) be modified by, for example, the deletion or addition of amino acids that have minimal influence on the immunogenicity, secondary structure and hydrophobic nature of the polypeptide.

As noted above, polypeptides may comprise a signal (or leader) sequence at the N-terminal end of the protein, which co-translationally or post-translationally directs transfer of the protein. The polypeptide may also be conjugated to a linker or other sequence for ease of synthesis, purification or identification of the polypeptide (*e.g.*, poly-His), or to enhance binding of the polypeptide to a solid support. For example, a polypeptide may be conjugated to an immunoglobulin Fc region.

15

Polypeptides may be prepared using any of a variety of well known techniques. Recombinant polypeptides encoded by DNA sequences as described above may be readily prepared from the DNA sequences using any of a variety of expression vectors known to those of ordinary skill in the art. Expression may be achieved in any appropriate host cell that has been transformed or transfected with an expression vector containing a DNA molecule that encodes a recombinant polypeptide. Suitable host cells include prokaryotes, yeast, and higher eukaryotic cells, such as mammalian cells and plant cells. Preferably, the host cells employed are *E. coli*, yeast or a mammalian cell line such as COS or CHO. Supernatants from suitable host/vector systems which secrete recombinant protein or polypeptide into culture media may be first concentrated using a commercially available filter. Following concentration, the concentrate may be applied to a suitable purification matrix such as an affinity matrix or an ion exchange

20

25

30

resin. Finally, one or more reverse phase HPLC steps can be employed to further purify a recombinant polypeptide.

Portions and other variants having less than about 100 amino acids, and generally less than about 50 amino acids, may also be generated by synthetic means, using techniques well known to those of ordinary skill in the art. For example, such polypeptides may be synthesized using any of the commercially available solid-phase techniques, such as the Merrifield solid-phase synthesis method, where amino acids are sequentially added to a growing amino acid chain. See Merrifield, *J. Am. Chem. Soc.* 85:2149-2146, 1963. Equipment for automated synthesis of polypeptides is commercially available from suppliers such as Perkin Elmer/Applied BioSystems Division (Foster City, CA), and may be operated according to the manufacturer's instructions.

Within certain specific embodiments, a polypeptide may be a fusion protein that comprises multiple polypeptides as described herein, or that comprises at least one polypeptide as described herein and an unrelated sequence, such as a known tumor protein. A fusion partner may, for example, assist in providing T helper epitopes (an immunological fusion partner), preferably T helper epitopes recognized by humans, or may assist in expressing the protein (an expression enhancer) at higher yields than the native recombinant protein. Certain preferred fusion partners are both immunological and expression enhancing fusion partners. Other fusion partners may be selected so as to increase the solubility of the protein or to enable the protein to be targeted to desired intracellular compartments. Still further fusion partners include affinity tags, which facilitate purification of the protein.

Fusion proteins may generally be prepared using standard techniques, including chemical conjugation. Preferably, a fusion protein is expressed as a recombinant protein, allowing the production of increased levels, relative to a non-fused protein, in an expression system. Briefly, DNA sequences encoding the polypeptide components may be assembled separately, and ligated into an appropriate expression vector. The 3' end of the DNA sequence encoding one polypeptide component is ligated, with or without a peptide linker, to the 5' end of a DNA sequence encoding the second polypeptide component so that the reading frames of the sequences are in phase.

This permits translation into a single fusion protein that retains the biological activity of both component polypeptides.

A peptide linker sequence may be employed to separate the first and second polypeptide components by a distance sufficient to ensure that each polypeptide folds into its secondary and tertiary structures. Such a peptide linker sequence is incorporated into the fusion protein using standard techniques well known in the art. Suitable peptide linker sequences may be chosen based on the following factors: (1) their ability to adopt a flexible extended conformation; (2) their inability to adopt a secondary structure that could interact with functional epitopes on the first and second polypeptides; and (3) the lack of hydrophobic or charged residues that might react with the polypeptide functional epitopes. Preferred peptide linker sequences contain Gly, Asn and Ser residues. Other near neutral amino acids, such as Thr and Ala may also be used in the linker sequence. Amino acid sequences which may be usefully employed as linkers include those disclosed in Maratea et al., *Gene* 40:39-46, 1985; Murphy et al., *Proc. Natl. Acad. Sci. USA* 83:8258-8262, 1986; U.S. Patent No. 4,935,233 and U.S. Patent No. 4,751,180. The linker sequence may generally be from 1 to about 50 amino acids in length. Linker sequences are not required when the first and second polypeptides have non-essential N-terminal amino acid regions that can be used to separate the functional domains and prevent steric interference.

The ligated DNA sequences are operably linked to suitable transcriptional or translational regulatory elements. The regulatory elements responsible for expression of DNA are located only 5' to the DNA sequence encoding the first polypeptides. Similarly, stop codons required to end translation and transcription termination signals are only present 3' to the DNA sequence encoding the second polypeptide.

Fusion proteins are also provided. Such proteins comprise a polypeptide as described herein together with an unrelated immunogenic protein. Preferably the immunogenic protein is capable of eliciting a recall response. Examples of such proteins include tetanus, tuberculosis and hepatitis proteins (see, for example, Stoute et al. *New Engl. J. Med.*, 336:86-91, 1997).

Within preferred embodiments, an immunological fusion partner is derived from protein D, a surface protein of the gram-negative bacterium *Haemophilus influenza B* (WO 91/18926). Preferably, a protein D derivative comprises approximately the first third of the protein (*e.g.*, the first N-terminal 100-110 amino acids), and a protein D derivative may be lipidated. Within certain preferred  
5       embodiments, the first 109 residues of a Lipoprotein D fusion partner is included on the N-terminus to provide the polypeptide with additional exogenous T-cell epitopes and to increase the expression level in *E. coli* (thus functioning as an expression enhancer). The lipid tail ensures optimal presentation of the antigen to antigen presenting cells.  
10      Other fusion partners include the non-structural protein from influenzae virus, NS1 (hemagglutinin). Typically, the N-terminal 81 amino acids are used, although different fragments that include T-helper epitopes may be used.

In another embodiment, the immunological fusion partner is the protein known as LYTA, or a portion thereof (preferably a C-terminal portion). LYTA is  
15      derived from *Streptococcus pneumoniae*, which synthesizes an N-acetyl-L-alanine amidase known as amidase LYTA (encoded by the *LytA* gene; *Gene* 43:265-292, 1986). LYTA is an autolysin that specifically degrades certain bonds in the peptidoglycan backbone. The C-terminal domain of the LYTA protein is responsible for the affinity to the choline or to some choline analogues such as DEAE. This  
20      property has been exploited for the development of *E. coli* C-LYTA expressing plasmids useful for expression of fusion proteins. Purification of hybrid proteins containing the C-LYTA fragment at the amino terminus has been described (*see Biotechnology* 10:795-798, 1992). Within a preferred embodiment, a repeat portion of LYTA may be incorporated into a fusion protein. A repeat portion is found in the C-  
25      terminal region starting at residue 178. A particularly preferred repeat portion incorporates residues 188-305.

In general, polypeptides (including fusion proteins) and polynucleotides as described herein are isolated. An "isolated" polypeptide or polynucleotide is one that is removed from its original environment. For example, a naturally-occurring protein is  
30      isolated if it is separated from some or all of the coexisting materials in the natural system. Preferably, such polypeptides are at least about 90% pure, more preferably at

least about 95% pure and most preferably at least about 99% pure. A polynucleotide is considered to be isolated if, for example, it is cloned into a vector that is not a part of the natural environment.

#### BINDING AGENTS

5           The present invention further provides agents, such as antibodies and antigen-binding fragments thereof, that specifically bind to a breast tumor protein. As used herein, an antibody, or antigen-binding fragment thereof, is said to "specifically bind" to a breast tumor protein if it reacts at a detectable level (within, for example, an ELISA) with a breast tumor protein, and does not react detectably with unrelated  
10 proteins under similar conditions. As used herein, "binding" refers to a noncovalent association between two separate molecules such that a complex is formed. The ability to bind may be evaluated by, for example, determining a binding constant for the formation of the complex. The binding constant is the value obtained when the concentration of the complex is divided by the product of the component  
15 concentrations. In general, two compounds are said to "bind," in the context of the present invention, when the binding constant for complex formation exceeds about  $10^3$  L/mol. The binding constant may be determined using methods well known in the art.

          Binding agents may be further capable of differentiating between patients with and without a cancer, such as breast cancer, using the representative assays  
20 provided herein. In other words, antibodies or other binding agents that bind to a breast tumor protein will generate a signal indicating the presence of a cancer in at least about 20% of patients with the disease, and will generate a negative signal indicating the absence of the disease in at least about 90% of individuals without the cancer. To determine whether a binding agent satisfies this requirement, biological samples (*e.g.*,  
25 blood, sera, sputum, urine and/or tumor biopsies) from patients with and without a cancer (as determined using standard clinical tests) may be assayed as described herein for the presence of polypeptides that bind to the binding agent. It will be apparent that a statistically significant number of samples with and without the disease should be assayed. Each binding agent should satisfy the above criteria; however, those of



ordinary skill in the art will recognize that binding agents may be used in combination to improve sensitivity.

Any agent that satisfies the above requirements may be a binding agent. For example, a binding agent may be a ribosome, with or without a peptide component, an RNA molecule or a polypeptide. In a preferred embodiment, a binding agent is an antibody or an antigen-binding fragment thereof. Antibodies may be prepared by any of a variety of techniques known to those of ordinary skill in the art. See, e.g., Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. In general, antibodies can be produced by cell culture techniques, including the generation of monoclonal antibodies as described herein, or via transfection of antibody genes into suitable bacterial or mammalian cell hosts, in order to allow for the production of recombinant antibodies. In one technique, an immunogen comprising the polypeptide is initially injected into any of a wide variety of mammals (e.g., mice, rats, rabbits, sheep or goats). In this step, the polypeptides of this invention may serve as the immunogen without modification. Alternatively, particularly for relatively short polypeptides, a superior immune response may be elicited if the polypeptide is joined to a carrier protein, such as bovine serum albumin or keyhole limpet hemocyanin. The immunogen is injected into the animal host, preferably according to a predetermined schedule incorporating one or more booster immunizations, and the animals are bled periodically. Polyclonal antibodies specific for the polypeptide may then be purified from such antisera by, for example, affinity chromatography using the polypeptide coupled to a suitable solid support.

Monoclonal antibodies specific for an antigenic polypeptide of interest may be prepared, for example, using the technique of Kohler and Milstein, *Eur. J. Immunol.* 6:511-519, 1976, and improvements thereto. Briefly, these methods involve the preparation of immortal cell lines capable of producing antibodies having the desired specificity (i.e., reactivity with the polypeptide of interest). Such cell lines may be produced, for example, from spleen cells obtained from an animal immunized as described above. The spleen cells are then immortalized by, for example, fusion with a myeloma cell fusion partner, preferably one that is syngeneic with the immunized animal. A variety of fusion techniques may be employed. For example, the spleen cells

and myeloma cells may be combined with a nonionic detergent for a few minutes and then plated at low density on a selective medium that supports the growth of hybrid cells, but not myeloma cells. A preferred selection technique uses HAT (hypoxanthine, aminopterin, thymidine) selection. After a sufficient time, usually about 1 to 2 weeks,  
5 colonies of hybrids are observed. Single colonies are selected and their culture supernatants tested for binding activity against the polypeptide. Hybridomas having high reactivity and specificity are preferred.

Monoclonal antibodies may be isolated from the supernatants of growing hybridoma colonies. In addition, various techniques may be employed to enhance the  
10 yield, such as injection of the hybridoma cell line into the peritoneal cavity of a suitable vertebrate host, such as a mouse. Monoclonal antibodies may then be harvested from the ascites fluid or the blood. Contaminants may be removed from the antibodies by conventional techniques, such as chromatography, gel filtration, precipitation, and extraction. The polypeptides of this invention may be used in the purification process  
15 in, for example, an affinity chromatography step.

Within certain embodiments, the use of antigen-binding fragments of antibodies may be preferred. Such fragments include Fab fragments, which may be prepared using standard techniques. Briefly, immunoglobulins may be purified from rabbit serum by affinity chromatography on Protein A bead columns (Harlow and Lane,  
20 *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988) and digested by papain to yield Fab and Fc fragments. The Fab and Fc fragments may be separated by affinity chromatography on protein A bead columns.

Monoclonal antibodies of the present invention may be coupled to one or more therapeutic agents. Suitable agents in this regard include radionuclides,  
25 differentiation inducers, drugs, toxins, and derivatives thereof. Preferred radionuclides include  $^{90}\text{Y}$ ,  $^{123}\text{I}$ ,  $^{125}\text{I}$ ,  $^{131}\text{I}$ ,  $^{186}\text{Re}$ ,  $^{188}\text{Re}$ ,  $^{211}\text{At}$ , and  $^{212}\text{Bi}$ . Preferred drugs include methotrexate, and pyrimidine and purine analogs. Preferred differentiation inducers include phorbol esters and butyric acid. Preferred toxins include ricin, abrin, diphtheria toxin, cholera toxin, gelonin, *Pseudomonas* exotoxin, *Shigella* toxin, and pokeweed  
30 antiviral protein.

A therapeutic agent may be coupled (*e.g.*, covalently bonded) to a suitable monoclonal antibody either directly or indirectly (*e.g.*, via a linker group). A direct reaction between an agent and an antibody is possible when each possesses a substituent capable of reacting with the other. For example, a nucleophilic group, such as an amino or sulfhydryl group, on one may be capable of reacting with a carbonyl-containing group, such as an anhydride or an acid halide, or with an alkyl group containing a good leaving group (*e.g.*, a halide) on the other.

Alternatively, it may be desirable to couple a therapeutic agent and an antibody via a linker group. A linker group can function as a spacer to distance an antibody from an agent in order to avoid interference with binding capabilities. A linker group can also serve to increase the chemical reactivity of a substituent on an agent or an antibody, and thus increase the coupling efficiency. An increase in chemical reactivity may also facilitate the use of agents, or functional groups on agents, which otherwise would not be possible.

It will be evident to those skilled in the art that a variety of bifunctional or polyfunctional reagents, both homo- and hetero-functional (such as those described in the catalog of the Pierce Chemical Co., Rockford, IL), may be employed as the linker group. Coupling may be effected, for example, through amino groups, carboxyl groups, sulfhydryl groups or oxidized carbohydrate residues. There are numerous references describing such methodology, *e.g.*, U.S. Patent No. 4,671,958, to Rodwell et al.

Where a therapeutic agent is more potent when free from the antibody portion of the immunoconjugates of the present invention, it may be desirable to use a linker group which is cleavable during or upon internalization into a cell. A number of different cleavable linker groups have been described. The mechanisms for the intracellular release of an agent from these linker groups include cleavage by reduction of a disulfide bond (*e.g.*, U.S. Patent No. 4,489,710, to Spitler), by irradiation of a photolabile bond (*e.g.*, U.S. Patent No. 4,625,014, to Senter et al.), by hydrolysis of derivatized amino acid side chains (*e.g.*, U.S. Patent No. 4,638,045, to Kohn et al.), by serum complement-mediated hydrolysis (*e.g.*, U.S. Patent No. 4,671,958, to Rodwell et al.), and acid-catalyzed hydrolysis (*e.g.*, U.S. Patent No. 4,569,789, to Blattler et al.).

It may be desirable to couple more than one agent to an antibody. In one embodiment, multiple molecules of an agent are coupled to one antibody molecule. In another embodiment, more than one type of agent may be coupled to one antibody. Regardless of the particular embodiment, immunoconjugates with more than one agent  
5 may be prepared in a variety of ways. For example, more than one agent may be coupled directly to an antibody molecule, or linkers that provide multiple sites for attachment can be used. Alternatively, a carrier can be used.

A carrier may bear the agents in a variety of ways, including covalent bonding either directly or via a linker group. Suitable carriers include proteins such as  
10 albumins (*e.g.*, U.S. Patent No. 4,507,234, to Kato et al.), peptides and polysaccharides such as aminodextran (*e.g.*, U.S. Patent No. 4,699,784, to Shih et al.). A carrier may also bear an agent by noncovalent bonding or by encapsulation, such as within a liposome vesicle (*e.g.*, U.S. Patent Nos. 4,429,008 and 4,873,088). Carriers specific for radionuclide agents include radiohalogenated small molecules and chelating  
15 compounds. For example, U.S. Patent No. 4,735,792 discloses representative radiohalogenated small molecules and their synthesis. A radionuclide chelate may be formed from chelating compounds that include those containing nitrogen and sulfur atoms as the donor atoms for binding the metal, or metal oxide, radionuclide. For example, U.S. Patent No. 4,673,562, to Davison et al. discloses representative chelating  
20 compounds and their synthesis.

A variety of routes of administration for the antibodies and immunoconjugates may be used. Typically, administration will be intravenous, intramuscular, subcutaneous or in the bed of a resected tumor. It will be evident that the precise dose of the antibody/immunoconjugate will vary depending upon the antibody  
25 used, the antigen density on the tumor, and the rate of clearance of the antibody.

#### T CELLS

Immunotherapeutic compositions may also, or alternatively, comprise T cells specific for a breast tumor protein. Such cells may generally be prepared *in vitro* or *ex vivo*, using standard procedures. For example, T cells may be isolated from bone  
30 marrow, peripheral blood, or a fraction of bone marrow or peripheral blood of a patient,

using a commercially available cell separation system, such as the Isolex™ System, available from Nexell Therapeutics, Inc. (Irvine, CA; see also U.S. Patent No. 5,240,856; U.S. Patent No. 5,215,926; WO 89/06280; WO 91/16116 and WO 92/07243). Alternatively, T cells may be derived from related or unrelated humans,  
5 non-human mammals, cell lines or cultures.

T cells may be stimulated with a breast tumor polypeptide, polynucleotide encoding a breast tumor polypeptide and/or an antigen presenting cell (APC) that expresses such a polypeptide. Such stimulation is performed under conditions and for a time sufficient to permit the generation of T cells that are specific  
10 for the polypeptide. Preferably, a breast tumor polypeptide or polynucleotide is present within a delivery vehicle, such as a microsphere, to facilitate the generation of specific T cells.

T cells are considered to be specific for a breast tumor polypeptide if the T cells specifically proliferate, secrete cytokines or kill target cells coated with the  
15 polypeptide or expressing a gene encoding the polypeptide. T cell specificity may be evaluated using any of a variety of standard techniques. For example, within a chromium release assay or proliferation assay, a stimulation index of more than two fold increase in lysis and/or proliferation, compared to negative controls, indicates T cell specificity. Such assays may be performed, for example, as described in Chen et al., *Cancer Res.* 54:1065-1070, 1994. Alternatively, detection of the proliferation of  
20 T cells may be accomplished by a variety of known techniques. For example, T cell proliferation can be detected by measuring an increased rate of DNA synthesis (e.g., by pulse-labeling cultures of T cells with tritiated thymidine and measuring the amount of tritiated thymidine incorporated into DNA). Contact with a breast tumor polypeptide  
25 (100 ng/ml - 100 µg/ml, preferably 200 ng/ml - 25 µg/ml) for 3 - 7 days should result in at least a two fold increase in proliferation of the T cells. Contact as described above for 2-3 hours should result in activation of the T cells, as measured using standard cytokine assays in which a two fold increase in the level of cytokine release (e.g., TNF or IFN-γ) is indicative of T cell activation (see Coligan et al., *Current Protocols in*  
30 *Immunology*, vol. 1, Wiley Interscience (Greene 1998)). T cells that have been activated in response to a breast tumor polypeptide, polynucleotide or polypeptide-

expressing APC may be CD4<sup>+</sup> and/or CD8<sup>+</sup>. Breast tumor protein-specific T cells may be expanded using standard techniques. Within preferred embodiments, the T cells are derived from a patient, a related donor or an unrelated donor, and are administered to the patient following stimulation and expansion.

5               For therapeutic purposes, CD4<sup>+</sup> or CD8<sup>+</sup> T cells that proliferate in response to a breast tumor polypeptide, polynucleotide or APC can be expanded in number either *in vitro* or *in vivo*. Proliferation of such T cells *in vitro* may be accomplished in a variety of ways. For example, the T cells can be re-exposed to a breast tumor polypeptide, or a short peptide corresponding to an immunogenic portion  
10 of such a polypeptide, with or without the addition of T cell growth factors, such as interleukin-2, and/or stimulator cells that synthesize a breast tumor polypeptide. Alternatively, one or more T cells that proliferate in the presence of a breast tumor protein can be expanded in number by cloning. Methods for cloning cells are well known in the art, and include limiting dilution.

## 15   PHARMACEUTICAL COMPOSITIONS

In additional embodiments, the present invention concerns formulation of one or more of the polynucleotide, polypeptide, T-cell and/or antibody compositions disclosed herein in pharmaceutically-acceptable solutions for administration to a cell or an animal, either alone, or in combination with one or more other modalities of therapy.

20               It will also be understood that, if desired, the nucleic acid segment, RNA, DNA or PNA compositions that express a polypeptide as disclosed herein may be administered in combination with other agents as well, such as, *e.g.*, other proteins or polypeptides or various pharmaceutically-active agents. In fact, there is virtually no limit to other components that may also be included, given that the additional agents do  
25 not cause a significant adverse effect upon contact with the target cells or host tissues. The compositions may thus be delivered along with various other agents as required in the particular instance. Such compositions may be purified from host cells or other biological sources, or alternatively may be chemically synthesized as described herein. Likewise, such compositions may further comprise substituted or derivatized RNA or  
30 DNA compositions.

Formulation of pharmaceutically-acceptable excipients and carrier solutions is well-known to those of skill in the art, as is the development of suitable dosing and treatment regimens for using the particular compositions described herein in a variety of treatment regimens, including *e.g.*, oral, parenteral, intravenous, intranasal, and intramuscular administration and formulation.

#### 1. ORAL DELIVERY

In certain applications, the pharmaceutical compositions disclosed herein may be delivered *via* oral administration to an animal. As such, these compositions may be formulated with an inert diluent or with an assimilable edible carrier, or they may be enclosed in hard- or soft-shell gelatin capsule, or they may be compressed into tablets, or they may be incorporated directly with the food of the diet.

The active compounds may even be incorporated with excipients and used in the form of ingestible tablets, buccal tables, troches, capsules, elixirs, suspensions, syrups, wafers, and the like (Mathiowitz *et al.*, 1997; Hwang *et al.*, 1998; U. S. Patent 5,641,515; U. S. Patent 5,580,579 and U. S. Patent 5,792,451, each specifically incorporated herein by reference in its entirety). The tablets, troches, pills, capsules and the like may also contain the following: a binder, as gum tragacanth, acacia, cornstarch, or gelatin; excipients, such as dicalcium phosphate; a disintegrating agent, such as corn starch, potato starch, alginic acid and the like; a lubricant, such as magnesium stearate; and a sweetening agent, such as sucrose, lactose or saccharin may be added or a flavoring agent, such as peppermint, oil of wintergreen, or cherry flavoring. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier. Various other materials may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, tablets, pills, or capsules may be coated with shellac, sugar, or both. A syrup or elixir may contain the active compound sucrose as a sweetening agent methyl and propylparabens as preservatives, a dye and flavoring, such as cherry or orange flavor. Of course, any material used in preparing any dosage unit form should be pharmaceutically pure and substantially non-toxic in the amounts employed. In

addition, the active compounds may be incorporated into sustained-release preparation and formulations.

Typically, these formulations may contain at least about 0.1% of the active compound or more, although the percentage of the active ingredient(s) may, of course, be varied and may conveniently be between about 1 or 2% and about 60% or 70% or more of the weight or volume of the total formulation. Naturally, the amount of active compound(s) in each therapeutically useful composition may be prepared in such a way that a suitable dosage will be obtained in any given unit dose of the compound. Factors such as solubility, bioavailability, biological half-life, route of administration, product shelf life, as well as other pharmacological considerations will be contemplated by one skilled in the art of preparing such pharmaceutical formulations, and as such, a variety of dosages and treatment regimens may be desirable.

For oral administration the compositions of the present invention may alternatively be incorporated with one or more excipients in the form of a mouthwash, dentifrice, buccal tablet, oral spray, or sublingual orally-administered formulation. For example, a mouthwash may be prepared incorporating the active ingredient in the required amount in an appropriate solvent, such as a sodium borate solution (Dobell's Solution). Alternatively, the active ingredient may be incorporated into an oral solution such as one containing sodium borate, glycerin and potassium bicarbonate, or dispersed in a dentifrice, or added in a therapeutically-effective amount to a composition that may include water, binders, abrasives, flavoring agents, foaming agents, and humectants. Alternatively the compositions may be fashioned into a tablet or solution form that may be placed under the tongue or otherwise dissolved in the mouth.

## 2. INJECTABLE DELIVERY

In certain circumstances it will be desirable to deliver the pharmaceutical compositions disclosed herein parenterally, intravenously, intramuscularly, or even intraperitoneally as described in U. S. Patent 5,543,158; U. S. Patent 5,641,515 and U. S. Patent 5,399,363 (each specifically incorporated herein by reference in its entirety). Solutions of the active compounds as free base or pharmacologically acceptable salts may be prepared in water suitably mixed with a surfactant, such as



hydroxypropylcellulose. Dispersions may also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

5           The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions (U. S. Patent 5,466,468, specifically incorporated herein by reference in its entirety). In all cases the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable  
10 under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms, such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (*e.g.*, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and/or vegetable oils. Proper fluidity may be maintained, for  
15 example, by the use of a coating, such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. The prevention of the action of microorganisms can be facilitated by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars  
20 or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

          For parenteral administration in an aqueous solution, for example, the solution should be suitably buffered if necessary and the liquid diluent first rendered  
25 isotonic with sufficient saline or glucose. These particular aqueous solutions are especially suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. In this connection, a sterile aqueous medium that can be employed will be known to those of skill in the art in light of the present disclosure. For example, one dosage may be dissolved in 1 ml of isotonic NaCl solution and either added to 1000 ml  
30 of hypodermoclysis fluid or injected at the proposed site of infusion, (see for example, "Remington's Pharmaceutical Sciences" 15th Edition, pages 1035-1038 and 1570-

1580). Some variation in dosage will necessarily occur depending on the condition of the subject being treated. The person responsible for administration will, in any event, determine the appropriate dose for the individual subject. Moreover, for human administration, preparations should meet sterility, pyrogenicity, and the general safety and purity standards as required by FDA Office of Biologics standards.

Sterile injectable solutions are prepared by incorporating the active compounds in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum-drying and freeze-drying techniques which yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

The compositions disclosed herein may be formulated in a neutral or salt form. Pharmaceutically-acceptable salts, include the acid addition salts (formed with the free amino groups of the protein) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, histidine, procaine and the like. Upon formulation, solutions will be administered in a manner compatible with the dosage formulation and in such amount as is therapeutically effective. The formulations are easily administered in a variety of dosage forms such as injectable solutions, drug-release capsules, and the like.

As used herein, "carrier" includes any and all solvents, dispersion media, vehicles, coatings, diluents, antibacterial and antifungal agents, isotonic and absorption delaying agents, buffers, carrier solutions, suspensions, colloids, and the like. The use of such media and agents for pharmaceutical active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active

ingredient, its use in the therapeutic compositions is contemplated. Supplementary active ingredients can also be incorporated into the compositions.

The phrase "pharmaceutically-acceptable" refers to molecular entities and compositions that do not produce an allergic or similar untoward reaction when administered to a human. The preparation of an aqueous composition that contains a protein as an active ingredient is well understood in the art. Typically, such compositions are prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid prior to injection can also be prepared. The preparation can also be emulsified.

### 10 3. NASAL DELIVERY

In certain embodiments, the pharmaceutical compositions may be delivered by intranasal sprays, inhalation, and/or other aerosol delivery vehicles. Methods for delivering genes, nucleic acids, and peptide compositions directly to the lungs *via* nasal aerosol sprays has been described *e.g.*, in U. S. Patent 5,756,353 and U. S. Patent 5,804,212 (each specifically incorporated herein by reference in its entirety). Likewise, the delivery of drugs using intranasal microparticle resins (Takenaga *et al.*, 1998) and lysophosphatidyl-glycerol compounds (U. S. Patent 5,725,871, specifically incorporated herein by reference in its entirety) are also well-known in the pharmaceutical arts. Likewise, transmucosal drug delivery in the form of a polytetrafluoroethylene support matrix is described in U. S. Patent 5,780,045 (specifically incorporated herein by reference in its entirety).

### 4. LIPOSOME-, NANOCAPSULE-, AND MICROPARTICLE-MEDIATED DELIVERY

In certain embodiments, the inventors contemplate the use of liposomes, nanocapsules, microparticles, microspheres, lipid particles, vesicles, and the like, for the introduction of the compositions of the present invention into suitable host cells. In particular, the compositions of the present invention may be formulated for delivery either encapsulated in a lipid particle, a liposome, a vesicle, a nanosphere, or a nanoparticle or the like.

Such formulations may be preferred for the introduction of pharmaceutically-acceptable formulations of the nucleic acids or constructs disclosed herein. The formation and use of liposomes is generally known to those of skill in the art (see for example, Couvreur *et al.*, 1977; Couvreur, 1988; Lasic, 1998; which  
5 describes the use of liposomes and nanocapsules in the targeted antibiotic therapy for intracellular bacterial infections and diseases). Recently, liposomes were developed with improved serum stability and circulation half-times (Gabizon and Papahadjopoulos, 1988; Allen and Choun, 1987; U. S. Patent 5,741,516, specifically incorporated herein by reference in its entirety). Further, various methods of liposome  
10 and liposome like preparations as potential drug carriers have been reviewed (Takakura, 1998; Chandran *et al.*, 1997; Margalit, 1995; U. S. Patent 5,567,434; U. S. Patent 5,552,157; U. S. Patent 5,565,213; U. S. Patent 5,738,868 and U. S. Patent 5,795,587, each specifically incorporated herein by reference in its entirety).

Liposomes have been used successfully with a number of cell types that  
15 are normally resistant to transfection by other procedures including T cell suspensions, primary hepatocyte cultures and PC 12 cells (Renneisen *et al.*, 1990; Muller *et al.*, 1990). In addition, liposomes are free of the DNA length constraints that are typical of viral-based delivery systems. Liposomes have been used effectively to introduce genes, drugs (Heath and Martin, 1986; Heath *et al.*, 1986; Balazsovits *et al.*, 1989; Fresta and  
20 Puglisi, 1996), radiotherapeutic agents (Pikul *et al.*, 1987), enzymes (Imaizumi *et al.*, 1990a; Imaizumi *et al.*, 1990b), viruses (Faller and Baltimore, 1984), transcription factors and allosteric effectors (Nicolau and Gersonde, 1979) into a variety of cultured cell lines and animals. In addition, several successful clinical trails examining the effectiveness of liposome-mediated drug delivery have been completed (Lopez-  
25 Berestein *et al.*, 1985a; 1985b; Coune, 1988; Sculier *et al.*, 1988). Furthermore, several studies suggest that the use of liposomes is not associated with autoimmune responses, toxicity or gonadal localization after systemic delivery (Mori and Fukatsu, 1992).

Liposomes are formed from phospholipids that are dispersed in an aqueous medium and spontaneously form multilamellar concentric bilayer vesicles (also  
30 termed multilamellar vesicles (MLVs). MLVs generally have diameters of from 25 nm to 4  $\mu$ m. Sonication of MLVs results in the formation of small unilamellar vesicles

(SUVs) with diameters in the range of 200 to 500 Å, containing an aqueous solution in the core.

Liposomes bear resemblance to cellular membranes and are contemplated for use in connection with the present invention as carriers for the peptide compositions. They are widely suitable as both water- and lipid-soluble substances can be entrapped, *i.e.* in the aqueous spaces and within the bilayer itself, respectively. It is possible that the drug-bearing liposomes may even be employed for site-specific delivery of active agents by selectively modifying the liposomal formulation.

In addition to the teachings of Couvreur *et al.* (1977; 1988), the following information may be utilized in generating liposomal formulations. Phospholipids can form a variety of structures other than liposomes when dispersed in water, depending on the molar ratio of lipid to water. At low ratios the liposome is the preferred structure. The physical characteristics of liposomes depend on pH, ionic strength and the presence of divalent cations. Liposomes can show low permeability to ionic and polar substances, but at elevated temperatures undergo a phase transition which markedly alters their permeability. The phase transition involves a change from a closely packed, ordered structure, known as the gel state, to a loosely packed, less-ordered structure, known as the fluid state. This occurs at a characteristic phase-transition temperature and results in an increase in permeability to ions, sugars and drugs.

In addition to temperature, exposure to proteins can alter the permeability of liposomes. Certain soluble proteins, such as cytochrome c, bind, deform and penetrate the bilayer, thereby causing changes in permeability. Cholesterol inhibits this penetration of proteins, apparently by packing the phospholipids more tightly. It is contemplated that the most useful liposome formations for antibiotic and inhibitor delivery will contain cholesterol.

The ability to trap solutes varies between different types of liposomes. For example, MLVs are moderately efficient at trapping solutes, but SUVs are extremely inefficient. SUVs offer the advantage of homogeneity and reproducibility in size distribution, however, and a compromise between size and trapping efficiency is

offered by large unilamellar vesicles (LUVs). These are prepared by ether evaporation and are three to four times more efficient at solute entrapment than MLVs.

In addition to liposome characteristics, an important determinant in entrapping compounds is the physicochemical properties of the compound itself. Polar compounds are trapped in the aqueous spaces and nonpolar compounds bind to the lipid bilayer of the vesicle. Polar compounds are released through permeation or when the bilayer is broken, but nonpolar compounds remain affiliated with the bilayer unless it is disrupted by temperature or exposure to lipoproteins. Both types show maximum efflux rates at the phase transition temperature.

Liposomes interact with cells *via* four different mechanisms: endocytosis by phagocytic cells of the reticuloendothelial system such as macrophages and neutrophils; adsorption to the cell surface, either by nonspecific weak hydrophobic or electrostatic forces, or by specific interactions with cell-surface components; fusion with the plasma cell membrane by insertion of the lipid bilayer of the liposome into the plasma membrane, with simultaneous release of liposomal contents into the cytoplasm; and by transfer of liposomal lipids to cellular or subcellular membranes, or vice versa, without any association of the liposome contents. It often is difficult to determine which mechanism is operative and more than one may operate at the same time.

The fate and disposition of intravenously injected liposomes depend on their physical properties, such as size, fluidity, and surface charge. They may persist in tissues for h or days, depending on their composition, and half lives in the blood range from min to several h. Larger liposomes, such as MLVs and LUVs, are taken up rapidly by phagocytic cells of the reticuloendothelial system, but physiology of the circulatory system restrains the exit of such large species at most sites. They can exit only in places where large openings or pores exist in the capillary endothelium, such as the sinusoids of the liver or spleen. Thus, these organs are the predominate site of uptake. On the other hand, SUVs show a broader tissue distribution but still are sequestered highly in the liver and spleen. In general, this *in vivo* behavior limits the potential targeting of liposomes to only those organs and tissues accessible to their large size. These include the blood, liver, spleen, bone marrow, and lymphoid organs.

Targeting is generally not a limitation in terms of the present invention. However, should specific targeting be desired, methods are available for this to be accomplished. Antibodies may be used to bind to the liposome surface and to direct the antibody and its drug contents to specific antigenic receptors located on a particular cell-type surface. Carbohydrate determinants (glycoprotein or glycolipid cell-surface components that play a role in cell-cell recognition, interaction and adhesion) may also be used as recognition sites as they have potential in directing liposomes to particular cell types. Mostly, it is contemplated that intravenous injection of liposomal preparations would be used, but other routes of administration are also conceivable.

Alternatively, the invention provides for pharmaceutically-acceptable nanocapsule formulations of the compositions of the present invention. Nanocapsules can generally entrap compounds in a stable and reproducible way (Henry-Michelland *et al.*, 1987; Quintanar-Guerrero *et al.*, 1998; Douglas *et al.*, 1987). To avoid side effects due to intracellular polymeric overloading, such ultrafine particles (sized around 0.1  $\mu\text{m}$ ) should be designed using polymers able to be degraded *in vivo*. Biodegradable polyalkyl-cyanoacrylate nanoparticles that meet these requirements are contemplated for use in the present invention. Such particles may be easily made, as described (Couvreur *et al.*, 1980; 1988; zur Muhlen *et al.*, 1998; Zambaux *et al.* 1998; Pinto-Alphandry *et al.*, 1995 and U. S. Patent 5,145,684, specifically incorporated herein by reference in its entirety).

#### VACCINES

In certain preferred embodiments of the present invention, vaccines are provided. The vaccines will generally comprise one or more pharmaceutical compositions, such as those discussed above, in combination with an immunostimulant.

An immunostimulant may be any substance that enhances or potentiates an immune response (antibody and/or cell-mediated) to an exogenous antigen. Examples of immunostimulants include adjuvants, biodegradable microspheres (*e.g.*, polylactic galactide) and liposomes (into which the compound is incorporated; *see e.g.*, Fullerton, U.S. Patent No. 4,235,877). Vaccine preparation is generally described in, for example, M.F. Powell and M.J. Newman, eds., "Vaccine Design (the subunit and adjuvant

approach)," Plenum Press (NY, 1995). Pharmaceutical compositions and vaccines within the scope of the present invention may also contain other compounds, which may be biologically active or inactive. For example, one or more immunogenic portions of other tumor antigens may be present, either incorporated into a fusion  
5 polypeptide or as a separate compound, within the composition or vaccine.

Illustrative vaccines may contain DNA encoding one or more of the polypeptides as described above, such that the polypeptide is generated *in situ*. As noted above, the DNA may be present within any of a variety of delivery systems known to those of ordinary skill in the art, including nucleic acid expression systems,  
10 bacteria and viral expression systems. Numerous gene delivery techniques are well known in the art, such as those described by Rolland, *Crit. Rev. Therap. Drug Carrier Systems* 15:143-198, 1998, and references cited therein. Appropriate nucleic acid expression systems contain the necessary DNA sequences for expression in the patient (such as a suitable promoter and terminating signal). Bacterial delivery systems involve  
15 the administration of a bacterium (such as *Bacillus-Calmette-Guerrin*) that expresses an immunogenic portion of the polypeptide on its cell surface or secretes such an epitope. In a preferred embodiment, the DNA may be introduced using a viral expression system (*e.g.*, vaccinia or other pox virus, retrovirus, or adenovirus), which may involve the use of a non-pathogenic (defective), replication competent virus. Suitable systems are  
20 disclosed, for example, in Fisher-Hoch et al., *Proc. Natl. Acad. Sci. USA* 86:317-321, 1989; Flexner et al., *Ann. N.Y. Acad. Sci.* 569:86-103, 1989; Flexner et al., *Vaccine* 8:17-21, 1990; U.S. Patent Nos. 4,603,112, 4,769,330, and 5,017,487; WO 89/01973; U.S. Patent No. 4,777,127; GB 2,200,651; EP 0,345,242; WO 91/02805; Berkner, *Biotechniques* 6:616-627, 1988; Rosenfeld et al., *Science* 252:431-434, 1991; Kolls et  
25 al., *Proc. Natl. Acad. Sci. USA* 91:215-219, 1994; Kass-Eisler et al., *Proc. Natl. Acad. Sci. USA* 90:11498-11502, 1993; Guzmán et al., *Circulation* 88:2838-2848, 1993; and Guzman et al., *Cir. Res.* 73:1202-1207, 1993. Techniques for incorporating DNA into such expression systems are well known to those of ordinary skill in the art. The DNA may also be "naked," as described, for example, in Ulmer et al., *Science* 259:1745-1749,  
30 1993 and reviewed by Cohen, *Science* 259:1691-1692, 1993. The uptake of naked DNA may be increased by coating the DNA onto biodegradable beads, which are



efficiently transported into the cells. It will be apparent that a vaccine may comprise both a polynucleotide and a polypeptide component. Such vaccines may provide for an enhanced immune response.

It will be apparent that a vaccine may contain pharmaceutically acceptable salts of the polynucleotides and polypeptides provided herein. Such salts may be prepared from pharmaceutically acceptable non-toxic bases, including organic bases (*e.g.*, salts of primary, secondary and tertiary amines and basic amino acids) and inorganic bases (*e.g.*, sodium, potassium, lithium, ammonium, calcium and magnesium salts).

While any suitable carrier known to those of ordinary skill in the art may be employed in the vaccine compositions of this invention, the type of carrier will vary depending on the mode of administration. Compositions of the present invention may be formulated for any appropriate manner of administration, including for example, topical, oral, nasal, intravenous, intracranial, intraperitoneal, subcutaneous or intramuscular administration. For parenteral administration, such as subcutaneous injection, the carrier preferably comprises water, saline, alcohol, a fat, a wax or a buffer. For oral administration, any of the above carriers or a solid carrier, such as mannitol, lactose, starch, magnesium stearate, sodium saccharine, talcum, cellulose, glucose, sucrose, and magnesium carbonate, may be employed. Biodegradable microspheres (*e.g.*, polylactate polyglycolate) may also be employed as carriers for the pharmaceutical compositions of this invention. Suitable biodegradable microspheres are disclosed, for example, in U.S. Patent Nos. 4,897,268; 5,075,109; 5,928,647; 5,811,128; 5,820,883; 5,853,763; 5,814,344 and 5,942,252. One may also employ a carrier comprising the particulate-protein complexes described in U.S. Patent No. 5,928,647, which are capable of inducing a class I-restricted cytotoxic T lymphocyte responses in a host.

Such compositions may also comprise buffers (*e.g.*, neutral buffered saline or phosphate buffered saline), carbohydrates (*e.g.*, glucose, mannose, sucrose or dextrans), mannitol, proteins, polypeptides or amino acids such as glycine, antioxidants, bacteriostats, chelating agents such as EDTA or glutathione, adjuvants (*e.g.*, aluminum hydroxide), solutes that render the formulation isotonic, hypotonic or weakly hypertonic

with the blood of a recipient, suspending agents, thickening agents and/or preservatives. Alternatively, compositions of the present invention may be formulated as a lyophilizate. Compounds may also be encapsulated within liposomes using well known technology.

5           Any of a variety of immunostimulants may be employed in the vaccines of this invention. For example, an adjuvant may be included. Most adjuvants contain a substance designed to protect the antigen from rapid catabolism, such as aluminum hydroxide or mineral oil, and a stimulator of immune responses, such as lipid A, *Bordetella pertussis* or *Mycobacterium tuberculosis* derived proteins. Suitable  
10 adjuvants are commercially available as, for example, Freund's Incomplete Adjuvant and Complete Adjuvant (Difco Laboratories, Detroit, MI); Merck Adjuvant 65 (Merck and Company, Inc., Rahway, NJ); AS-2 (SmithKline Beecham, Philadelphia, PA); aluminum salts such as aluminum hydroxide gel (alum) or aluminum phosphate; salts of calcium, iron or zinc; an insoluble suspension of acylated tyrosine; acylated sugars;  
15 cationically or anionically derivatized polysaccharides; polyphosphazenes; biodegradable microspheres; monophosphoryl lipid A and quil A. Cytokines, such as GM-CSF or interleukin-2, -7, or -12, may also be used as adjuvants.

          Within the vaccines provided herein, the adjuvant composition is preferably designed to induce an immune response predominantly of the Th1 type.  
20 High levels of Th1-type cytokines (e.g., IFN- $\gamma$ , TNF $\alpha$ , IL-2 and IL-12) tend to favor the induction of cell mediated immune responses to an administered antigen. In contrast, high levels of Th2-type cytokines (e.g., IL-4, IL-5, IL-6 and IL-10) tend to favor the induction of humoral immune responses. Following application of a vaccine as provided herein, a patient will support an immune response that includes Th1- and Th2-  
25 type responses. Within a preferred embodiment, in which a response is predominantly Th1-type, the level of Th1-type cytokines will increase to a greater extent than the level of Th2-type cytokines. The levels of these cytokines may be readily assessed using standard assays. For a review of the families of cytokines, see Mosmann and Coffman, *Ann. Rev. Immunol.* 7:145-173, 1989.

30           Preferred adjuvants for use in eliciting a predominantly Th1-type response include, for example, a combination of monophosphoryl lipid A, preferably 3-

de-O-acylated monophosphoryl lipid A (3D-MPL), together with an aluminum salt. MPL adjuvants are available from Corixa Corporation (Seattle, WA; *see* US Patent Nos. 4,436,727; 4,877,611; 4,866,034 and 4,912,094). CpG-containing oligonucleotides (in which the CpG dinucleotide is unmethylated) also induce a predominantly Th1  
5 response. Such oligonucleotides are well known and are described, for example, in WO 96/02555, WO 99/33488 and U.S. Patent Nos. 6,008,200 and 5,856,462. Immunostimulatory DNA sequences are also described, for example, by Sato et al., *Science* 273:352, 1996. Another preferred adjuvant is a saponin, preferably QS21 (Aquila Biopharmaceuticals Inc., Framingham, MA), which may be used alone or in  
10 combination with other adjuvants. For example, an enhanced system involves the combination of a monophosphoryl lipid A and saponin derivative, such as the combination of QS21 and 3D-MPL as described in WO 94/00153, or a less reactogenic composition where the QS21 is quenched with cholesterol, as described in WO 96/33739. Other preferred formulations comprise an oil-in-water emulsion and  
15 tocopherol. A particularly potent adjuvant formulation involving QS21, 3D-MPL and tocopherol in an oil-in-water emulsion is described in WO 95/17210.

Other preferred adjuvants include Montanide ISA 720 (Seppic, France), SAF (Chiron, California, United States), ISCOMS (CSL), MF-59 (Chiron), the SBAS series of adjuvants (*e.g.*, SBAS-2 or SBAS-4, available from SmithKline Beecham,  
20 Rixensart, Belgium), Detox (Corixa, Hamilton, MT), RC-529 (Corixa, Hamilton, MT) and other aminoalkyl glucosaminide 4-phosphates (AGPs), such as those described in pending U.S. Patent Application Serial Nos. 08/853,826 and 09/074,720, the disclosures of which are incorporated herein by reference in their entireties.

Any vaccine provided herein may be prepared using well known  
25 methods that result in a combination of antigen, immune response enhancer and a suitable carrier or excipient. The compositions described herein may be administered as part of a sustained release formulation (*i.e.*, a formulation such as a capsule, sponge or gel (composed of polysaccharides, for example) that effects a slow release of compound following administration). Such formulations may generally be prepared using well  
30 known technology (*see, e.g.*, Coombes et al., *Vaccine* 14:1429-1438, 1996) and administered by, for example, oral, rectal or subcutaneous implantation, or by

implantation at the desired target site. Sustained-release formulations may contain a polypeptide, polynucleotide or antibody dispersed in a carrier matrix and/or contained within a reservoir surrounded by a rate controlling membrane.

Carriers for use within such formulations are biocompatible, and may also be biodegradable; preferably the formulation provides a relatively constant level of active component release. Such carriers include microparticles of poly(lactide-co-glycolide), polyacrylate, latex, starch, cellulose, dextran and the like. Other delayed-release carriers include supramolecular biovectors, which comprise a non-liquid hydrophilic core (e.g., a cross-linked polysaccharide or oligosaccharide) and, optionally, an external layer comprising an amphiphilic compound, such as a phospholipid (*see e.g.*, U.S. Patent No. 5,151,254 and PCT applications WO 94/20078, WO/94/23701 and WO 96/06638). The amount of active compound contained within a sustained release formulation depends upon the site of implantation, the rate and expected duration of release and the nature of the condition to be treated or prevented.

Any of a variety of delivery vehicles may be employed within pharmaceutical compositions and vaccines to facilitate production of an antigen-specific immune response that targets tumor cells. Delivery vehicles include antigen presenting cells (APCs), such as dendritic cells, macrophages, B cells, monocytes and other cells that may be engineered to be efficient APCs. Such cells may, but need not, be genetically modified to increase the capacity for presenting the antigen, to improve activation and/or maintenance of the T cell response, to have anti-tumor effects *per se* and/or to be immunologically compatible with the receiver (*i.e.*, matched HLA haplotype). APCs may generally be isolated from any of a variety of biological fluids and organs, including tumor and peritumoral tissues, and may be autologous, allogeneic, syngeneic or xenogeneic cells.

Certain preferred embodiments of the present invention use dendritic cells or progenitors thereof as antigen-presenting cells. Dendritic cells are highly potent APCs (Banchereau and Steinman, *Nature* 392:245-251, 1998) and have been shown to be effective as a physiological adjuvant for eliciting prophylactic or therapeutic antitumor immunity (*see* Timmerman and Levy, *Ann. Rev. Med.* 50:507-529, 1999). In general, dendritic cells may be identified based on their typical shape (stellate *in situ*,

with marked cytoplasmic processes (dendrites) visible *in vitro*), their ability to take up, process and present antigens with high efficiency and their ability to activate naïve T cell responses. Dendritic cells may, of course, be engineered to express specific cell-surface receptors or ligands that are not commonly found on dendritic cells *in vivo* or *ex vivo*, and such modified dendritic cells are contemplated by the present invention. As an alternative to dendritic cells, secreted vesicles antigen-loaded dendritic cells (called exosomes) may be used within a vaccine (*see* Zitvogel et al., *Nature Med.* 4:594-600, 1998).

Dendritic cells and progenitors may be obtained from peripheral blood, bone marrow, tumor-infiltrating cells, peritumoral tissues-infiltrating cells, lymph nodes, spleen, skin, umbilical cord blood or any other suitable tissue or fluid. For example, dendritic cells may be differentiated *ex vivo* by adding a combination of cytokines such as GM-CSF, IL-4, IL-13 and/or TNF $\alpha$  to cultures of monocytes harvested from peripheral blood. Alternatively, CD34 positive cells harvested from peripheral blood, umbilical cord blood or bone marrow may be differentiated into dendritic cells by adding to the culture medium combinations of GM-CSF, IL-3, TNF $\alpha$ , CD40 ligand, LPS, flt3 ligand and/or other compound(s) that induce differentiation, maturation and proliferation of dendritic cells.

Dendritic cells are conveniently categorized as "immature" and "mature" cells, which allows a simple way to discriminate between two well characterized phenotypes. However, this nomenclature should not be construed to exclude all possible intermediate stages of differentiation. Immature dendritic cells are characterized as APC with a high capacity for antigen uptake and processing, which correlates with the high expression of Fc $\gamma$  receptor and mannose receptor. The mature phenotype is typically characterized by a lower expression of these markers, but a high expression of cell surface molecules responsible for T cell activation such as class I and class II MHC, adhesion molecules (*e.g.*, CD54 and CD11) and costimulatory molecules (*e.g.*, CD40, CD80, CD86 and 4-1BB).

APCs may generally be transfected with a polynucleotide encoding a breast tumor protein (or portion or other variant thereof) such that the breast tumor polypeptide, or an immunogenic portion thereof, is expressed on the cell surface. Such

transfection may take place *ex vivo*, and a composition or vaccine comprising such transfected cells may then be used for therapeutic purposes, as described herein. Alternatively, a gene delivery vehicle that targets a dendritic or other antigen presenting cell may be administered to a patient, resulting in transfection that occurs *in vivo*. *In vivo* and *ex vivo* transfection of dendritic cells, for example, may generally be performed using any methods known in the art, such as those described in WO 97/24447, or the gene gun approach described by Mahvi et al., *Immunology and cell Biology* 75:456-460, 1997. Antigen loading of dendritic cells may be achieved by incubating dendritic cells or progenitor cells with the breast tumor polypeptide, DNA (naked or within a plasmid vector) or RNA; or with antigen-expressing recombinant bacterium or viruses (e.g., vaccinia, fowlpox, adenovirus or lentivirus vectors). Prior to loading, the polypeptide may be covalently conjugated to an immunological partner that provides T cell help (e.g., a carrier molecule). Alternatively, a dendritic cell may be pulsed with a non-conjugated immunological partner, separately or in the presence of the polypeptide.

Vaccines and pharmaceutical compositions may be presented in unit-dose or multi-dose containers, such as sealed ampoules or vials. Such containers are preferably hermetically sealed to preserve sterility of the formulation until use. In general, formulations may be stored as suspensions, solutions or emulsions in oily or aqueous vehicles. Alternatively, a vaccine or pharmaceutical composition may be stored in a freeze-dried condition requiring only the addition of a sterile liquid carrier immediately prior to use.

#### CANCER THERAPY

In further aspects of the present invention, the compositions described herein may be used for immunotherapy of cancer, such as breast cancer. Within such methods, pharmaceutical compositions and vaccines are typically administered to a patient. As used herein, a "patient" refers to any warm-blooded animal, preferably a human. A patient may or may not be afflicted with cancer. Accordingly, the above pharmaceutical compositions and vaccines may be used to prevent the development of a cancer or to treat a patient afflicted with a cancer. A cancer may be diagnosed using

criteria generally accepted in the art, including the presence of a malignant tumor. Pharmaceutical compositions and vaccines may be administered either prior to or following surgical removal of primary tumors and/or treatment such as administration of radiotherapy or conventional chemotherapeutic drugs. Administration may be by any  
5 suitable method, including administration by intravenous, intraperitoneal, intramuscular, subcutaneous, intranasal, intradermal, anal, vaginal, topical and oral routes.

Within certain embodiments, immunotherapy may be active immunotherapy, in which treatment relies on the *in vivo* stimulation of the endogenous  
10 host immune system to react against tumors with the administration of immune response-modifying agents (such as polypeptides and polynucleotides as provided herein).

Within other embodiments, immunotherapy may be passive immunotherapy, in which treatment involves the delivery of agents with established  
15 tumor-immune reactivity (such as effector cells or antibodies) that can directly or indirectly mediate antitumor effects and does not necessarily depend on an intact host immune system. Examples of effector cells include T cells as discussed above, T lymphocytes (such as CD8<sup>+</sup> cytotoxic T lymphocytes and CD4<sup>+</sup> T-helper tumor-infiltrating lymphocytes), killer cells (such as Natural Killer cells and lymphokine-activated killer cells), B cells and antigen-presenting cells (such as dendritic cells and  
20 macrophages) expressing a polypeptide provided herein. T cell receptors and antibody receptors specific for the polypeptides recited herein may be cloned, expressed and transferred into other vectors or effector cells for adoptive immunotherapy. The polypeptides provided herein may also be used to generate antibodies or anti-idiotypic  
25 antibodies (as described above and in U.S. Patent No. 4,918,164) for passive immunotherapy.

Effector cells may generally be obtained in sufficient quantities for adoptive immunotherapy by growth *in vitro*, as described herein. Culture conditions for expanding single antigen-specific effector cells to several billion in number with  
30 retention of antigen recognition *in vivo* are well known in the art. Such *in vitro* culture conditions typically use intermittent stimulation with antigen, often in the presence of

cytokines (such as IL-2) and non-dividing feeder cells. As noted above, immunoreactive polypeptides as provided herein may be used to rapidly expand antigen-specific T cell cultures in order to generate a sufficient number of cells for immunotherapy. In particular, antigen-presenting cells, such as dendritic, macrophage, monocyte, fibroblast and/or B cells, may be pulsed with immunoreactive polypeptides or transfected with one or more polynucleotides using standard techniques well known in the art. For example, antigen-presenting cells can be transfected with a polynucleotide having a promoter appropriate for increasing expression in a recombinant virus or other expression system. Cultured effector cells for use in therapy must be able to grow and distribute widely, and to survive long term *in vivo*. Studies have shown that cultured effector cells can be induced to grow *in vivo* and to survive long term in substantial numbers by repeated stimulation with antigen supplemented with IL-2 (*see*, for example, Cheever et al., *Immunological Reviews* 157:177, 1997).

Alternatively, a vector expressing a polypeptide recited herein may be introduced into antigen presenting cells taken from a patient and clonally propagated *ex vivo* for transplant back into the same patient. Transfected cells may be reintroduced into the patient using any means known in the art, preferably in sterile form by intravenous, intracavitary, intraperitoneal or intratumor administration.

Routes and frequency of administration of the therapeutic compositions described herein, as well as dosage, will vary from individual to individual, and may be readily established using standard techniques. In general, the pharmaceutical compositions and vaccines may be administered by injection (*e.g.*, intracutaneous, intramuscular, intravenous or subcutaneous), intranasally (*e.g.*, by aspiration) or orally. Preferably, between 1 and 10 doses may be administered over a 52 week period. Preferably, 6 doses are administered, at intervals of 1 month, and booster vaccinations may be given periodically thereafter. Alternate protocols may be appropriate for individual patients. A suitable dose is an amount of a compound that, when administered as described above, is capable of promoting an anti-tumor immune response, and is at least 10-50% above the basal (*i.e.*, untreated) level. Such response can be monitored by measuring the anti-tumor antibodies in a patient or by vaccine-dependent generation of cytolytic effector cells capable of killing the patient's tumor



cells *in vitro*. Such vaccines should also be capable of causing an immune response that leads to an improved clinical outcome (*e.g.*, more frequent remissions, complete or partial or longer disease-free survival) in vaccinated patients as compared to non-vaccinated patients. In general, for pharmaceutical compositions and vaccines  
5 comprising one or more polypeptides, the amount of each polypeptide present in a dose ranges from about 25 µg to 5 mg per kg of host. Suitable dose sizes will vary with the size of the patient, but will typically range from about 0.1 mL to about 5 mL.

In general, an appropriate dosage and treatment regimen provides the active compound(s) in an amount sufficient to provide therapeutic and/or prophylactic  
10 benefit. Such a response can be monitored by establishing an improved clinical outcome (*e.g.*, more frequent remissions, complete or partial, or longer disease-free survival) in treated patients as compared to non-treated patients. Increases in preexisting immune responses to a breast tumor protein generally correlate with an improved clinical outcome. Such immune responses may generally be evaluated using  
15 standard proliferation, cytotoxicity or cytokine assays, which may be performed using samples obtained from a patient before and after treatment.

#### CANCER DETECTION AND DIAGNOSIS

In general, a cancer may be detected in a patient based on the presence of one or more breast tumor proteins and/or polynucleotides encoding such proteins in a  
20 biological sample (for example, blood, sera, sputum urine and/or tumor biopsies) obtained from the patient. In other words, such proteins may be used as markers to indicate the presence or absence of a cancer such as breast cancer. In addition, such proteins may be useful for the detection of other cancers. The binding agents provided herein generally permit detection of the level of antigen that binds to the agent in the  
25 biological sample. Polynucleotide primers and probes may be used to detect the level of mRNA encoding a tumor protein, which is also indicative of the presence or absence of a cancer. In general, a breast tumor sequence should be present at a level that is at least three fold higher in tumor tissue than in normal tissue

There are a variety of assay formats known to those of ordinary skill in  
30 the art for using a binding agent to detect polypeptide markers in a sample. *See, e.g.*,

Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. In general, the presence or absence of a cancer in a patient may be determined by (a) contacting a biological sample obtained from a patient with a binding agent; (b) detecting in the sample a level of polypeptide that binds to the binding agent; and (c) 5 comparing the level of polypeptide with a predetermined cut-off value.

In a preferred embodiment, the assay involves the use of binding agent immobilized on a solid support to bind to and remove the polypeptide from the remainder of the sample. The bound polypeptide may then be detected using a detection reagent that contains a reporter group and specifically binds to the binding 10 agent/polypeptide complex. Such detection reagents may comprise, for example, a binding agent that specifically binds to the polypeptide or an antibody or other agent that specifically binds to the binding agent, such as an anti-immunoglobulin, protein G, protein A or a lectin. Alternatively, a competitive assay may be utilized, in which a polypeptide is labeled with a reporter group and allowed to bind to the immobilized 15 binding agent after incubation of the binding agent with the sample. The extent to which components of the sample inhibit the binding of the labeled polypeptide to the binding agent is indicative of the reactivity of the sample with the immobilized binding agent. Suitable polypeptides for use within such assays include full length breast tumor proteins and portions thereof to which the binding agent binds, as described above.

20 The solid support may be any material known to those of ordinary skill in the art to which the tumor protein may be attached. For example, the solid support may be a test well in a microtiter plate or a nitrocellulose or other suitable membrane. Alternatively, the support may be a bead or disc, such as glass, fiberglass, latex or a plastic material such as polystyrene or polyvinylchloride. The support may also be a 25 magnetic particle or a fiber optic sensor, such as those disclosed, for example, in U.S. Patent No. 5,359,681. The binding agent may be immobilized on the solid support using a variety of techniques known to those of skill in the art, which are amply described in the patent and scientific literature. In the context of the present invention, the term "immobilization" refers to both noncovalent association, such as adsorption, 30 and covalent attachment (which may be a direct linkage between the agent and functional groups on the support or may be a linkage by way of a cross-linking agent).

Immobilization by adsorption to a well in a microtiter plate or to a membrane is preferred. In such cases, adsorption may be achieved by contacting the binding agent, in a suitable buffer, with the solid support for a suitable amount of time. The contact time varies with temperature, but is typically between about 1 hour and about 1 day. In  
5 general, contacting a well of a plastic microtiter plate (such as polystyrene or polyvinylchloride) with an amount of binding agent ranging from about 10 ng to about 10  $\mu$ g, and preferably about 100 ng to about 1  $\mu$ g, is sufficient to immobilize an adequate amount of binding agent.

Covalent attachment of binding agent to a solid support may generally be  
10 achieved by first reacting the support with a bifunctional reagent that will react with both the support and a functional group, such as a hydroxyl or amino group, on the binding agent. For example, the binding agent may be covalently attached to supports having an appropriate polymer coating using benzoquinone or by condensation of an aldehyde group on the support with an amine and an active hydrogen on the binding  
15 partner (*see, e.g.,* Pierce Immunotechnology Catalog and Handbook, 1991, at A12-A13).

In certain embodiments, the assay is a two-antibody sandwich assay. This assay may be performed by first contacting an antibody that has been immobilized on a solid support, commonly the well of a microtiter plate, with the sample, such that  
20 polypeptides within the sample are allowed to bind to the immobilized antibody. Unbound sample is then removed from the immobilized polypeptide-antibody complexes and a detection reagent (preferably a second antibody capable of binding to a different site on the polypeptide) containing a reporter group is added. The amount of detection reagent that remains bound to the solid support is then determined using a  
25 method appropriate for the specific reporter group.

More specifically, once the antibody is immobilized on the support as described above, the remaining protein binding sites on the support are typically blocked. Any suitable blocking agent known to those of ordinary skill in the art, such as bovine serum albumin or Tween 20<sup>TM</sup> (Sigma Chemical Co., St. Louis, MO). The  
30 immobilized antibody is then incubated with the sample, and polypeptide is allowed to bind to the antibody. The sample may be diluted with a suitable diluent, such as

phosphate-buffered saline (PBS) prior to incubation. In general, an appropriate contact time (*i.e.*, incubation time) is a period of time that is sufficient to detect the presence of polypeptide within a sample obtained from an individual with breast cancer. Preferably, the contact time is sufficient to achieve a level of binding that is at least about 95% of that achieved at equilibrium between bound and unbound polypeptide. Those of ordinary skill in the art will recognize that the time necessary to achieve equilibrium may be readily determined by assaying the level of binding that occurs over a period of time. At room temperature, an incubation time of about 30 minutes is generally sufficient.

10               Unbound sample may then be removed by washing the solid support with an appropriate buffer, such as PBS containing 0.1% Tween 20™. The second antibody, which contains a reporter group, may then be added to the solid support. Preferred reporter groups include those groups recited above.

                  The detection reagent is then incubated with the immobilized antibody-polypeptide complex for an amount of time sufficient to detect the bound polypeptide. An appropriate amount of time may generally be determined by assaying the level of binding that occurs over a period of time. Unbound detection reagent is then removed and bound detection reagent is detected using the reporter group. The method employed for detecting the reporter group depends upon the nature of the reporter group. For radioactive groups, scintillation counting or autoradiographic methods are generally appropriate. Spectroscopic methods may be used to detect dyes, luminescent groups and fluorescent groups. Biotin may be detected using avidin, coupled to a different reporter group (commonly a radioactive or fluorescent group or an enzyme). Enzyme reporter groups may generally be detected by the addition of substrate (generally for a specific period of time), followed by spectroscopic or other analysis of the reaction products.

                  To determine the presence or absence of a cancer, such as breast cancer, the signal detected from the reporter group that remains bound to the solid support is generally compared to a signal that corresponds to a predetermined cut-off value. In one preferred embodiment, the cut-off value for the detection of a cancer is the average mean signal obtained when the immobilized antibody is incubated with samples from

patients without the cancer. In general, a sample generating a signal that is three standard deviations above the predetermined cut-off value is considered positive for the cancer. In an alternate preferred embodiment, the cut-off value is determined using a Receiver Operator Curve, according to the method of Sackett et al., *Clinical*  
5 *Epidemiology: A Basic Science for Clinical Medicine*, Little Brown and Co., 1985, p. 106-7. Briefly, in this embodiment, the cut-off value may be determined from a plot of pairs of true positive rates (*i.e.*, sensitivity) and false positive rates (100%-specificity) that correspond to each possible cut-off value for the diagnostic test result. The cut-off value on the plot that is the closest to the upper left-hand corner (*i.e.*, the value that  
10 encloses the largest area) is the most accurate cut-off value, and a sample generating a signal that is higher than the cut-off value determined by this method may be considered positive. Alternatively, the cut-off value may be shifted to the left along the plot, to minimize the false positive rate, or to the right, to minimize the false negative rate. In general, a sample generating a signal that is higher than the cut-off value determined by  
15 this method is considered positive for a cancer.

In a related embodiment, the assay is performed in a flow-through or strip test format, wherein the binding agent is immobilized on a membrane, such as nitrocellulose. In the flow-through test, polypeptides within the sample bind to the immobilized binding agent as the sample passes through the membrane. A second,  
20 labeled binding agent then binds to the binding agent-polypeptide complex as a solution containing the second binding agent flows through the membrane. The detection of bound second binding agent may then be performed as described above. In the strip test format, one end of the membrane to which binding agent is bound is immersed in a solution containing the sample. The sample migrates along the membrane through a  
25 region containing second binding agent and to the area of immobilized binding agent. Concentration of second binding agent at the area of immobilized antibody indicates the presence of a cancer. Typically, the concentration of second binding agent at that site generates a pattern, such as a line, that can be read visually. The absence of such a pattern indicates a negative result. In general, the amount of binding agent immobilized  
30 on the membrane is selected to generate a visually discernible pattern when the biological sample contains a level of polypeptide that would be sufficient to generate a

positive signal in the two-antibody sandwich assay, in the format discussed above. Preferred binding agents for use in such assays are antibodies and antigen-binding fragments thereof. Preferably, the amount of antibody immobilized on the membrane ranges from about 25 ng to about 1 $\mu$ g, and more preferably from about 50 ng to about 500 ng. Such tests can typically be performed with a very small amount of biological sample.

Of course, numerous other assay protocols exist that are suitable for use with the tumor proteins or binding agents of the present invention. The above descriptions are intended to be exemplary only. For example, it will be apparent to those of ordinary skill in the art that the above protocols may be readily modified to use breast tumor polypeptides to detect antibodies that bind to such polypeptides in a biological sample. The detection of such breast tumor protein specific antibodies may correlate with the presence of a cancer.

A cancer may also, or alternatively, be detected based on the presence of T cells that specifically react with a breast tumor protein in a biological sample. Within certain methods, a biological sample comprising CD4<sup>+</sup> and/or CD8<sup>+</sup> T cells isolated from a patient is incubated with a breast tumor polypeptide, a polynucleotide encoding such a polypeptide and/or an APC that expresses at least an immunogenic portion of such a polypeptide, and the presence or absence of specific activation of the T cells is detected. Suitable biological samples include, but are not limited to, isolated T cells. For example, T cells may be isolated from a patient by routine techniques (such as by Ficoll/Hypaque density gradient centrifugation of peripheral blood lymphocytes). T cells may be incubated *in vitro* for 2-9 days (typically 4 days) at 37°C with polypeptide (e.g., 5 - 25  $\mu$ g/ml). It may be desirable to incubate another aliquot of a T cell sample in the absence of breast tumor polypeptide to serve as a control. For CD4<sup>+</sup> T cells, activation is preferably detected by evaluating proliferation of the T cells. For CD8<sup>+</sup> T cells, activation is preferably detected by evaluating cytolytic activity. A level of proliferation that is at least two fold greater and/or a level of cytolytic activity that is at least 20% greater than in disease-free patients indicates the presence of a cancer in the patient.

As noted above, a cancer may also, or alternatively, be detected based on the level of mRNA encoding a breast tumor protein in a biological sample. For example, at least two oligonucleotide primers may be employed in a polymerase chain reaction (PCR) based assay to amplify a portion of a breast tumor cDNA derived from a biological sample, wherein at least one of the oligonucleotide primers is specific for (i.e., hybridizes to) a polynucleotide encoding the breast tumor protein. The amplified cDNA is then separated and detected using techniques well known in the art, such as gel electrophoresis. Similarly, oligonucleotide probes that specifically hybridize to a polynucleotide encoding a breast tumor protein may be used in a hybridization assay to detect the presence of polynucleotide encoding the tumor protein in a biological sample.

To permit hybridization under assay conditions, oligonucleotide primers and probes should comprise an oligonucleotide sequence that has at least about 60%, preferably at least about 75% and more preferably at least about 90%, identity to a portion of a polynucleotide encoding a breast tumor protein that is at least 10 nucleotides, and preferably at least 20 nucleotides, in length. Preferably, oligonucleotide primers and/or probes hybridize to a polynucleotide encoding a polypeptide described herein under moderately stringent conditions, as defined above. Oligonucleotide primers and/or probes which may be usefully employed in the diagnostic methods described herein preferably are at least 10-40 nucleotides in length. In a preferred embodiment, the oligonucleotide primers comprise at least 10 contiguous nucleotides, more preferably at least 15 contiguous nucleotides, of a DNA molecule having a sequence recited in SEQ ID NO: 1-38, 42-204, 205, 207 and 210-290. Techniques for both PCR based assays and hybridization assays are well known in the art (see, for example, Mullis et al., *Cold Spring Harbor Symp. Quant. Biol.*, 51:263, 1987; Erlich ed., *PCR Technology*, Stockton Press, NY, 1989).

One preferred assay employs RT-PCR, in which PCR is applied in conjunction with reverse transcription. Typically, RNA is extracted from a biological sample, such as biopsy tissue, and is reverse transcribed to produce cDNA molecules. PCR amplification using at least one specific primer generates a cDNA molecule, which may be separated and visualized using, for example, gel electrophoresis. Amplification may be performed on biological samples taken from a test patient and from an

individual who is not afflicted with a cancer. The amplification reaction may be performed on several dilutions of cDNA spanning two orders of magnitude. A two-fold or greater increase in expression in several dilutions of the test patient sample as compared to the same dilutions of the non-cancerous sample is typically considered  
5 positive.

In another embodiment, the compositions described herein may be used as markers for the progression of cancer. In this embodiment, assays as described above for the diagnosis of a cancer may be performed over time, and the change in the level of reactive polypeptide(s) or polynucleotide(s) evaluated. For example, the assays  
10 may be performed every 24-72 hours for a period of 6 months to 1 year, and thereafter performed as needed. In general, a cancer is progressing in those patients in whom the level of polypeptide or polynucleotide detected increases over time. In contrast, the cancer is not progressing when the level of reactive polypeptide or polynucleotide either remains constant or decreases with time.

Certain *in vivo* diagnostic assays may be performed directly on a tumor.  
15 One such assay involves contacting tumor cells with a binding agent. The bound binding agent may then be detected directly or indirectly via a reporter group. Such binding agents may also be used in histological applications. Alternatively, polynucleotide probes may be used within such applications.

As noted above, to improve sensitivity, multiple breast tumor protein  
20 markers may be assayed within a given sample. It will be apparent that binding agents specific for different proteins provided herein may be combined within a single assay. Further, multiple primers or probes may be used concurrently. The selection of tumor protein markers may be based on routine experiments to determine combinations that  
25 results in optimal sensitivity. In addition, or alternatively, assays for tumor proteins provided herein may be combined with assays for other known tumor antigens.

#### **DIAGNOSTIC KITS**

The present invention further provides kits for use within any of the above diagnostic methods. Such kits typically comprise two or more components  
30 necessary for performing a diagnostic assay. Components may be compounds,



reagents, containers and/or equipment. For example, one container within a kit may contain a monoclonal antibody or fragment thereof that specifically binds to a breast tumor protein. Such antibodies or fragments may be provided attached to a support material, as described above. One or more additional containers may enclose elements,  
5 such as reagents or buffers, to be used in the assay. Such kits may also, or alternatively, contain a detection reagent as described above that contains a reporter group suitable for direct or indirect detection of antibody binding.

Alternatively, a kit may be designed to detect the level of mRNA encoding a breast tumor protein in a biological sample. Such kits generally comprise at  
10 least one oligonucleotide probe or primer, as described above, that hybridizes to a polynucleotide encoding a breast tumor protein. Such an oligonucleotide may be used, for example, within a PCR or hybridization assay. Additional components that may be present within such kits include a second oligonucleotide and/or a diagnostic reagent or container to facilitate the detection of a polynucleotide encoding a breast tumor protein.

15 The following Examples are offered by way of illustration and not by way of limitation.

## EXAMPLE 1

IDENTIFICATION OF BREAST TUMOR PROTEIN cDNAS USING  
SUBTRACTION METHODOLOGY

This Example illustrates the identification of cDNA molecules encoding  
5 breast tumor proteins.

A human metastatic breast tumor cDNA expression library was constructed from metastatic breast tumor poly A<sup>+</sup> RNA using a Superscript Plasmid System for cDNA Synthesis and Plasmid Cloning kit (BRL Life Technologies, Gaithersburg, MD 20897) following the manufacturer's protocol. Specifically, breast  
10 tumor tissues were homogenized with polytron (Kinematica, Switzerland) and total RNA was extracted using Trizol reagent (BRL Life Technologies) as directed by the manufacturer. The poly A<sup>+</sup> RNA was then purified using a Qiagen oligotex spin column mRNA purification kit (Qiagen, Santa Clarita, CA 91355) according to the manufacturer's protocol. First-strand cDNA was synthesized using the NotI/Oligo-  
15 dT18 primer. Double-stranded cDNA was synthesized, ligated with EcoRI/BstX I adaptors (Invitrogen, Carlsbad, CA) and digested with NotI. Following size fractionation with Chroma Spin-1000 columns (Clontech, Palo Alto, CA 94303), the cDNA was ligated into the EcoRI/NotI site of pCDNA3.1 (Invitrogen, Carlsbad, CA) and transformed into ElectroMax *E. coli* DH10B cells (BRL Life Technologies) by  
20 electroporation.

Using the same procedure, a normal human breast cDNA expression library was prepared from a pool of four normal breast tissue specimens. The cDNA libraries were characterized by determining the number of independent colonies, the percentage of clones that carried insert, the average insert size and by sequence analysis.  
25 Sequencing analysis showed both libraries to contain good complex cDNA clones that were synthesized from mRNA, with minimal rRNA and mitochondrial DNA contamination sequencing.

A cDNA subtracted library (referred to as BS3) was prepared using the above metastatic breast tumor and normal breast cDNA libraries, as described by Hara  
30 *et al.* (*Blood*, 84:189-199, 1994) with some modifications. Specifically, a breast tumor-

specific subtracted cDNA library was generated as follows. Normal breast cDNA library (70 µg) was digested with EcoRI, NotI, and SfuI, followed by a filling-in reaction with DNA polymerase Klenow fragment. After phenol-chloroform extraction and ethanol precipitation, the DNA was dissolved in 100 µl of H<sub>2</sub>O, heat-denatured and  
5 mixed with 100 µl (100 µg) of Photoprobe biotin (Vector Laboratories, Burlingame, CA), the resulting mixture was irradiated with a 270 W sunlamp on ice for 20 minutes. Additional Photoprobe biotin (50 µl) was added and the biotinylation reaction was repeated. After extraction with butanol five times, the DNA was ethanol-precipitated and dissolved in 23 µl H<sub>2</sub>O to form the driver DNA.

10 To form the tracer DNA, 10 µg breast tumor cDNA library was digested with BamHI and XhoI, phenol chloroform extracted and passed through Chroma spin-400 columns (Clontech). Following ethanol precipitation, the tracer DNA was dissolved in 5 µl H<sub>2</sub>O. Tracer DNA was mixed with 15 µl driver DNA and 20 µl of 2 x hybridization buffer (1.5 M NaCl/10 mM EDTA/50 mM HEPES pH 7.5/0.2% sodium  
15 dodecyl sulfate), overlaid with mineral oil, and heat-denatured completely. The sample was immediately transferred into a 68 °C water bath and incubated for 20 hours (long hybridization [LH]). The reaction mixture was then subjected to a streptavidin treatment followed by phenol/chloroform extraction. This process was repeated three more times. Subtracted DNA was precipitated, dissolved in 12 µl H<sub>2</sub>O, mixed with 8 µl  
20 driver DNA and 20 µl of 2 x hybridization buffer, and subjected to a hybridization at 68 °C for 2 hours (short hybridization [SH]). After removal of biotinylated double-stranded DNA, subtracted cDNA was ligated into BamHI/XhoI site of chloramphenicol resistant pBCSK<sup>+</sup> (Stratagene, La Jolla, CA 92037) and transformed into ElectroMax *E. coli* DH10B cells by electroporation to generate a breast tumor specific subtracted  
25 cDNA library.

To analyze the subtracted cDNA library, plasmid DNA was prepared from independent clones, randomly picked from the subtracted breast tumor specific library and characterized by DNA sequencing with a Perkin Elmer/Applied Biosystems Division Automated Sequencer Model 373A (Foster City, CA).

30 A second cDNA subtraction library containing cDNA from breast tumor subtracted with normal breast cDNA, and known as BT, was constructed as follows.

Total RNA was extracted from primary breast tumor tissues using Trizol reagent (Gibco BRL Life Technologies, Gaithersburg, MD) as described by the manufacturer. The polyA<sup>+</sup> RNA was purified using an oligo(dT) cellulose column according to standard protocols. First strand cDNA was synthesized using the primer supplied in a Clontech  
5 PCR-Select cDNA Subtraction Kit (Clontech, Palo Alto, CA). The driver DNA consisted of cDNAs from two normal breast tissues with the tester cDNA being from three primary breast tumors. Double-stranded cDNA was synthesized for both tester and driver, and digested with a combination of endonucleases (MluI, MscI, PvuII, Sall and StuI) which recognize six base pairs DNA. This modification increased the average  
10 cDNA size dramatically compared with cDNAs generated according to the protocol of Clontech. The digested tester cDNAs were ligated to two different adaptors and the subtraction was performed according to Clontech's protocol. The subtracted cDNAs were subjected to two rounds of PCR amplification, following the manufacturer's protocol. The resulting PCR products were subcloned into the TA cloning vector,  
15 pCRII (Invitrogen, San Diego, CA) and transformed into ElectroMax *E. coli* DH10B cells (Gibco BRL Life, Technologies) by electroporation. DNA was isolated from independent clones and sequenced using a Perkin Elmer/Applied Biosystems Division (Foster City, CA) Automated Sequencer Model 373A.

Two additional subtracted cDNA libraries were prepared from cDNA  
20 from breast tumors subtracted with a pool of cDNA from six normal tissues (liver, brain, stomach, small intestine, kidney and heart; referred to as 2BT and BC6) using the PCR-subtraction protocol of Clontech, described above. A fourth subtracted library (referred to as Bt-Met) was prepared using the protocol of Clontech from cDNA from metastatic breast tumors subtracted with cDNA from five normal tissues (brain, lung,  
25 PBMC, pancreas and normal breast).

cDNA clones isolated in the breast subtractions BS3, BT, 2BT, BC6 and BT-Met, described above, were colony PCR amplified and their mRNA expression levels in breast tumor, normal breast and various other normal tissues were determined using microarray technology. Briefly, the PCR amplification products were dotted onto  
30 slides in an array format, with each product occupying a unique location in the array. mRNA was extracted from the tissue sample to be tested, reverse transcribed, and

fluorescent-labeled cDNA probes were generated. The microarrays were probed with the labeled cDNA probes, the slides scanned and fluorescence intensity was measured. This intensity correlates with the hybridization intensity.

The determined cDNA sequences of 131 clones determined to be over-expressed in breast tumor tissue compared to other tissues tested by a visual analysis of the microarray data are provided in SEQ ID NO: 1-35 and 42-137. Comparison of these cDNA sequences with known sequences in the gene bank using the EMBL and GenBank databases revealed no significant homologies to the sequences provided in SEQ ID NO: 7, 10, 21, 26, 30, 63, 81 and 104. The sequences of SEQ ID NO: 2-5, 8, 9, 13, 15, 16, 22, 25, 27, 28, 33, 35, 72, 73, 103, 107, 109, 118, 128, 129 134 and 136 showed some homology to previously isolated expressed sequences tags (ESTs), while the sequences of SEQ ID NO: 1, 6, 11, 12, 14, 17-20, 23, 24, 29, 31, 32, 34, 42-62, 64-71, 74-80, 82-102, 105, 106, 108, 110-117, 119-127, 130-133, 135 and 137 showed some homology to previously identified genes.

The determined cDNA sequences of an additional 45 clones isolated from the BT-Met library as described above and found to be over-expressed in breast tumors and metastatic breast tumors compared to other tissues tested, are provided in SEQ ID NO: 138-182. Comparison of the sequences of SEQ ID NO: 159-161, 164 and 181 revealed no significant homologies to previously identified sequences. The sequences of SEQ ID NO: 138-158, 162, 163, 165-180 and 182 showed some homology to previously identified genes.

In further studies, suppression subtractive hybridization (Clontech) was preformed using a pool of cDNA from 3 unique human breast tumors as the tester and a pool of cDNA from 6 other normal human tissues (liver, brain, stomach, small intestine, heart and kidney) as the driver. The isolated cDNA fragments were subcloned and characterized by DNA sequencing. The determined cDNA sequences of 22 isolated clones are provided in SEQ ID NO: 183-204. Comparison of these sequences with those in the public databases revealed no significant homologies to previously identified sequences.

The determined cDNA sequences of 71 additional breast-specific genes isolated during characterization of breast tumor cDNA libraries are provided in SEQ ID

NO: 210-290. Comparison of these sequences with those in the GenBank and Geneseq databases revealed no significant homologies.

## EXAMPLE 2

### 5     IDENTIFICATION OF BREAST TUMOR PROTEIN cDNAS BY RT-PCR

GABA<sub>A</sub> receptor clones were isolated from human breast cancer cDNA libraries by first preparing cDNA libraries from breast tumor samples from different patients as described above. PCR primers were designed based on the GABA<sub>A</sub> receptor subunit sequences described by Hedblom and Kirkness (*Jnl. Biol. Chem.* 272:15346-10 15350, 1997) and used to amplify sequences from the breast tumor cDNA libraries by RT-PCR. The determined cDNA sequences of three GABA<sub>A</sub> receptor clones are provided in SEQ ID NO: 36-38, with the corresponding amino acid sequences being provided in SEQ ID NO: 39-41.

The clone with the longest open reading frame (ORF; SEQ ID NO: 36) 15 showed homology to the GABA<sub>A</sub> receptor of Hedblom and Kirkness, with four potential transmembrane regions at the C-terminal part of the protein, while the clones of SEQ ID NO: 37 and 38 retained either no transmembrane region or only the first transmembrane region. Some patients were found to have only the clones with the shorter ORFs while others had both the clones with longer and shorter ORFs.

20

## EXAMPLE 3

### EXPRESSION OF OVARIAN TUMOR-DERIVED ANTIGENS

#### IN BREAST

Isolation of the antigens O772P and O8E from ovarian tumor tissue is 25 described in US Patent Application No. 09/338,933, filed June 23, 1999. The determined cDNA sequence for O772P is provided in SEQ ID NO: 205, with the corresponding amino acid sequence being provided in SEQ ID NO: 206. The full-length cDNA sequence for O8E is provided in SEQ ID NO: 207. Two protein sequences may be translated from the full length O8E. Form "A" (SEQ ID NO: 208)

begins with a putative start methionine. A second form "B" (SEQ ID NO: 209) includes 27 additional upstream residues to the 5' end of the nucleotide sequence.

The expression levels of O772P and O8E in a variety of tumor and normal tissues, including metastatic breast tumors, were analyzed by real time PCR. Both genes were found to have increased mRNA expression in 30-50% of breast tumors. For O772P, elevated expression was also observed in normal trachea, ureter, uterus and ovary. For O8E, elevated expression was also observed in normal trachea, kidney and ovary. Additional analysis employing a panel of tumor cell lines demonstrated increased expression of O8E in the breast tumor cell lines SKBR3, MDA-MB-415 and BT474, and increased expression of O772P in SKBR3. Collectively, the data indicate that O772P and O8E may be useful in the diagnosis and therapy of breast cancer.

#### EXAMPLE 4

##### SYNTHESIS OF POLYPEPTIDES

Polypeptides may be synthesized on a Perkin Elmer/Applied Biosystems Division 430A peptide synthesizer using Fmoc chemistry with HPTU (O-Benzotriazole-N,N,N',N'-tetramethyluronium hexafluorophosphate) activation. A Gly-Cys-Gly sequence may be attached to the amino terminus of the peptide to provide a method of conjugation, binding to an immobilized surface, or labeling of the peptide. Cleavage of the peptides from the solid support may be carried out using the following cleavage mixture: trifluoroacetic acid:ethanedithiol:thioanisole:water:phenol (40:1:2:2:3). After cleaving for 2 hours, the peptides may be precipitated in cold methyl-t-butyl-ether. The peptide pellets may then be dissolved in water containing 0.1% trifluoroacetic acid (TFA) and lyophilized prior to purification by C18 reverse phase HPLC. A gradient of 0%-60% acetonitrile (containing 0.1% TFA) in water (containing 0.1% TFA) may be used to elute the peptides. Following lyophilization of the pure fractions, the peptides may be characterized using electrospray or other types of mass spectrometry and by amino acid analysis.

From the foregoing it will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.



## CLAIMS

1. An isolated polypeptide, comprising at least an immunogenic portion of a breast tumor protein, or a variant thereof that differs in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with antigen-specific antisera is not substantially diminished, wherein the tumor protein comprises an amino acid sequence that is encoded by a polynucleotide sequence selected from the group consisting of:

(a) sequences recited in SEQ ID NOs: 2-5, 7, 10, 13, 15, 16, 21, 22, 25, 28, 30, 33, 35, 63, 72, 73, 81, 103, 104, 107, 109, 118, 128, 129, 134, 136, 159-161, 164, 181, 183-204 and 210-290;

(b) sequences that hybridize to a sequence recited in any one of SEQ ID NOs: 2-5, 7, 10, 13, 15, 16, 21, 22, 25, 28, 30, 33, 35, 63, 72, 73, 81, 103, 104, 107, 109, 118, 128, 129, 134, 136, 159-161, 164, 181, 183-204 and 210-290 under moderately stringent conditions; and

(c) complements of sequences of (a) or (b).

2. An isolated polypeptide according to claim 1, wherein the polypeptide comprises an amino acid sequence that is encoded by a polynucleotide sequence recited in any one of SEQ ID NOs: 2-5, 7, 10, 13, 15, 16, 21, 22, 25, 28, 30, 33, 35, 63, 72, 73, 81, 103, 104, 107, 109, 118, 128, 129, 134, 136, 159-161, 164, 181, 183-204 and 210-290 or a complement of any of the foregoing polynucleotide sequences.

3. An isolated polynucleotide encoding at least 15 amino acid residues of a breast tumor protein, or a variant thereof that differs in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with antigen-specific antisera is not substantially diminished, wherein the tumor protein comprises an amino acid sequence that is encoded by a polynucleotide comprising a sequence recited in any one of SEQ ID NOs: 2-5, 7, 10, 13, 15, 16, 21, 22,

25, 28, 30, 33, 35, 63, 72, 73, 81, 103, 104, 107, 109, 118, 128, 129, 134, 136, 159-161, 164, 181, 183-204 and 210-290 or a complement of any of the foregoing sequences.

4. An isolated polynucleotide encoding a breast tumor protein, or a variant thereof, wherein the tumor protein comprises an amino acid sequence that is encoded by a polynucleotide comprising a sequence recited in any one of SEQ ID NOs: 2-5, 7, 10, 13, 15, 16, 21, 22, 25, 28, 30, 33, 35, 63, 72, 73, 81, 103, 104, 107, 109, 118, 128, 129, 134, 136, 159-161, 164, 181, 183-204 and 210-290 or a complement of any of the foregoing sequences.

5. An isolated polynucleotide, comprising a sequence recited in any one of SEQ ID NOs: 2-5, 7, 10, 13, 15, 16, 21, 22, 25, 28, 30, 33, 35, 63, 72, 73, 81, 103, 104, 107, 109, 118, 128, 129, 134, 136, 159-161, 164, 181, 183-204 and 210-290.

6. An isolated polynucleotide, comprising a sequence that hybridizes to a sequence recited in any one of SEQ ID NOs: 2-5, 7, 10, 13, 15, 16, 21, 22, 25, 28, 30, 33, 35, 63, 72, 73, 81, 103, 104, 107, 109, 118, 128, 129, 134, 136, 159-161, 164, 181, 183-204 and 210-290 under moderately stringent conditions.

7. An isolated polynucleotide complementary to a polynucleotide according to any one of claims 3-6.

8. An expression vector, comprising a polynucleotide according to any one of claims 3-7.

9. A host cell transformed or transfected with an expression vector according to claim 8.

10. An isolated antibody, or antigen-binding fragment thereof, that specifically binds to a breast tumor protein that comprises an amino acid sequence that is encoded by a polynucleotide sequence recited in any one of SEQ ID NOs: 2-5, 7, 10, 13, 15, 16, 21, 22, 25, 28, 30, 33, 35, 63, 72, 73, 81, 103, 104, 107, 109, 118, 128, 129,

134, 136, 159-161, 164, 181, 183-204 and 210-290 or a complement of any of the foregoing polynucleotide sequences.

11. A fusion protein, comprising at least one polypeptide according to claim 1.

12. A fusion protein according to claim 11, wherein the fusion protein comprises an expression enhancer that increases expression of the fusion protein in a host cell transfected with a polynucleotide encoding the fusion protein.

13. A fusion protein according to claim 11, wherein the fusion protein comprises a T helper epitope that is not present within the polypeptide of claim 1.

14. A fusion protein according to claim 11, wherein the fusion protein comprises an affinity tag.

15. An isolated polynucleotide encoding a fusion protein according to claim 11.

16. A pharmaceutical composition, comprising a physiologically acceptable carrier and at least one component selected from the group consisting of:

- (a) a polypeptide according to claim 1;
- (b) a polynucleotide according to claim 3;
- (c) an antibody according to claim 10;
- (d) a fusion protein according to claim 11; and
- (e) a polynucleotide according to claim 15.

17. A vaccine comprising an immunostimulant and at least one component selected from the group consisting of:

- (a) a polypeptide according to claim 1;

- (b) a polynucleotide according to claim 3;
- (c) an antibody according to claim 10;
- (d) a fusion protein according to claim 11; and
- (e) a polynucleotide according to claim 15.

18. A vaccine according to claim 17, wherein the immunostimulant is an adjuvant.

19. A vaccine according to any claim 17, wherein the immunostimulant induces a predominantly Type I response.

20. A method for inhibiting the development of a cancer in a patient, comprising administering to a patient an effective amount of a pharmaceutical composition according to claim 16.

21. A method for inhibiting the development of a cancer in a patient, comprising administering to a patient an effective amount of a vaccine according to claim 17.

22. A pharmaceutical composition comprising an antigen-presenting cell that expresses a polypeptide according to claim 1, in combination with a pharmaceutically acceptable carrier or excipient.

23. A pharmaceutical composition according to claim 22, wherein the antigen presenting cell is a dendritic cell or a macrophage.

24. A vaccine comprising an antigen-presenting cell that expresses a polypeptide comprising at least an immunogenic portion of a breast tumor protein, or a variant thereof, wherein the tumor protein comprises an amino acid sequence that is encoded by a polynucleotide sequence selected from the group consisting of:

(a) sequences recited in SEQ ID NOs: 1-38, 42-204, 205, 207 and 210-290;

(b) sequences that hybridize to a sequence recited in any one of SEQ ID NOs: 1-38, 42-204, 205, 207 and 210-290 under moderately stringent conditions; and

(c) complements of sequences of (i) or (ii);  
in combination with an immunostimulant.

25. A vaccine according to claim 24, wherein the immunostimulant is an adjuvant.

26. A vaccine according to claim 24, wherein the immunostimulant induces a predominantly Type I response.

27. A vaccine according to claim 24, wherein the antigen-presenting cell is a dendritic cell.

28. A method for inhibiting the development of a cancer in a patient, comprising administering to a patient an effective amount of an antigen-presenting cell that expresses a polypeptide comprising at least an immunogenic portion of a breast tumor protein, or a variant thereof, wherein the tumor protein comprises an amino acid sequence that is encoded by a polynucleotide sequence selected from the group consisting of:

(a) sequences recited in SEQ ID NOs: 1-38, 42-204, 205, 207 and 210-290;

(b) sequences that hybridize to a sequence recited in any one of SEQ ID NOs: 1-38, 42-204, 205, 207 and 210-290 under moderately stringent conditions; and

(c) complements of sequences of (i) or (ii) encoded by a polynucleotide recited in any one of SEQ ID NOs: 1-38, 42-204, 205, 207 and 210-290; and thereby inhibiting the development of a cancer in the patient.

29. A method according to claim 28, wherein the antigen-presenting cell is a dendritic cell.

30. A method according to any one of claims 20, 21 and 28, wherein the cancer is breast cancer.

31. A method for removing tumor cells from a biological sample, comprising contacting a biological sample with T cells that specifically react with a breast tumor protein, wherein the tumor protein comprises an amino acid sequence that is encoded by a polynucleotide sequence selected from the group consisting of:

(i) polynucleotides recited in any one of SEQ ID NOs: 1-38, 42-204, 205, 207 and 210-290; and

(ii) complements of the foregoing polynucleotides;

wherein the step of contacting is performed under conditions and for a time sufficient to permit the removal of cells expressing the antigen from the sample.

32. A method according to claim 31, wherein the biological sample is blood or a fraction thereof.

33. A method for inhibiting the development of a cancer in a patient, comprising administering to a patient a biological sample treated according to the method of claim 31.

34. A method for stimulating and/or expanding T cells specific for a breast tumor protein, comprising contacting T cells with at least one component selected from the group consisting of:

(a) polypeptides comprising at least an immunogenic portion of a breast tumor protein, or a variant thereof, wherein the tumor protein comprises an amino acid sequence that is encoded by a polynucleotide sequence selected from the group consisting of:

- (i) sequences recited in SEQ ID NOs: 1-38, 42-204, 205, 207 and 210-290;
  - (ii) sequences that hybridize to a sequence recited in any one of SEQ ID NOs: 1-38, 42-204, 205, 207 and 210-290 under moderately stringent conditions; and
  - (iii) complements of sequences of (i) or (ii);
- (b) polynucleotides encoding a polypeptide of (a); and
- (c) antigen presenting cells that express a polypeptide of (a);
- under conditions and for a time sufficient to permit the stimulation and/or expansion of T cells.

35. An isolated T cell population, comprising T cells prepared according to the method of claim 34.

36. A method for inhibiting the development of a cancer in a patient, comprising administering to a patient an effective amount of a T cell population according to claim 35.

37. A method for inhibiting the development of a cancer in a patient, comprising the steps of:

(a) incubating CD4<sup>+</sup> and/or CD8<sup>+</sup> T cells isolated from a patient with at least one component selected from the group consisting of:

(i) polypeptides comprising at least an immunogenic portion of a breast tumor protein, or a variant thereof, wherein the tumor protein comprises an amino acid sequence that is encoded by a polynucleotide sequence selected from the group consisting of:

(1) sequences recited in SEQ ID NOs: 1-38, 42-204, 205, 207 and 210-290;

(2) sequences that hybridize to a sequence recited in any one of SEQ ID NOs: 1-38, 42-204, 205, 207 and 210-290 under moderately stringent conditions; and

- (3) complements of sequences of (1) or (2);
- (ii) polynucleotides encoding a polypeptide of (i); and
- (iii) antigen presenting cells that expresses a polypeptide of (i);

such that T cells proliferate; and

- (b) administering to the patient an effective amount of the proliferated T cells, and thereby inhibiting the development of a cancer in the patient.

38. A method for inhibiting the development of a cancer in a patient, comprising the steps of:

- (a) incubating CD4<sup>+</sup> and/or CD8<sup>+</sup> T cells isolated from a patient with at least one component selected from the group consisting of:

- (i) polypeptides comprising at least an immunogenic portion of a breast tumor protein, or a variant thereof, wherein the tumor protein comprises an amino acid sequence that is encoded by a polynucleotide sequence selected from the group consisting of:

- (1) sequences recited in SEQ ID NOs: 1-38, 42-204, 205, 207 and 210-290;

- (2) sequences that hybridize to a sequence recited in any one of SEQ ID NOs: 1-38, 42-204, 205, 207 and 210-290 under moderately stringent conditions; and

- (3) complements of sequences of (1) or (2);

- (ii) polynucleotides encoding a polypeptide of (i); and

- (iii) antigen presenting cells that express a polypeptide of (i);

such that T cells proliferate;

- (b) cloning at least one proliferated cell to provide cloned T cells;

and

- (c) administering to the patient an effective amount of the cloned T cells, and thereby inhibiting the development of a cancer in the patient.



39. A method for determining the presence or absence of a cancer in a patient, comprising the steps of:

(a) contacting a biological sample obtained from a patient with a binding agent that binds to a breast tumor protein, wherein the tumor protein comprises an amino acid sequence that is encoded by a polynucleotide sequence recited in any one of SEQ ID NOs: 1-38, 42-204, 205, 207 and 210-290 or a complement of any of the foregoing polynucleotide sequences;

(b) detecting in the sample an amount of polypeptide that binds to the binding agent; and

(c) comparing the amount of polypeptide to a predetermined cut-off value, and therefrom determining the presence or absence of a cancer in the patient.

40. A method according to claim 39, wherein the binding agent is an antibody.

41. A method according to claim 40, wherein the antibody is a monoclonal antibody.

42. A method according to claim 40, wherein the cancer is breast cancer.

43. A method for monitoring the progression of a cancer in a patient, comprising the steps of:

(a) contacting a biological sample obtained from a patient at a first point in time with a binding agent that binds to a breast tumor protein, wherein the tumor protein comprises an amino acid sequence that is encoded by a polynucleotide sequence recited in any one of SEQ ID NOs: 1-38, 42-204, 205, 207 and 210-290 or a complement of any of the foregoing polynucleotide sequences;

(b) detecting in the sample an amount of polypeptide that binds to the binding agent;

(c) repeating steps (a) and (b) using a biological sample obtained from the patient at a subsequent point in time; and

(d) comparing the amount of polypeptide detected in step (c) to the amount detected in step (b) and therefrom monitoring the progression of the cancer in the patient.

44. A method according to claim 43, wherein the binding agent is an antibody.

45. A method according to claim 44, wherein the antibody is a monoclonal antibody.

46. A method according to claim 43, wherein the cancer is a breast cancer.

47. A method for determining the presence or absence of a cancer in a patient, comprising the steps of:

(a) contacting a biological sample obtained from a patient with an oligonucleotide that hybridizes to a polynucleotide that encodes a breast tumor protein, wherein the tumor protein comprises an amino acid sequence that is encoded by a polynucleotide sequence recited in any one of SEQ ID NO: 1-38, 42-204, 205, 207 and 210-290 or a complement of any of the foregoing polynucleotide sequences;

(b) detecting in the sample an amount of a polynucleotide that hybridizes to the oligonucleotide; and

(c) comparing the amount of polynucleotide that hybridizes to the oligonucleotide to a predetermined cut-off value, and therefrom determining the presence or absence of a cancer in the patient.

48. A method according to claim 47, wherein the amount of polynucleotide that hybridizes to the oligonucleotide is determined using a polymerase chain reaction.

49. A method according to claim 47, wherein the amount of polynucleotide that hybridizes to the oligonucleotide is determined using a hybridization assay.

50. A method for monitoring the progression of a cancer in a patient, comprising the steps of:

(a) contacting a biological sample obtained from a patient with an oligonucleotide that hybridizes to a polynucleotide that encodes a breast tumor protein, wherein the tumor protein comprises an amino acid sequence that is encoded by a polynucleotide sequence recited in any one of SEQ ID NO: 1-38, 42-204, 205, 207 and 210-290 or a complement of any of the foregoing polynucleotide sequences;

(b) detecting in the sample an amount of a polynucleotide that hybridizes to the oligonucleotide;

(c) repeating steps (a) and (b) using a biological sample obtained from the patient at a subsequent point in time; and

(d) comparing the amount of polynucleotide detected in step (c) to the amount detected in step (b) and therefrom monitoring the progression of the cancer in the patient.

51. A method according to claim 50, wherein the amount of polynucleotide that hybridizes to the oligonucleotide is determined using a polymerase chain reaction.

52. A method according to claim 50, wherein the amount of polynucleotide that hybridizes to the oligonucleotide is determined using a hybridization assay.

53. A diagnostic kit, comprising:

(a) one or more antibodies according to claim 10; and

(b) a detection reagent comprising a reporter group.

54. A kit according to claim 53, wherein the antibodies are immobilized on a solid support.

55. A kit according to claim 53, wherein the detection reagent comprises an anti-immunoglobulin, protein G, protein A or lectin.

56. A kit according to claim 53, wherein the reporter group is selected from the group consisting of radioisotopes, fluorescent groups, luminescent groups, enzymes, biotin and dye particles.

57. An oligonucleotide comprising 10 to 40 contiguous nucleotides that hybridize under moderately stringent conditions to a polynucleotide that encodes a breast tumor protein, wherein the tumor protein comprises an amino acid sequence that is encoded by a polynucleotide sequence recited in any one of SEQ ID NOs: 2-5, 7, 10, 13, 15, 16, 21, 22, 25, 28, 30, 33, 35, 63, 72, 73, 81, 103, 104, 107, 109, 118, 128, 129, 134, 136, 159-161, 164, 181, 183-204 and 210-290 or a complement of any of the foregoing polynucleotides.

58. A oligonucleotide according to claim 57, wherein the oligonucleotide comprises 10-40 contiguous nucleotides recited in any one of SEQ ID NOs: 2-5, 7, 10, 13, 15, 16, 21, 22, 25, 28, 30, 33, 35, 63, 72, 73, 81, 103, 104, 107, 109, 118, 128, 129, 134, 136, 159-161, 164, 181, 183-204 and 210-290.

59. A diagnostic kit, comprising:  
(a) an oligonucleotide according to claim 58; and  
(b) a diagnostic reagent for use in a polymerase chain reaction or hybridization assay.

## SEQUENCE LISTING

<110> Corixa Corporation  
 Dillon, Davin C.  
 Day, Craig H.  
 Jiang, Yuqiu  
 Wang, Aijun  
 Houghton, Raymond L.  
 Mitcham, Jennifer L.

<120> COMPOSITIONS AND METHODS FOR THERAPY AND  
 DIAGNOSIS OF BREAST CANCER

<130> 210121.491PC

<140> PCT  
 <141> 2000-11-29

<160> 290

<170> FastSEQ for Windows Version 3.0

<210> 1  
 <211> 298  
 <212> DNA  
 <213> Homo sapien

<400> 1

ctgaacagtg tcagctccgt gctggagaca gtccctgctga tcacctgaat gctgaacatg	60
cttcgtgggg ctatcttttg tttctctgt agtctcttg gtgatctcat ctgcttttct	120
gctcgagtga tgacagcctt gaaccttgtc cttccttgtc tcagagggga aaaaggaatt	180
ggatttcctc agggctctggg gctgggctg tggcttgagg ttccgagact gatgaatcca	240
agcatgcttg agggcctggt ccggggctcat gcgaagagaa ggttcccata ccaaacac	298

<210> 2  
 <211> 276  
 <212> DNA  
 <213> Homo sapien

<400> 2

tggaaggtgt ggtgactaag ggccacgggt attgggtgaa atttgagatt gtaggccaac	60
tgtattttca agctttctgaa cttaggcaaa atattcatcg caaagtctct agcgtcatat	120
ttttctcacc taaattacgt ttccacgaga ttatttatat atagttggtc tatctctgca	180
gtccttgaag gtgaagttgt gtgttactag gctgtgtttt gggatgtcag cagtggcctg	240
aagtgagttg tgcaataaat gttaagttga aacctc	276

<210> 3  
 <211> 405  
 <212> DNA  
 <213> Homo sapien

<220>  
 <221> misc\_feature  
 <222> (1)...(405)  
 <223> n = A,T,C or G

<400> 3

```

tcacatggct atttcattta tttagtagtt ttgaaatggt agcaaataata aggtatttgt      60
aaagcatctt tcattataaa gagatttagta atattcacca atcatgccaa tgagattata      120
cactctgcc aagactacta naaaaaatttg atcattatta aattcaatgt tatttgacag      180
tgtgaactct atgtaacagc acaaaattctg gactttgaat ctggctgctg tcttcacctg      240
aaccattaaa atgaccttgt taacaaggaa ggaatcaatg gggaaatatac acaaccagag      300
attggctgtg tgtccaaggg tgccttgtct tgttgccagg atcagactgt gaaatcacag      360
aggcaagctg atgtcatcag aggtgactct gcccacaaca caatg                      405

```

<210> 4  
 <211> 696  
 <212> DNA  
 <213> Homo sapien

```

<400> 4
cattgtgttg ggcactgtta cagtgaacag gaaacgtgga aaatcacagc caaactgtgc      60
tctgaaagaa cactctatgt ctaatatagc cagcgtcaag agtccttatg aggcggagaa      120
ctcgggggaa gagctggatc agaggtattc caaggccaag ccaatgtgta acacatgtgg      180
gaaagtgttt tcagaagcca gcagtttgag aaggcacatg agaatacata aaggagtcaa      240
accttacgtc tgccacttat gtggaaggc atttacccea tgtaaccagc tgaaaacgca      300
tgtaagaact catacagggt agaagccata caaatgtgaa ttgtgtgata aaggatttgc      360
tcagaaatgt cagctagtct tccatagtct catgcatcat ggtgaagaaa aaccctataa      420
atgtgatgta tgcaacttac agtttgcaac ttctagcaat ctcaagattc atgcaaggaa      480
gcatagtgga gagaagccat atgtctgtga taggtgtgga cagagatttg ctcaaggcag      540
cacactgacc tatcatgtcc gtaggcatac tggagaaaag ccttatgtat gtgatacctg      600
tgggaaggca tttgtctgtc ctagtctctc tatcactcat tctcgaaaac atacaggtaa      660
gtttgacagg gagagactgc ttaaaataaaa gttata                             696

```

<210> 5  
 <211> 580  
 <212> DNA  
 <213> Homo sapien

<220>  
 <221> misc\_feature  
 <222> (1)... (580)  
 <223> n = A,T,C or G

```

<400> 5
acatcaaaaa ggaaatatatt ttgacttgct tttcttctgt aaatcctccc atctcactaa      60
tatttacaac aatccagagt agcgtttatg agacactgaa aaagacaggg aggaaatcct      120
ttttcaagat atgaagtcag aacctgaatg tagacatcgg acagagaagt cctcaaccac      180
aaacctgtcc tccagctcta gagagagtaa ggctgtattt ccaaccttga gatttttcat      240
tacatthttcc cctttttggg tgttaaattc ttccaagaa tgctgtactt gtaaaaatga      300
ttttattcta gctacaaaac atttcattta anaaaaccgc attttatatc cttgtgtgaa      360
atgctcccaa aagccatcaa gatatggaga caacagattt taaaaacata aatctaataca      420
tatgggcttg aaacagtatg aacatttaac agagtgcac gatatcatta ttatatattgt      480
ttgtcatgag atgaaaggcc tggaggcaga tgggtattaa tcataattcc tgagcttcta      540
cagaaatttt aaaaatgaaat tactaactgc ttaaaattat                             580

```

<210> 6  
 <211> 557  
 <212> DNA  
 <213> Homo sapien

```

<400> 6
attacattca agataaaaga tttattcaca ccacaaaaag ataatcacia caaaatatac      60
actaacttaa aaaacaaaag attatagtga cataaaatgt tatattctct ttttaagtgg      120
gtaaaagtat tttgtttgct tctacataaa tttctattca tgagagaata acaaataatta      180

```

aaatacagtg	atagtttgca	tttcttctat	agaatgaaca	tagacataac	cctgaagctt	240
ttagtttaca	gggagtttcc	atgaagccac	aaactaaact	aattatcaaa	cacattagtt	300
atttccagac	tcaaatagat	acacattcaa	ccaataaact	gagaaagaag	catttcatgt	360
tctctttcat	tttctataaa	agcatttttt	cttttgacta	aatgcaaagt	gagaaattgt	420
attttttctc	cttttaattg	acctcagaag	atgcactatc	taattcatga	gaaatacgaa	480
atttcagggtg	tttatcttct	tccttacttt	tgggggtctac	aaccagcata	tcttcatggc	540
tgtgaaattc	atggctg					557

&lt;210&gt; 7

&lt;211&gt; 653

&lt;212&gt; DNA

&lt;213&gt; Homo sapien

&lt;400&gt; 7

cattgtgttg	ggggaagtag	ggaatattat	tgaggcaggg	taagaaatgg	tttacaattc	60
tgaaaggatg	atcaaagaaa	aactcattgt	tgagaaagta	atatgagtag	agacctgaaa	120
taagttaggg	agtgacgggt	tatgtccagg	gcaataatgt	ttctgacaga	ggggagagtc	180
atttcagaag	cctagaggca	tgtgtaaagc	tgttagaatg	ccagacagtc	accaggccaa	240
gatgtgcaga	tatccataag	tgaaggggaa	agaaatacaa	aatgaaggca	gagaaatcac	300
aaaattggat	aagtgggtgc	ttgtaggcca	tgatgatttt	agttcatact	aaaattgagt	360
taggctgcc	ttgtagggtt	tgtgagctca	gggataacat	ggtctgaatt	ttattttctaa	420
aaggatcact	ccaagtgtta	cattgcaaag	aataacgtaa	ggtggctggt	gtagtagact	480
aaagtggaat	atagtaacag	tgaaatacat	tttgtggtta	agcttggtag	atttgaccac	540
acaaaattgt	gaaattacct	gtggcacaaa	aaatatcaaa	ggtacatata	gacagaagaa	600
ccttgcgatt	gtttattaat	gtccttaatt	tataatgtta	ataccagtag	aag	653

&lt;210&gt; 8

&lt;211&gt; 456

&lt;212&gt; DNA

&lt;213&gt; Homo sapien

&lt;400&gt; 8

cattgtgttg	ggctaactct	tgtctctat	ccacctgcc	tagcaattta	tctcaaagct	60
tcaagttcct	gccatctaca	tgtgcccagg	tcaaccaatc	aatggctcag	acagataagc	120
caacatgcat	cccgccggag	ctgccgaaaa	tgtgtaagga	gtttgccaaa	gccgccattc	180
gggcgcagcc	gcaggacctc	atccagtggg	gggcccatta	ttttgaggcc	ctgtcccgtg	240
gagagacgcc	tccggtgaga	gagcgggtctg	agcaggtcgc	tttgtgtaac	tgggcagagc	300
taacacctga	gctgtttaaag	atcctgcatt	ctcaggttgc	tggcagactg	atcatccgtg	360
cagaggagct	ggcccagatg	tggaaagtgg	tgaatctccc	aacagatctg	tttaatagtg	420
tgatgaatgt	gggtcgcttc	acggaggaga	tcgagt			456

&lt;210&gt; 9

&lt;211&gt; 512

&lt;212&gt; DNA

&lt;213&gt; Homo sapien

&lt;400&gt; 9

gtttttgatt	cattttat	taacaatgtt	taacaatgta	agtccacata	taagataccc	60
aagctttaaa	tatctatata	tataaactga	tttcaacatc	tttggettca	aaacagtaaa	120
attgtttttc	caatatcaaa	caagtcaa	ttggaaaagg	cataaatctg	tatgaacatc	180
ctgtatccat	ggagatgtca	tgactaaatt	cagaaatagc	ctcatctctc	tttgtttttg	240
ctttcttatg	tctgagttct	gcattccaatt	ctgtttatta	catagttttc	tataagattg	300
tacctctttt	aaacagtgtc	tattgatata	tattctaggt	gtctggaagt	ctttttctat	360
agtcggctct	tggttgtctc	tgggaatatg	aatggaagga	gcagagtga	aataaatctg	420
agggcaatat	tcataaataa	tccaagagct	acactgtagt	caactctccc	cagagcctga	480
ccacagtgtt	tcctctctc	ctcctcccaa	cc			512

&lt;210&gt; 10

<211> 308  
 <212> DNA  
 <213> Homo sapien

<220>  
 <221> misc\_feature  
 <222> (1)...(308)  
 <223> n = A,T,C or G

<400> 10  
 atgtttatga agacctttaa atatttatat agaaacaaaa tgtcattgca acctaacatc 60  
 atccattaaa aataaaagga aaggaaaacg gcagggaata gtgcagtaat aacaaatggg 120  
 gacatgcttg gtcttaagca tcatagcaaa ctcatatttt ccaatgaaac aaggattttt 180  
 agacccatct ttggaaatga ttcccaaatt aganaaccat cagggtctcaa aaaaggaagg 240  
 gtcacaaag tccatccagc ccagccaccc tgaggngcct gtatctcttc aacaagccca 300  
 acacaatg 308

<210> 11  
 <211> 510  
 <212> DNA  
 <213> Homo sapien

<220>  
 <221> misc\_feature  
 <222> (1)...(510)  
 <223> n = A,T,C or G

<400> 11  
 attatatgaa tattttaatg caaaatgctt aacacttaaa attagcaaag cgtcatttaa 60  
 attaaaattc catttaacta aagatgggta accccaanaa attgtacagt agttgatttc 120  
 tgctatataa tgccagtcct atgccatata ataagaactg caacattagc tgcacttcc 180  
 tccattgctc ttctggaccc taagggatga gggaggggac tcagacacaa aacacaaccc 240  
 aaataaaactg tgcagtgtatt cctaatagtt ataaacccaa tctaagttgt ccaaacagct 300  
 gaagaataac tgcaggtatt gttccanagc tgatacgagg ttttgctttt acagcctggg 360  
 aaaagtctcg cactaggtga gaagtcacag tttaaggatg catgttctgt aaatagttac 420  
 tacatatata catttactgt ctgtaaacac tagaaatata cattagacag agtaccctca 480  
 caagttgggt acagtttaaa aaagaagatg 510

<210> 12  
 <211> 611  
 <212> DNA  
 <213> Homo sapien

<220>  
 <221> misc\_feature  
 <222> (1)...(611)  
 <223> n = A,T,C or G

<400> 12  
 agttttataa aatattttat ttacagtaga gctttacaaa aatagtctta aattaatata 60  
 aatccctttt gcaatataac ttatatgact atcttctcaa aaacgtgaca ttcgattata 120  
 acacataaac tacatttata gttgttaagt caccttgtag tataaatatg ttttcatctt 180  
 ttttttgtaa taaggnacat accaataaca atgaacaatg gacaacaaat cttattttgt 240  
 tattcttcca atgtaaaatt catctctggc caaaacaaaa ttaaccaaag aaaagtaaaa 300  
 caattgtccc tctgttcaac aatacagtc tttttaatta ttgagagtt tatctgacag 360  
 agacacagca ttaactgaa agcaccatgg cataaagtct agtaacatta tcctcaaaag 420  
 ctttttccaa tgtctttcct tcaactgttt attcagtatt tggccagtag aaataaagat 480  
 tggctcaca tctctcttcc attagttctca agtgttccta ttatgcactg agttttcaga 540



ccttcccaac tggcatgtgt ttttaagtgtg agtttctttc tttggcttca agtggagttt 600  
cacaacattt a 611

<210> 13  
<211> 394  
<212> DNA  
<213> Homo sapien  
  
<220>  
<221> misc\_feature  
<222> (1)...(394)  
<223> n = A,T,C or G

<400> 13  
caatgttttag attcatttta ttagtggcat atacaaagca ccatataata tatgaaacgt 60  
anaacaatca tgactatgta attaaactgta naaataaactg ctaanaaaat atagcaatat 120  
ttaacacagg atttctaaaa ccattatatt ttcattactt tccccaaagc taatgtccca 180  
tgttttatatt tatanacttt gtttatcaag atttatatgc atttggcacc tttttgggct 240  
gaaaatagtt gatgtactct gtacagtaat gttacagttt tatacaaaat tcanaaatat 300  
tgcatttgga atagtcttta tggctcctct ccaagtattc agtttcacac aacagcaaac 360  
actctgaatg cctttcctcc tgcccaaacac aatg 394

<210> 14  
<211> 361  
<212> DNA  
<213> Homo sapien  
  
<220>  
<221> misc\_feature  
<222> (1)...(361)  
<223> n = A,T,C or G

<400> 14  
agcaggnaact ataattttat aattaatttt acaattcatg tagcaaatgg aaaatcatat 60  
agagaggcca atgtatataa ataagagttt atacagaaac tgccaattca caaaacagca 120  
ctgcatgggt tctatattgc aagcacaaga catggtcaca tggttccact gtacaggtag 180  
aaacaagccc acagacaata catagagtac cacctgaaac gaggcccttg gagctgctca 240  
gcttcttana aaataganaa ctttcaatgg tcataataca ttttgattca aaatgtcttc 300  
taaaatgttt tcattgtggg agaaaattaa gaaggggcaa aaatccatct atggaaacttc 360  
t 361

<210> 15  
<211> 537  
<212> DNA  
<213> Homo sapien  
  
<220>  
<221> misc\_feature  
<222> (1)...(537)  
<223> n = A,T,C or G

<400> 15  
acttacaaaa ttaattttat tttgcaaaac tcaacaaata cacgttcaga tctggtttct 60  
cttcaaaaaca tgtgtttgtt tttttaacaa acatgcaagt taatttggca tgccaaacat 120  
ctttctctct agctcgctt ggaaaaattt ttttcataac acaaaacaagg gtgcaaatat 180  
tgtccaaacc tatttacatt tttacctctt agaattacat acattaatat ttattgggag 240  
gaaagcaaaa ctgcaaaaca tagtcttttg cattcacatt tgcttcagca gtataattaa 300  
aaccttatat ttgttttaaa gataaacagt ttgaaggaaa ttttaataat cttgttttgg 360

ctctgcaaag gagccactat atcaaagcat ttaactggag ctgttgagtt cctgctggta	420
gaatattact tccagcctat ttattagctt gtcttcgggn ggccaatac atgctttttt	480
ccctctacac tgaatgaaag tacaaaaaga aaaccatttc ttttcccaa cacaatg	537

&lt;210&gt; 16

&lt;211&gt; 547

&lt;212&gt; DNA

&lt;213&gt; Homo sapien

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (1)...(547)

&lt;223&gt; n = A,T,C or G

&lt;400&gt; 16

gggtgtggng atgtatttat tcataatata ttttcagaac acattaataa tggagaataa	60
cacttattca tatactgaat ataacttttc ctggagcact cttagagcttg tttggagttg	120
gagaatactg ccaggctttt cctaattctt ttggtctttg gaagtgggca gggtttctca	180
aaccaagtgt ctccatggg ccattggcaa aggtctccct tcatcagctt ggaggggcag	240
aaagaccatg gcttcagcac ttccattttg gaaagaagta acaaaaaagt gaattaatga	300
gcaatcgga agactcaaag cattttgtac tccacagttc atttcttcac acaaacgtcc	360
attactgcag cgggcatgaa aaccggcagg gtgttaggct catggcctga agagaagtca	420
catcaccagc cgatgttttc atgcaaaaagg caatcgtgat gattcanaac ctgggtctga	480
atttctccag gtgtgctcgt gagctgaagg tcatgcccac tctgtgcac ctgtgcccaa	540
cacaatg	547

&lt;210&gt; 17

&lt;211&gt; 342

&lt;212&gt; DNA

&lt;213&gt; Homo sapien

&lt;400&gt; 17

acattaagaa gctcctcttc tagcatgtcc ttaagaagcc tgtcttgag cactttcata	60
tcttctttca tcaaacacat ctggatgta aaaacagttt cttcactatc agtattacag	120
aagacacttt tagccaatga agttttcaaa agaagaaagc ctctgttgtt cgcttttttg	180
atatgcactg aacttctgaa atatcttttc ccaaaagtcc acaaattcct tttccaaatc	240
ttttaagac tgtgaatctt tttcaaaatt ctccagctcc tctatgataa tgaattggaa	300
tttatcaagt tttttaatcc tagagtctgt actttggatg at	342

&lt;210&gt; 18

&lt;211&gt; 279

&lt;212&gt; DNA

&lt;213&gt; Homo sapien

&lt;400&gt; 18

catcataagg ttttattcat atatatacag ggtattaaga attaagagga tgctgggctc	60
tggtcttggc ttggaagatt ctatttaatt gaaactctct gtccagaaag caataacttt	120
gtctcggtcc tggtgggctg aaccctaagg tgagtgtgca gtacagtgtg tgggggtgaa	180
atggagattt ggaattgaac tctctgctg taaatgttcc ccaataaatt gttgtgtgta	240
tgatacgtgt ataataaaag tattcttggt agaactctga	279

&lt;210&gt; 19

&lt;211&gt; 239

&lt;212&gt; DNA

&lt;213&gt; Homo sapien

&lt;400&gt; 19

ctgccagcgt ttttgtgtgg ctgcagtgtg cctggggcca gctcacgggc agtgggtgga	60
---	----

```

cctaactgcc caggcaggcg agagctactt ccagagcctt ccagtgcatt ggagggcagg      120
gctaggtgta gcggtgtctc ctctttgaaa ttaagaacta tctttcttgt agcaaagctg      180
cacctgatga tgtgcctctt cctctctgtg ttgtctgggc ccttgtttac aagcacgcg      239

```

```

<210> 20
<211> 527
<212> DNA
<213> Homo sapien

```

```

<400> 20
ctgaaccatt atgggataaa ctgggtgcaa ttctttgcct tctctacttc tctactgattg      60
aacataagct tccagggctc cctgatgag gaggagcctg tccttttcag atggatggtc      120
atccagccac tgagagaagc gtgtgtggga ccactctgcc ctctggaaag gagatttcag      180
ttcagcgggt gctctcgtga acaaaaactg aataatgatg ctgaacggaa tcacatcccc      240
caatgcagga ctactggcta catgttcact tgcctggaag agcagaggct tgaatgatct      300
cagcatccga taggactttc ctaaatcaga tactcgtcta cagaatgaac ccacagccaa      360
ctccatctgt gcaaaatcag cagcaagtcg cattttccca ccttcaccaa gaggtcttat      420
gagactggca tggcggataa aaagttcaac agctctttgg gcaataacct cagtgttgct      480
aaagacaaaa tccaagcatt caaagtgttt aaaatagtc ctcataa      527

```

```

<210> 21
<211> 399
<212> DNA
<213> Homo sapien

```

```

<400> 21
ctgcaatggg tgcaagtgtt atttcacact agctctgact ctccacttct aaccagacaa      60
acagccaacc aaccaatcaa catgtattta ataaccacct atgggggtgca aagcacaaaa      120
gggcactcat cttgaaaagg aaagaccaag aatgtgctag agtaaagaga cagagaccag      180
accctactct caagatcaag agacttcagt ctgggagaca tctgccattt ctctcttctt      240
aataaacctc atttgccctt aaaaatacat ttgctttggg ggcccagaat caagaaagga      300
aactttacaa agtaaacaga agttactccc cacagggagg cagaagcaga ttaaccccaa      360
cagcagacat ctgcccggaa gagcaaacct cacatctgg      399

```

```

<210> 22
<211> 532
<212> DNA
<213> Homo sapien

```

```

<400> 22
ccagaagggt aagaaaagtt atctgataat gctcaaagtg cagtagaaat acttttaacc      60
attgatgata caaagagagc tggaatgaaa gagctaaaac gtcacccctt cttcagtgat      120
gtggactggg aaaatctgca gcatcagact atgcctttca tccccagcc agatgatgaa      180
acagatacct cctattttga agccaggaat actgctcagc acctgaccgt atctggattt      240
agtctgtagc acaaaaattt tccttttagt ctagcctcgt gttatagaat gaacttgcat      300
aattatatac tccttaatac tagattgatc taagggggaa agatcattat ttaacctagt      360
tcaatgtgct tttaatgtac gttacagctt tcacagagtt aaaaggtgga aaggaatata      420
gtcagtaatt tatcttaacc tcaaaactgt atataaatct tcaaaagctt tttcatctat      480
ttattttggt tattgcactt tatgaaaact gaagcatcaa taaaattaga gg      532

```

```

<210> 23
<211> 215
<212> DNA
<213> Homo sapien

```

```

<400> 23
tgcaataaag ggctgctgtt tcgacgacac cgttcgtggg gtccctgggt gcttctatcc      60
taataaccac gacgtccctc cagaagagga gtgtgaattt tagacacttc tgcagggatc      120

```

```

tgctgcctc ctgacacggg gccgtcccca gcacgggtgat tagtcccaga gctcggctgc 180
cacctccacc ggacacctca gacacgcttc tgcag 215

```

```

<210> 24
<211> 215
<212> DNA
<213> Homo sapien

```

```

<400> 24
cctgaggctc caggctaaga agtagccaag ttccacctgg agagaagagt agagggactt 60
cccaaatctt ttctctgaact cagctctgat actcagaagg tcagtctcac atcgagagat 120
aaggatgcga atcaggactt ggttaattggg ctccagtttcc tagtagggga agaaagagat 180
ggggggtagt tagtgagagt ctccactgaga gtagg 215

```

```

<210> 25
<211> 530
<212> DNA
<213> Homo sapien

```

```

<400> 25
ttttttttct agtaagacta gatttattca ataccctagt aaaagttttg attataagta 60
tccaacagta taaaaagtac aaaacagatc tgtagatttc taatatatta atacaaagtg 120
catgactaca tacagtacat cctacaggca aagagagggt gaaggggaaa aagaagactg 180
tggttgagggt ctagtaataa ataaataaat acagaagtag agatgatcca tattatagta 240
tattctacca ccaatactgc agccaaaatg tacaaaaaaa atcatttcaa ataactcagg 300
aggatgataa tggtctggact tttgtaattc acctcaaaga ctgtgggaga gccactcaa 360
ctcactgtat agtctgtgca tatggtgggt ttagcatgt aggttttttc caaaagaagg 420
aaatataaaa tgttttagatt aagaactata aaactacagg gtgcctataa aaggtggctt 480
actccttatt gttattatac tatccaattt ttaaaatgca gtttaaaaaa 530

```

```

<210> 26
<211> 366
<212> DNA
<213> Homo sapien

```

```

<400> 26
ccagcagttc tcggacctcc tctgggggca gggagaggcc attgggtcag gggctggacc 60
caggaggagt tggaatgggt gaaagatggg gagcaagttt ttaggggtaca gggtaggcct 120
aagatgggtc agtagacaga tgggagcaca gaggagggca gggggtgagg tcaagtgagg 180
gccacaggat gtgctgaggg ctcccaggga gccctaccca ggctcacgtc ctccgtgtca 240
ccacctgtac tgtctggggt ccacaggggt tgggcgttgc caggagcac tgggagggcc 300
tcggtagggt ccacctgtag ggagaggatg tcaggaccac tagcctctgg gcaagggcag 360
aggagg 366

```

```

<210> 27
<211> 331
<212> DNA
<213> Homo sapien

```

```

<220>
<221> misc_feature
<222> (1)...(331)
<223> n = A,T,C or G

```

```

<400> 27
ccaaactcag agatgggtacc agccaggggc aagcatgacc agagccaggg accctgtggc 60
tctgatcccc catttatcca ccccatgtgc ctccaggacta gaggtagcaa tcatacctta 120
taaagtactt ttgtgccttt ctgctccagt ctcaaaattt cctacacctg ccagttcttt 180

```

```

acatttttcc aaggaaagga aaacggaagc aggggttcttg cctggtagct ccaggaccca      240
nctctgcagg caccacaaga ccctctgtgt ccagcctctt ccttgagttc tcggaacctc      300
ctccctaatt ctcccttctt tccccacaag g                                331

```

```

<210> 28
<211> 530
<212> DNA
<213> Homo sapien

```

```

<400> 28
ccatgaatgc ccaacaagat aatattctat accagactgt tacaggattg aagaaagatt      60
tgtcaggagt tcagaaggtc cctgcactcc tagaaaatca agtggaggaa aggacttggt      120
ctgattcaga agatattgga agctctgagt gctctgacac agattctgaa gagcagggag      180
accatgcccc cccaagaaa cacaccacgg accctgacat tgataaaaaa gaaagaaaaa      240
agatgggtcaa ggaagccag agagagaaaa gaaaaaacia aattcctaaa catgtgaaaa      300
aaagaaagga gaagacagcc aagacgaaaa aaggcaaata gaatgagaac catattatgt      360
acagtcattt tcctcagttc cttttctcgc ctgaactctt aagctgcac tcggaagatgg      420
cttattgggt ttaaccagat tgtcatcgtg gactgtctg tgaagacgga ttcaaagtgt      480
ttcatgtaac tatgtaaaaa gctctaagct ctagagtcta gatccagtca                    530

```

```

<210> 29
<211> 571
<212> DNA
<213> Homo sapien

```

```

<220>
<221> misc_feature
<222> (1) ... (571)
<223> n = A,T,C or G

```

```

<400> 29
ccataatatt ctgatgatca aggagcacac atatacaaaa gttattggat tactgcaatt      60
ctcagaggca caaaacctga catggtgtga tatagtatat aatcagtcac gggggggaaa      120
agaacattaa gtcttttaaa aggcttagga agacataaac agtaaattct tgtttttcta      180
ccttcctttg gacagtgtta tatttcactt tcttctttgc aaaatgtttc caaattcatt      240
tgctcaggat ttattttaaga taataactta aaacaactaa cagtgtgtta tgctatatgc      300
atatcatgca tgttctactg gttcaaggac aaaattaaaa caagatcttc tctgtaaagc      360
aaatatattt attatgcact ttcatatata cagggttttt ttgagtacca angggataaa      420
ataaaacttt tacaatgtga aattcaatgt acatttttgg ctattttacat acctcaaac      480
aagggaaaaa taaaaagaaa gcatttgttt gcaactacat ttgctgagaa gtgtaaattg      540
aggacattaa gcaaaacaaa ttttgcata g                                571

```

```

<210> 30
<211> 917
<212> DNA
<213> Homo sapien

```

```

<400> 30
actgccagag agtatgattt gaaggagatg ggagcagatg taattcttgg ctggaatctc      60
tcatttcaaa atcacttcac ataatggtgt catcatttaa acacttaaca gtcagtgcaa      120
ctgccactgt aacatctagt tggacaaaac cacaaggagg gggaggagaa aatgccatca      180
ctattatggt aacaaacatt taatttaaat gggttgctgca ctagttaaatt tctgcagaaa      240
acagttttac ccgccccctt tcacagttcc aaattaatca aggatgcttt tctataatct      300
gatgcttagc aaattagctc atgattcaaa ttttgccttc ttgaagcaca tatacctttt      360
attttaaaag tccattatag agaatttgga atatataagg tatttgaatt gcagaacacc      420
cctctaattc tgtaaatata gcaaagacaa aacagtatca tatacatcaa gatcatactt      480
ttaagtaag tttaaaggtc tcaattgccc agatattaaa tttatatttt ccttctatta      540
aaaaatatta catttcaatt ttgtaatttt gtaacatatt ttaagatgac cagcaagacc      600

```

```

tagtcaattt gaaaataccc ttgcattcca tacacaagct ataccataag taataaccca 660
agtatatgat gtgtaaaagt tgggtgaagg cataatactg aatttttttg caaatgtaaa 720
ctgctttcca agtaatcagc accatttttt actagactac attttaatca ctcccttagc 780
tgcttacaac ctctacttag gcataaataa aagaatctga aattggtata tttccccttc 840
ctgctgtgtt aaccaaaaat actatttgac ttaaagatca aagagtcttt ttcctgaagg 900
tttttgtttt taaatgt

```

```

<210> 31
<211> 367
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(367)
<223> n = A,T,C or G

```

```

<400> 31
tcttttcttt ctgtatttcc caaattacag ggagctatgc ccttgggtatt gcacacagta 60
cactgcaaaa gattcacaag gttagttgaa agtcattttt gccctgggtga ttcaaagctc 120
aaanaatttt ctagcataaa gtcttattaa aaattttaat caaaaatatta tttgagttta 180
agttaataaa aacaatacca ctatatatac tctcaacaac ttcattatat aatcagtcct 240
atgagggtgt acttgctttt catatcacac tgattaagga caaaaataat tttgatgtac 300
atgtaccata cactgatatg caatctacac actgatgcat ttacatacat acaaccccaa 360
cacaatg

```

```

<210> 32
<211> 847
<212> DNA
<213> Homo sapien

```

```

<400> 32
cattgtgttg ggctggcagg atagaagcag cggetcactt ggactttttc accagggaaa 60
tcagagacaa tgatggggct ctccccaga actacagggg ctctggccat ctccgtggta 120
agtccctggat tttcctaata atcacaaact tcctgtcttc ctccctgtgt aaagaatatt 180
atatttgatt gcacaatctt tattataaat tctaaaagga gtgcagtgga aatcaacact 240
ttgaaatgaa atcgtgaaga ttaccaatct ccttcttttg ttgtttttta tgttgatttt 300
tacaatagaaa aataaaccag aaagaaatga gttttaaaaa ccatttagaa ttttttttag 360
ttaatgaatt aagtaatctt aatcacagg tttattttcc acaacatttt cactttcttt 420
aaagtattgc ttttactagt ttttctaacc cacaacaag aacacaggag ccacttctat 480
tttccaagat tacatgtctc ttagcatata gctaagaact ctacacgcct gggcttgata 540
ctgacacgc tttttaaagt aaaaaatcgc agaattaaaa tcaaagcagt gtttgactct 600
agagaagttg ggaggattat taagtaagta tttatgttta gctattatgt gccaaaagaa 660
aatgtcagcc tttggggatg gggggaaaga catacaacat tttaaagcca tttttttcag 720
aaaagtaata cttctgttga ttgagaaagt cgtacatagt attatctaaa agagaaacgg 780
aatgttacag actgtttaaa acctggatgt tacagactaa cttactcctt aactgtgttc 840
ttatagc

```

```

<210> 33
<211> 863
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(863)
<223> n = A,T,C or G

```

&lt;400&gt; 33

cattgtgttg	ggcttttatt	tgagtttatg	aacagaaata	gaaagtatgg	tgcttggggt	60
ttgccctttc	ttactcctga	aagttaaatc	agaagacact	gatttcattt	tgtgaaattt	120
agctcagaga	ctattgatct	tttgtttcat	taatatgaac	aactattagt	aaaaaatagc	180
tttaacagca	tttctgctga	tatctagtaa	tctattcttt	taatgtgaaa	ataagataaa	240
atgtcctgga	gctaattcta	gcttaaattt	gccagtattt	ctgtatgtca	ttaaagtttt	300
ttcctctaag	ggttgtaata	naattttgtt	aatctttgca	tacctgatgg	catctatgtc	360
aatgctgatt	gggtaattat	aaattctgtg	ctaattttaa	acttaatttg	cctcttaagg	420
tgattgtcct	ctgagtaatg	attgtagtta	aatgaagtat	agcttgcaac	tatactatca	480
catgggtcgt	taagtaaaaa	taaataaacc	aaatttgtct	gagacaggct	aagatcaatc	540
ttctcatcaa	accaattttt	ctntaagagc	aatttcactt	tcagttttag	ggtggacatt	600
nttgaatgcc	tcaaattaaa	cgttatctat	ttaatcttcc	tggaatagtc	tgtgaccaa	660
aaggagggtg	tgatatattt	agggtgaaat	atatcacata	tatgggtgta	tatatgtggg	720
atttatatat	tcagctcatt	ctctgtgaag	aagtcttcct	gactaaaatt	ggtttcaaga	780
taaactaatt	tctgttagta	tttctactct	gcctaccatg	tatgcctttt	tgttagaaac	840
taataaatgt	atcagtcnct	agc				863

&lt;210&gt; 34

&lt;211&gt; 432

&lt;212&gt; DNA

&lt;213&gt; Homo sapien

&lt;400&gt; 34

agtgcatctt	ctcttgattt	gtctgggtta	aaaccattcc	ttttgtatga	aatgttttga	60
cttaggaatc	attttatgta	cttggttctac	ctggattgtc	aacaactgaa	agtacatatt	120
tcattccaaat	caagtaaaaa	tgtattttaag	ttgattctga	gagtacagggt	cagtaagcct	180
cattatlttg	aatttgagag	aagggtatagg	tgatcggatc	tgtttcattt	ataaaaaggtc	240
cagtttttag	gactagtaca	ttcctgttat	tttctggggt	ttatcatttt	gcctaaaata	300
ggatataaaa	gggacaaaaa	ataagtagac	tgtttttatg	tgtgaattat	atttctacta	360
aatgtttttg	tatgactgtg	ttatacttga	taatatatat	atatatatat	atatatatca	420
acttggttaa	tt					432

&lt;210&gt; 35

&lt;211&gt; 350

&lt;212&gt; DNA

&lt;213&gt; Homo sapien

&lt;400&gt; 35

ccagaggggt	gtttatctta	gggttggaat	gtttctgatt	atgctgacaa	tagccattag	60
gctgatgttt	tggggctgga	tttaggcagt	ttttaataaa	aagagaactt	aaaatgggtg	120
tgtttgcca	agatgggtgat	gttcctgctg	tcaattagca	taaacaaaag	agaattctga	180
taccctgttg	gaatgtcctc	attcctctga	gcttctccac	tcacaggata	aatgcaggag	240
tggttccccc	tcattggacac	ctgcaaattgc	agagtgtggg	ggctctcctg	gccctgcac	300
actagcaaga	gcaaaaagctg	ctccgagtct	tgtttttaga	acctgggtcga		350

&lt;210&gt; 36

&lt;211&gt; 1082

&lt;212&gt; DNA

&lt;213&gt; Homo sapien

&lt;400&gt; 36

atgaactaca	gcctccactt	ggccttcgtg	tgtctgagtc	tcttctactga	gaggatgtgc	60
atccagggga	gtcagttcaa	cgctcagggtc	ggcagaagtg	acaagctttc	cctgcctggc	120
tttgagaacc	tcacagcagg	atataacaaa	tttctcaggc	ccaatttttg	tggagaaccc	180
gtacagatag	cgctgactct	ggacattgca	agtatctcta	gcatttcaga	gagtaacatg	240
gactacacag	ccaccatata	cctccgacag	cgctggatgg	accagcggct	ggtgtttgaa	300
ggcaacaaga	gcttctactct	ggatgcccgc	ctcgtggagt	tcctctgggt	gccagatact	360
tacattgtgg	agtccaagaa	gtccttctctc	catgaagtca	ctgtgggaaa	caggctcatc	420

cgccctcttct	ccaatggcac	ggtcctgtat	gccctcagaa	tcacgacaac	tggtgcatgt	480
aacatggatc	tgtctaaata	ccccatggac	acacagacat	gcaagttgca	gctggaaagc	540
tggggctatg	atggaaatga	tgtggagttc	acctggctga	gagggaaacga	ctctgtgcgt	600
ggactggaac	acctgcggct	tgctcagtac	accatagagc	ggatatttcac	cttagtcacc	660
agatcgagc	aggagacagg	aaattacact	agattgggtct	tacagtttga	gcttcggagg	720
aatgttctgt	atttcatttt	ggatctctct	cgattcagtc	cctgcaagaa	cctgcattgg	780
ggacaacaaa	ggaagtagaa	gaagtcagta	ttactaatat	catcaacagc	tccatctcca	840
gctttaaacg	gaagatcagc	tttgccagca	ttgaaatttc	cagcgacaac	gttgactaca	900
gtgacttgac	aatgaaaacc	agcgacaagt	taaagtttgt	cttccgagaa	aagatgggca	960
ggattgttga	ttatttcaca	attcaaaacc	ccagtaatgt	tgatcactat	tccaaactac	1020
tgtttccctt	gatttttatg	ctagccaatg	tattttactg	ggcatactac	atgtattttt	1080
ga						1082

&lt;210&gt; 37

&lt;211&gt; 1135

&lt;212&gt; DNA

&lt;213&gt; Homo sapien

&lt;400&gt; 37

atgaactaca	gcttccactt	ggccttcgtg	tgtctgagtc	tcttactga	gaggatgtgc	60
atccagggga	gtcagttcaa	cgctcaggtc	ggcagaagtg	acaagcttcc	cctgcctggc	120
tttgagaacc	tcacagcagg	atataacaaa	tttctcaggc	ccaattttgg	tggagaaccc	180
gtacagatag	cgctgactct	ggacattgca	agtatctcta	gcatttcaga	gagtaacatg	240
gactacacag	ccaccatata	cctccgacag	cgctggatgg	accagcggct	ggtgtttgaa	300
ggcaacaaga	gcttactctt	ggatgcccgc	ctcgtggagt	tcctctgggt	gccagatact	360
tacattgtgg	agtccaagaa	gtccttcctc	catgaagtca	ctgtgggaaa	caggctcatc	420
cgccctcttct	ccaatggcac	ggtcctgtat	gccctcagaa	tcacgacaac	tggtgcatgt	480
aacatggatc	tgtctaaata	ccccatggac	acacagacat	gcaagttgca	gctggaaagc	540
tggggctatg	atggaaatga	tgtggagttc	acctggctga	gagggaaacga	ctctgtgcgt	600
ggactggaac	acctgcggct	tgctcagtac	accatagagc	ggatatttcac	cttagtcacc	660
agatcgagc	aggagacagg	aaattacact	agattgggtct	tacagtttga	gcttcggagg	720
aatgttctgt	atttcatttt	ggaaacctac	gttccctcca	ctttcctggg	ggtgttgtcc	780
tgggtttcat	tttgatcttc	tctcgattca	gtccctgcaa	gaaccgcgat	tggggacaac	840
aaaggaagta	gaagaagtca	gtattactaa	tatcatcaac	agctccatct	ccagctttaa	900
acggaagatc	agcttttgcca	gcattgaaat	ttccagcgac	aacgttgact	acagtgaact	960
gacaatgaaa	accagcgaca	agttaaagtt	tgtcttccga	gaaaagatgg	gcaggattgt	1020
tgattatttc	acaattcaaa	accccgagta	tggtgatcac	tattccaaac	tactgtttcc	1080
tttgattttt	atgctagcca	atgtatttta	ctgggcatcc	tacatgtatt	tttga	1135

&lt;210&gt; 38

&lt;211&gt; 1323

&lt;212&gt; DNA

&lt;213&gt; Homo sapien

&lt;400&gt; 38

atgaactaca	gcttccactt	ggccttcgtg	tgtctgagtc	tcttactga	gaggatgtgc	60
atccagggga	gtcagttcaa	cgctcaggtc	ggcagaagtg	acaagcttcc	cctgcctggc	120
tttgagaacc	tcacagcagg	atataacaaa	tttctcaggc	ccaattttgg	tggagaaccc	180
gtacagatag	cgctgactct	ggacattgca	agtatctcta	gcatttcaga	gagtaacatg	240
gactacacag	ccaccatata	cctccgacag	cgctggatgg	accagcggct	ggtgtttgaa	300
ggcaacaaga	gcttactctt	ggatgcccgc	ctcgtggagt	tcctctgggt	gccagatact	360
tacattgtgg	agtccaagaa	gtccttcctc	catgaagtca	ctgtgggaaa	caggctcatc	420
cgccctcttct	ccaatggcac	ggtcctgtat	gccctcagaa	tcacgacaac	tggtgcatgt	480
aacatggatc	tgtctaaata	ccccatggac	acacagacat	gcaagttgca	gctggaaagc	540
tggggctatg	atggaaatga	tgtggagttc	acctggctga	gagggaaacga	ctctgtgcgt	600
ggactggaac	acctgcggct	tgctcagtac	accatagagc	ggatatttcac	cttagtcacc	660
agatcgagc	aggagacagg	aaattacact	agattgggtct	tacagtttga	gcttcggagg	720
aatgttctgt	atttcatttt	ggaaacctac	gttccctcca	ctttcctggg	ggtgttgtcc	780



```

tgggtttcat tttggatctc tctcgattca gtccctgcaa gaacctgcat tggagtgcag 840
accgtgttat caatgaccac actgatgatc ggggtcccgca cttctcttcc caacaccaac 900
tgcttcatca aggccatcga tgtgtacctg gggatctgct ttagctttgt gtttggggcc 960
ttgctagaat atgcagttgc tcaactacagt tccttacagc agatggcagc caaagatagg 1020
gggacaacaa aggaagtaga agaagtcagt attactaata tcatcaacag ctccatctcc 1080
agcttttaaac ggaagatcag ctttgccagc attgaaattt ccagcgacaa cgttgactac 1140
agtgacttga caatgaaaac cagcgacaag ttcaagtttg tcttccgaga aaagatgggc 1200
aggattgttg attatttcac aattcaaac cccagtaatg ttgatcacta ttccaaacta 1260
ctgtttcctt tgatttttat gctagccaat gtattttact gggcatacta catgtatttt 1320
tga 1323

```

&lt;210&gt; 39

&lt;211&gt; 440

&lt;212&gt; PRT

&lt;213&gt; Homo sapien

&lt;400&gt; 39

```

Met Asn Tyr Ser Leu His Leu Ala Phe Val Cys Leu Ser Leu Phe Thr
  1          5          10          15
Glu Arg Met Cys Ile Gln Gly Ser Gln Phe Asn Val Glu Val Gly Arg
  20          25          30
Ser Asp Lys Leu Ser Leu Pro Gly Phe Glu Asn Leu Thr Ala Gly Tyr
  35          40          45
Asn Lys Phe Leu Arg Pro Asn Phe Gly Gly Glu Pro Val Gln Ile Ala
  50          55          60
Leu Thr Leu Asp Ile Ala Ser Ile Ser Ser Ile Ser Glu Ser Asn Met
  65          70          75          80
Asp Tyr Thr Ala Thr Ile Tyr Leu Arg Gln Arg Trp Met Asp Gln Arg
  85          90          95
Leu Val Phe Glu Gly Asn Lys Ser Phe Thr Leu Asp Ala Arg Leu Val
 100          105          110
Glu Phe Leu Trp Val Pro Asp Thr Tyr Ile Val Glu Ser Lys Lys Ser
 115          120          125
Phe Leu His Glu Val Thr Val Gly Asn Arg Leu Ile Arg Leu Phe Ser
 130          135          140
Asn Gly Thr Val Leu Tyr Ala Leu Arg Ile Thr Thr Thr Val Ala Cys
 145          150          155          160
Asn Met Asp Leu Ser Lys Tyr Pro Met Asp Thr Gln Thr Cys Lys Leu
 165          170          175
Gln Leu Glu Ser Trp Gly Tyr Asp Gly Asn Asp Val Glu Phe Thr Trp
 180          185          190
Leu Arg Gly Asn Asp Ser Val Arg Gly Leu Glu His Leu Arg Leu Ala
 195          200          205
Gln Tyr Thr Ile Glu Arg Tyr Phe Thr Leu Val Thr Arg Ser Gln Gln
 210          215          220
Glu Thr Gly Asn Tyr Thr Arg Leu Val Leu Gln Phe Glu Leu Arg Arg
 225          230          235          240
Asn Val Leu Tyr Phe Ile Leu Glu Thr Tyr Val Pro Ser Thr Phe Leu
 245          250          255
Val Val Leu Ser Trp Val Ser Phe Trp Ile Ser Leu Asp Ser Val Pro
 260          265          270
Ala Arg Thr Cys Ile Gly Val Thr Thr Val Leu Ser Met Thr Thr Leu
 275          280          285
Met Ile Gly Ser Arg Thr Ser Leu Pro Asn Thr Asn Cys Phe Ile Lys
 290          295          300
Ala Ile Asp Val Tyr Leu Gly Ile Cys Phe Ser Phe Val Phe Gly Ala
 305          310          315          320
Leu Leu Glu Tyr Ala Val Ala His Tyr Ser Ser Leu Gln Gln Met Ala

```

			325				330				335				
Ala	Lys	Asp	Arg	Gly	Thr	Thr	Lys	Glu	Val	Glu	Glu	Val	Ser	Ile	Thr
			340				345				350				
Asn	Ile	Ile	Asn	Ser	Ser	Ile	Ser	Ser	Phe	Lys	Arg	Lys	Ile	Ser	Phe
			355				360				365				
Ala	Ser	Ile	Glu	Ile	Ser	Ser	Asp	Asn	Val	Asp	Tyr	Ser	Asp	Leu	Thr
			370				375				380				
Met	Lys	Thr	Ser	Asp	Lys	Phe	Lys	Phe	Val	Phe	Arg	Glu	Lys	Met	Gly
			385				390				395				
Arg	Ile	Val	Asp	Tyr	Phe	Thr	Ile	Gln	Asn	Pro	Ser	Asn	Val	Asp	His
			405				410				415				
Tyr	Ser	Lys	Leu	Leu	Phe	Pro	Leu	Ile	Phe	Met	Leu	Ala	Asn	Val	Phe
			420				425				430				
Tyr	Trp	Ala	Tyr	Tyr	Met	Tyr	Phe								
			435				440								

```
<210> 40
<211> 289
<212> PRT
<213> Homo sapien
```

	<400>	40														
Met	Asn	Tyr	Ser	Leu	His	Leu	Ala	Phe	Val	Cys	Leu	Ser	Leu	Phe	Thr	
1			5						10					15		
Glu	Arg	Met	Cys	Ile	Gln	Gly	Ser	Gln	Phe	Asn	Val	Glu	Val	Gly	Arg	
			20					25					30			
Ser	Asp	Lys	Leu	Ser	Leu	Pro	Gly	Phe	Glu	Asn	Leu	Thr	Ala	Gly	Tyr	
		35					40					45				
Asn	Lys	Phe	Leu	Arg	Pro	Asn	Phe	Gly	Gly	Glu	Pro	Val	Gln	Ile	Ala	
	50					55					60					
Leu	Thr	Leu	Asp	Ile	Ala	Ser	Ile	Ser	Ser	Ile	Ser	Glu	Ser	Asn	Met	
65					70					75					80	
Asp	Tyr	Thr	Ala	Thr	Ile	Tyr	Leu	Arg	Gln	Arg	Trp	Met	Asp	Gln	Arg	
				85					90					95		
Leu	Val	Phe	Glu	Gly	Asn	Lys	Ser	Phe	Thr	Leu	Asp	Ala	Arg	Leu	Val	
			100					105					110			
Glu	Phe	Leu	Trp	Val	Pro	Asp	Thr	Tyr	Ile	Val	Glu	Ser	Lys	Lys	Ser	
		115					120					125				
Phe	Leu	His	Glu	Val	Thr	Val	Gly	Asn	Arg	Leu	Ile	Arg	Leu	Phe	Ser	
	130					135					140					
Asn	Gly	Thr	Val	Leu	Tyr	Ala	Leu	Arg	Ile	Thr	Thr	Thr	Val	Ala	Cys	
145					150					155					160	
Asn	Met	Asp	Leu	Ser	Lys	Tyr	Pro	Met	Asp	Thr	Gln	Thr	Cys	Lys	Leu	
				165					170					175		
Gln	Leu	Glu	Ser	Trp	Gly	Tyr	Asp	Gly	Asn	Asp	Val	Glu	Phe	Thr	Trp	
			180					185					190			
Leu	Arg	Gly	Asn	Asp	Ser	Val	Arg	Gly	Leu	Glu	His	Leu	Arg	Leu	Ala	
		195					200					205				
Gln	Tyr	Thr	Ile	Glu	Arg	Tyr	Phe	Thr	Leu	Val	Thr	Arg	Ser	Gln	Gln	
	210					215					220					
Glu	Thr	Gly	Asn	Tyr	Thr	Arg	Leu	Val	Leu	Gln	Phe	Glu	Leu	Arg	Arg	
225					230					235					240	
Asn	Val	Leu	Tyr	Phe	Ile	Leu	Glu	Thr	Tyr	Val	Pro	Ser	Thr	Phe	Leu	
				245					250					255		
Val	Val	Leu	Ser	Trp	Val	Ser	Phe	Trp	Ile	Ser	Leu	Asp	Ser	Val	Pro	
		260						265					270			
Ala	Arg	Thr	Arg	Ile	Gly	Asp	Asn	Lys	Gly	Ser	Arg	Arg	Ser	Gln	Tyr	
		275					280					285				

Tyr

<210> 41  
 <211> 265  
 <212> PRT  
 <213> Homo sapien

<400> 41  
 Met Asn Tyr Ser Leu His Leu Ala Phe Val Cys Leu Ser Leu Phe Thr  
 1 5 10 15  
 Glu Arg Met Cys Ile Gln Gly Ser Gln Phe Asn Val Glu Val Gly Arg  
 20 25 30  
 Ser Asp Lys Leu Ser Leu Pro Gly Phe Glu Asn Leu Thr Ala Gly Tyr  
 35 40 45  
 Asn Lys Phe Leu Arg Pro Asn Phe Gly Gly Glu Pro Val Gln Ile Ala  
 50 55 60  
 Leu Thr Leu Asp Ile Ala Ser Ile Ser Ser Ile Ser Glu Ser Asn Met  
 65 70 75 80  
 Asp Tyr Thr Ala Thr Ile Tyr Leu Arg Gln Arg Trp Met Asp Gln Arg  
 85 90 95  
 Leu Val Phe Glu Gly Asn Lys Ser Phe Thr Leu Asp Ala Arg Leu Val  
 100 105 110  
 Glu Phe Leu Trp Val Pro Asp Thr Tyr Ile Val Glu Ser Lys Lys Ser  
 115 120 125  
 Phe Leu His Glu Val Thr Val Gly Asn Arg Leu Ile Arg Leu Phe Ser  
 130 135 140  
 Asn Gly Thr Val Leu Tyr Ala Leu Arg Ile Thr Thr Thr Val Ala Cys  
 145 150 155 160  
 Asn Met Asp Leu Ser Lys Tyr Pro Met Asp Thr Gln Thr Cys Lys Leu  
 165 170 175  
 Gln Leu Glu Ser Trp Gly Tyr Asp Gly Asn Asp Val Glu Phe Thr Trp  
 180 185 190  
 Leu Arg Gly Asn Asp Ser Val Arg Gly Leu Glu His Leu Arg Leu Ala  
 195 200 205  
 Gln Tyr Thr Ile Glu Arg Tyr Phe Thr Leu Val Thr Arg Ser Gln Gln  
 210 215 220  
 Glu Thr Gly Asn Tyr Thr Arg Leu Val Leu Gln Phe Glu Leu Arg Arg  
 225 230 235 240  
 Asn Val Leu Tyr Phe Ile Leu Asp Leu Ser Arg Phe Ser Pro Cys Lys  
 245 250 255  
 Asn Leu His Trp Gly Gln Gln Arg Lys  
 260 265

<210> 42  
 <211> 574  
 <212> DNA  
 <213> Homo sapien

<220>  
 <221> misc\_feature  
 <222> (1)...(574)  
 <223> n = A,T,C or G

<400> 42  
 accaacanag cttagtaatt tctaaaaaga aaaaatgatac tttttccgac ttctaaacaa 60  
 gtgactatac tagcataaat cattcttcta gtaaaacagc taaggatatag acatttcta 120  
 aatttgggaa aacctatgat tacaagttaa aactcagaaa tgcaaagatg ttgggttttt 180

```

gtttctcagt ctgcttttagc ttttaactct ggaaacgcat gcacactgaa ctctgctcag      240
tgctaaacag tcaccagcag gtctctcagg gtttcagccc taaaatgtaa aacctggata      300
atcagtgtat gttgcaccag aatcagcatt ttttttttaa ctgcaaaaaa tgatggctc      360
atctctgaat ttatatctct cattcttttg aacatactat agctaataa ttttatgttg      420
ctaaattgct tctatctagc atgttaaaca aagataatat actttcgatg aaagtaaatt      480
ataggaaaaa aattaactgt tttaaaaaga acttgattat gttttatgat ttcaggcaag      540
tattcatttt taacttgcta cctactttta aata                                     574

```

```

<210> 43
<211> 467
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(467)
<223> n = A,T,C or G

```

```

<400> 43
tttttttttt ttttttattg ccatcaattt attaaaataa acatgtatag caggtttcaa      60
caattgtctt gtagtttgta gtaaaaagac ataagaaaga gaagggtgtg tttgcagcaa      120
tccgtagctg gtttctcacc ataccctgca gttctgtgag ccaaaggctc tgcagaaagt      180
taaaataaat cacaaagact gctgtcatat attaatgca taaacacctc aacattgctc      240
anagtttcat cgttttggtt aanaaaacat tccttcaatt catctatggc attttagtg      300
gcattgtcgt ctatgaactc ttgaagaagt tctttgtatt cagtcttaga cacttgtgga      360
ttgattgtct tggaaatcac attctccaat aaggggcagc cagagcctgc gtagcagtgc      420
tgggagaggg ccgccagcat gaggaccatc agcaacttca tgggtgag                    467

```

```

<210> 44
<211> 613
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(613)
<223> n = A,T,C or G

```

```

<400> 44
tttttttttt ttttttttag ttttaaaata ttttcaactt attattatgc ttataatatt      60
atccaacag actgtattaa aggcagtgat cactaacaca gaacacgaca ggcgaagag      120
gcagccgggc cgattgcagg acgtggcctg tcgggccagg gtcgctgaca tgcacgctgg      180
tagctcatal actgctacct tcagcacagg ctgcaggaat agggacaaga cagatgccgc      240
cggactctta gaagctatth aataaatatc atccaaaaac aaaatggaaa agaaacaaga      300
aacctccga gcacaaccac cttaggccaa ctgaatgtaa tctagttht tcaacaaaa      360
attgagagag aaggaaaata ttgaaacaaa caaacgaaag aaagcagttc ttaagactag      420
cagtaataaa atttatacaa cagttcggtc tgtataatat gatgaaataa atctacatct      480
tttcttattt tggngctttg aattatacat acaaaacaaca attacagga cttgttcaca      540
aagcatgtag gcctanaaaa aggtctctctg aaacctcaa tggcaactgg tgaacggtaa      600
cactgattgc cca                                     613

```

```

<210> 45
<211> 334
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature

```

<222> (1)...(334)

<223> n = A,T,C or G

<400> 45

accagaccaa	gtgaatgcga	cagggaatta	tttcctgtgt	tgataattca	tgaagtagaa	60
cagtataatc	aaaatcaatt	gtatcatcat	tagttttcca	ctgcctcaca	ctagttagct	120
gtgccaagta	gtagtgtgac	acctgtgttg	tcattttcca	catcacgtaa	gagcttccaa	180
ggaaagccaa	atcccagatg	agttctcagag	agggatcaat	atgtccatga	ttatcaggta	240
tgctgactat	ttccaagggg	tttttcagtt	gcttcatttg	cttgtaaagc	aggtaatcct	300
cttggtgtnt	tttctttttc	tcgatgagcc	gtgt			334

<210> 46

<211> 429

<212> DNA

<213> Homo sapien

<220>

<221> misc\_feature

<222> (1)...(429)

<223> n = A,T,C or G

<400> 46

acaattttnt	taaacaagca	gaatagcact	aggcagaata	aaaaattgca	cagacgtatg	60
caattttcca	agatagcatt	ctttaaattc	agtattcagc	ttccaaagat	tggttgccca	120
taatagactt	aaacataata	tgatggctaa	aaaaataaag	tatacgaaaa	tgtaaaaaag	180
gaaatgtaag	tccactctca	atctcataaa	agggtgagag	aaggatgcta	aagcaaaaata	240
aatgtagggt	ctttttttct	atttcggtt	atcatgcagt	ctgcttcttt	gatatgcctt	300
agggttacc	atttaagtta	gaggttgtaa	tgcaatggtg	ggaatgaaaa	ttgatcaaat	360
atacaccttg	tcatttcatt	tcaaattgcg	gntggaaact	tccaaaaaaa	gggtaggcat	420
gaagaaaaa						429

<210> 47

<211> 394

<212> DNA

<213> Homo sapien

<220>

<221> misc\_feature

<222> (1)...(394)

<223> n = A,T,C or G

<400> 47

acgcgaantt	gtgttatgac	tgatagcctt	cagctacaaa	angataggac	tgacctggtt	60
taaagtgttc	tattttgtaa	atcattccat	ttgagtcttt	ctgatgaact	tggtataact	120
gaaatctgtt	attttagtga	ggctccaaaa	tgagcaaaagc	taggcctgat	tagagtagag	180
tgactattaa	aaaacataac	tttctaggag	ctataaatca	aagttttaaa	aagatgtttg	240
gatatatattg	agtattccga	tcagtgaaac	agaaattgcc	ctgcctacta	caaggacaga	300
ctgatgggaa	attatgcacc	tggtcaactt	agcttttaag	cagacgatgc	tgtaaaaaaca	360
aacggcttct	ctgatattta	ttgtaagttt	tagt			394

<210> 48

<211> 486

<212> DNA

<213> Homo sapien

<400> 48

acaaggaac	cgaggggtga	ccacctctga	gatgtccttg	actttgtcat	agcctggggc	60
atattgagca	tctctctcac	agctgccttt	cttatcccca	ttcttgatgt	agacctcctt	120

```

ccgagtcagc tttttctcct cctcagacac aaacagagct ttgatatacct gtgcagggag      180
cagctcttcc ttttgttgct ggcaagtggg agttggagga agcctcaaag ctcgagttgt      240
tccctcggtg caggggagac aaatgggcct gatagtctgg ccatatttca gcttattctt      300
gagcttgatc agggcaacgt catagtcata aaattcagga attcctgctt cttttttccc      360
attaatgttg tagttggggg gaaataggac tacttctatc tccaggtecc gcttctcccc      420
tcccttgatt gagtgttcc tgtcatccac agtgaaacaa tgtgtgtgctg tcagcacaaa      480
gtacct                                           486

```

&lt;210&gt; 49

&lt;211&gt; 487

&lt;212&gt; DNA

&lt;213&gt; Homo sapien

&lt;400&gt; 49

```

acgggctgac agagaagatt cccgagagta aatcatcttt ccaatccaga ggaacaagca      60
tgtctctctg ccaagatcca tctaaactgg agtgatgta gcagaccag cttagagttc      120
ttctttcttt cttaagccct ttgctctgga ggaagtcttc cagcttcagc tcaactcaca      180
gcttctccaa gcatcaccct gggagtttcc tgagggtttt ctcataaatg agggctgcac      240
attgctgttt ctgcttcgaa gtattcaata ccgctcagta ttttaaatga agtgattcta      300
agatttggtt tgggatcaat aggaaagcat atgcagccaa ccaagatgca aatgttttga      360
aatgatatga ccaaaatttt aagtaggaaa gtcacccaaa cacttctgct ttcacttaag      420
tgtctggccc gcaatactgt aggaacaagc atgatcttgt tactgtgata ttttaaatat      480
ccacagt                                           487

```

&lt;210&gt; 50

&lt;211&gt; 460

&lt;212&gt; DNA

&lt;213&gt; Homo sapien

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (1)...(460)

&lt;223&gt; n = A,T,C or G

&lt;400&gt; 50

```

acatattttg gttgaagaca ccagactgaa gtaaacagct gtgcatccaa tttattatag      60
ttttgtaagt aacaatatgt aatcaaactt ctagggtgact tgagagtgga acctcctata      120
tcattattta gcaccgttta tgacagtaac catttcagtg tattgtttat tataccactt      180
atatcaactt atttttcacc aggttaaaat tttaatttct acaaaataac attctgaatc      240
aagcacactg tatgttcagt aggttgaact atgaacactg tcatcaatgt tcagttcaaa      300
agcctgaaag tttagatcta gaagctggta aaaatgacaa tatcaatcac attaggggaa      360
ccattgttgt cttcacttaa tccatttagc actattgaaa ataagcacac caagntatat      420
gactaatata acttgaaaat tttttatact gagggggtng      460

```

&lt;210&gt; 51

&lt;211&gt; 529

&lt;212&gt; DNA

&lt;213&gt; Homo sapien

&lt;400&gt; 51

```

acacttgaaa ccaaatttct aaaacttggt tttcttaaaa aatagttggt gtaacattaa      60
accataacct aatcagtggt ttcactatgc ttccacacta gccagtcttc tcacacttct      120
tctggtttca agtctcaagg cctgacagac agaagggtct ggagattttt tttctttaca      180
attcagttct cagcaacttg agagctttct tcatgttgct aagcaacaga gctgtatctg      240
caggttcgta agcatagaga cggtttgaat atcttccagt gatatcggct ctaactgtca      300
gagatgggtc aacaaacata atcctgggga catactggcc atcaggagaa aggtgtttgt      360
cagttgtttc ataaaccaga ttgaggagga caaactgctc tgccaatttc tggattttct      420
tattttcagc aaacactttc tttaaagctt gactgtgtgg gcactcatcc aagtgatgaa      480

```

taaatcatca aggggttgggt gcttgtcttg gatttatata gagcttctt 529

<210> 52  
 <211> 379  
 <212> DNA  
 <213> Homo sapien

<400> 52  
 acttttccaa gcagtaaagg atccaggaga tagcactgga tgtgggtgtca tgtcctgcaa 60  
 acatgaacgt tttcacttca gcctggagat ctgcttcaga gaaatctttg gtgttttcgc 120  
 tttttggcact caaaagtatg tccagaaaat cccagcgccct tttctgagta gtatcttggt 180  
 ttagcttata ctttaagagac tccttccgggt cctggattac tttctctgtg aactgatgaa 240  
 gttcttggtt aaatttagaa aagatttggc cttgagagct gaatttgaaa accaggtcgt 300  
 tgtgatgtag aaaattgttc atgcgctgggt tggagatttt gctaagggtg aacactgctt 360  
 tcaggatatga gtccagggt 379

<210> 53  
 <211> 380  
 <212> DNA  
 <213> Homo sapien

<220>  
 <221> misc\_feature  
 <222> (1)...(380)  
 <223> n = A,T,C or G

<400> 53  
 acttttatct taaaagggtg gtagttttcc ctaaaatact tattatgtaa gggtcattag 60  
 acaaatgtct tgaagtagac atggaattta tgaatgggtc tttatcattt ctcttcccc 120  
 tttttggcat cctggcttgc ctccagtttt aggtccttta gtttgcttct gtaagcaacg 180  
 ggaacacctg ctgagggggc tctttccctc atgtatactt caagtaagat caagaatctt 240  
 ttgtgaaatt atagaaatn actatgtaaa tgcttgatgg aatnntttcc tgctagtgtg 300  
 gcttctgaaa ggcgctttct ccattttattt aaaactacc atgcaattaa aaggtacctt 360  
 gccgcgacca cnctaanggc 380

<210> 54  
 <211> 245  
 <212> DNA  
 <213> Homo sapien

<400> 54  
 gcgcggcgct tcacttcttc aacttccgggt ccggctcgcc cagcgcgctg cgagtgcctg 60  
 ccgaggtgca ggaggccgc gcgtggatta atccaaaaga gggatgtaaa gttcacgtgg 120  
 tcttcagcac agagcgctac aaccagagct ctttacttca ggaagggtgag ggacgtttgg 180  
 ggaatgttc tgctcgagtg tttttcaaga atcagaaacc cagaccaacc atcaatgtaa 240  
 cttgt 245

<210> 55  
 <211> 556  
 <212> DNA  
 <213> Homo sapien

<400> 55  
 acagaagatg aataataatg aaaaactgtg attttttgac tatcacatac atttgtgttaa 60  
 aaaacaggta aatataatga ctattactgt taagaaagac aaggaggaaa actgtttcaa 120  
 tgttcagggt taaatactaa gcacaaaaat ataacaaatt ctgtgtctac aataattttt 180  
 gaagtgtata caagtgcatt gcaaatgagc tcttttaaat ttaaagtcca tttccctttt 240  
 agccaagcat atgtctacat ttatgatttc tttctcttat tttaaagtct cttctggttt 300

```

agttttttaa aaagtttcat catggctgtc atcttggaaat ctagcctcca gctcaaagct    360
gagacttcac gcatacatat tctcctttct gggtgcatct tcacctagtt tctccaagta    420
ttcagagtta aatagcaca cttcttttat atgttcactt ttgtccacat gtagtggcag    480
tgctgctgct tcagtaggct ttctcacaca cctctttcct tctttcaaca gcagtcacca    540
aacgttcaca acacaa                                     556

```

```

<210> 56
<211> 166
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(166)
<223> n = A,T,C or G

```

```

<400> 56
atgggccctg attacatcat tatgaactac tcagggnaac atcccaaata cgcacctngg    60
gaaagacttg gtccgagatg tggatcatcca tacaggctac ctcttcacaga gncagggncc  120
caagagctgc ntnatcacct acctggccca ggtggacccc anaggg                    166

```

```

<210> 57
<211> 475
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(475)
<223> n = A,T,C or G

```

```

<400> 57
acatccncat gttcctccaa atgacgtttg gggctcctgct tgccaacatt ctttattgcc    60
agctgttcag gtgtcatctt atcttcttct tctacagcct tattgtaatt ctgggctaatt  120
tccaacatct cttttaccac tgattcattg cgtttacaat gttcactgta gtccctgaagt  180
gtcaaacctt ccacccaact cttcttatgc aaatttagca acatcttctg ttccagttca    240
tttttccgat agttaatagt aatggagtaa taatgtctgt ttagtccatg aattaatgcc    300
tggatagatg gcttggttaa gtgaccacaga ttcgaaagtg tttgtcttgg ttcatgtcct  360
aagaccatca tattagcatt gatcaatctg aaggcatcaa taacaacctt tctttttaca  420
ctctgaatgg gatccacaac cactgccaca gntctctccg ataaggcttc aaagc        475

```

```

<210> 58
<211> 520
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(520)
<223> n = A,T,C or G

```

```

<400> 58
actgttnatg tgctacttgc atttgtccct cttcctgtgc actaaagacc ccactcactt    60
ccctagtgtt cagcagtgga tgacctctag tcaagacctt tgcactagga tagttaatgt  120
gaaccatggc aactgatcac aacaatgtct ttcagatcag atccatttta tctccttgt  180
tttacagcaa gggatattaa ttacctatgt tacctttccc tgggactatg aatgtgcaaa  240
attccaatgt tcatggtctc tccctttaaa cctatatctc acccctttta cattatagaa  300
aggaatgctg gaaaccacaga gtccttctct tgggactctt aatgtgtatt tctaattatc  360

```



catgactctt aatgtgcata ttttcaattg cctaattgat ttcaattgtc taagacattt	420
caaatgtcta attggggaga actgagtctt ttatatcaag ctaatatcta gcttttata	480
caagctaata tcttgacttc tcagcatcat agaagggggt	520

<210> 59  
 <211> 214  
 <212> DNA  
 <213> Homo sapien

<220>  
 <221> misc\_feature  
 <222> (1)...(214)  
 <223> n = A,T,C or G

<400> 59	
ctggcaggaa atgcatcaaa agacttaaag gtanagecga ttacccctcg tcacttgcaa	60
cttgctattc gtggagatga agaattggat tctctcatca aggtacaat tgctggtggn	120
ggtgtcattc cacacateca caaatctctg atngggaana aaggacaaca naagactgnc	180
taanggatgc ctgnatnctt tggaatctca tgac	214

<210> 60  
 <211> 360  
 <212> DNA  
 <213> Homo sapien

<220>  
 <221> misc\_feature  
 <222> (1)...(360)  
 <223> n = A,T,C or G

<400> 60	
gcatacaaca tggcagcagg gcctcgggaa gangggtagg aggaccgagc agcattctct	60
gtagaggaag acaggaaagg agaccctctt ggcacacatt tatggagggt tgtccctgaa	120
gagaagggca ggtgggagag gttccctggt acttaagaga aggcaccagt ggcaaagagc	180
acaatgaaga ggatgatgat aaaaacaatc acgcagataa ggacaatcat cttcacgttc	240
ttccaccaga attttcgagc caccttctgc gatgtcgtct tgaagtgtc agatgtgggt	300
tccagatcct ctgtcttgtt gcggagatgt tccaagtttt ccccccgggc caggatccgc	360

<210> 61  
 <211> 391  
 <212> DNA  
 <213> Homo sapien

<220>  
 <221> misc\_feature  
 <222> (1)...(391)  
 <223> n = A,T,C or G

<400> 61	
tntgggatcg tactcgatta aacagagcca cttttgttcc tgaggcaatg cataantcan	60
catttttcaa tgactgcttc tttttggaag gnttggagat gacttttatc cgcttgctga	120
ggaacacacc aatgncatca ctgttgccat agaacatctt tacagacaac atgaantgct	180
ttcgcttgct tgagtcagat atatacaatg ttttggtctg gcaatagtgc tttccttcca	240
agtttagctg ctgcatttct tggncactat ttcctatccc aataaatgca cacggttgag	300
actcttgntc agaacaacca tcncgttcca tttgttcttt ttttntcttc catccactgc	360
ccataagata tacacannga ggtgggcaaa a	391

<210> 62

<211> 324  
<212> DNA  
<213> Homo sapien

<220>  
<221> misc\_feature  
<222> (1)...(324)  
<223> n = A,T,C or G

<400> 62  
acaattttat ttttaacagat ttcaagagtc cattttttta aaaatgagca ataaagaacc 60  
tctatcagtg agacttctca ttttatagca aatacatttt tgcagcttaa attttcttga 120  
attcatatac gcttctgtca ttttaacaaa cttccagaga aaactggtct ctatataatt 180  
aagtaacaaa tttgacaaaa tacatatatta tacatatata ganctctaata ataaatatta 240  
aatttgaaaa aatcaaatgt gaagcagaaa ctgctataca agtatattgt ntaatatcta 300  
tntnatacat taaagnnttc cggg 324

<210> 63  
<211> 360  
<212> DNA  
<213> Homo sapien

<220>  
<221> misc\_feature  
<222> (1)...(360)  
<223> n = A,T,C or G

<400> 63  
acaganncc tgaatatgtt gtggttccct cattatggcc cttcattccc ttctgtgtta 60  
atagtaaagc atgttgccca ataaactaca ccctgaccaa atttgggccc ggatctcatg 120  
ggtcacgtgg agtttttaaat acgattttta atttacttgg gtaattgagc tgaatcttta 180  
gttttcagat tactttttta aacagatagg ctcttagaac aaattattaa aaacataata 240  
ccccattgga ggggaatctg gattaactac ccaactgttc ccccccccc aacttttgaa 300  
aaattttggc catatagaat gcatgaaaaa tcaggataga tcttatgagg actttatagt 360

<210> 64  
<211> 491  
<212> DNA  
<213> Homo sapien

<220>  
<221> misc\_feature  
<222> (1)...(491)  
<223> n = A,T,C or G

<400> 64  
nctgactgtg atgtccactt gttccctgat ttttacacat catgtcaaag ataacagctg 60  
ttcccaccca ccagttccctc taagcacata ctctgetttt ctgtcaacat cccatttttg 120  
ggaaaggaaa agtcatattt attcccgcac ccagttttt taacttggtc tcccagttgt 180  
ccccctcttc tctgggtgta agaagggaaa ttggaaaaaa attatatata tattctcctt 240  
ttaatggtgg ggggtactct gagaggagag acagcaagtc caccctaact tgttacacag 300  
cacataccac aggttctgga attctcatct tcgaacctag agaaataggt gctataaaca 360  
gggaattaaag caaaatgctg gatgctatag atcttttaatt tgncttaatt ttttttctat 420  
tattaaacta caggctgtag atntcttagg tctcacagaa cttntatcat tttaaactga 480  
cttgatatt t 491

<210> 65  
<211> 484

<212> DNA  
<213> Homo sapien

<220>  
<221> misc\_feature  
<222> (1)...(484)  
<223> n = A,T,C or G

<400> 65  
accagcacac cggcgccgctc ctggactgcg ccttctacga tccaacgcat gcctggagtg 60  
gaggactaga tcatcaattg aaaatgcatg atttgaacac tgatcaagaa aatcttggtg 120  
ggacccatga tgcccctatc agatgtgttg aatactgtcc agaagtgaat gtgatggtca 180  
ctggaagtgt ggatcagaca gctaaactgt gggatcccag aactccttgt aatgctggga 240  
ccttctctca gcctgaaaag gtatataccc tctcagtgtc tggagaccgg ctgattgtgg 300  
gaacagcagg ccgcagagng ttggtgtggg acttacggaa catgggttac gtgcagcagc 360  
gcaggggagt cagcctgaaa taccagactc gctgcatacg agcgtttcca aacaagcagg 420  
gttatgtatt aagctctatt gaaggccgag tggcagttga gtatttgac ccaagccctg 480  
aggt 484

<210> 66  
<211> 355  
<212> DNA  
<213> Homo sapien  
  
<220>  
<221> misc\_feature  
<222> (1)...(355)  
<223> n = A,T,C or G

<400> 66  
ngaagaaagt atgggtggag gtgaaggtaa tcacagagct gctgattctc aaaacagtgg 60  
tgaaggaaat acaggtgctg cagaatcttc tttttctcag gaggtttcta gagaacaaca 120  
gccatcatca gcatttgaaa gacaggcccc tcgagcacct cagtcaccga gacgccacc 180  
acatccactt cccccaagac tgaccattca tgcccaccc caggagttgg gaccaccagt 240  
tcagagaatt cagatgaccc gaaggcagtc tgtaggagct ggccttcagt tgactccagg 300  
aataggtggc acgcaacagc atttttttga tgatgaagac agaacagttc caagt 355

<210> 67  
<211> 417  
<212> DNA  
<213> Homo sapien

<400> 67  
acgacacccc tcaagagggtg gcogaagctt tcctgtcttc cctgacagag accatagaag 60  
gagtcgatgc tgaggatggg cacagcccag gggacaaca gaagcggaag atcgctctgg 120  
acccttcagg ctccatgaac atctacctgg tgctagatgg atcagacagc attggggcca 180  
gcaacttcac aggagccaaa aagtgtctag tcaacttaat tgagaagggtg gcaagttatg 240  
gtgtgaagtc aagatatggt ctagtacat atgccacata ccccaaaatt tgggtcaaag 300  
tgtctgaagc agacagcagt aatgcagact gggtcacgaa gcagctcaat gaaatcaatt 360  
atgaagacca caagttgaag tcagggacta acaccaagaa ggccctccag gcagtgt 417

<210> 68  
<211> 223  
<212> DNA  
<213> Homo sapien

<220>  
<221> misc\_feature

<222> (1)...(223)

<223> n = A,T,C or G

<400> 68

cacttgcaag	cttgcttaca	gagacctgnt	aaacaaagaa	cagacagatt	ctataaaatc	60
agttatatca	acatataaag	gagtgtgatt	ttcagtttgt	ttttttaagt	aaatatgacc	120
aaactgacta	aataagaagg	caaaacaaaa	aattatgctt	ccttgacaag	gcctttggag	180
taaacaaaat	gctttaaggc	tectggtgaa	tggggttgca	agg		223

<210> 69

<211> 396

<212> DNA

<213> Homo sapien

<400> 69

accttttttc	tctccaaagg	aacagtttct	aaagttttct	ggggggaaaa	aaaacttaca	60
tcaaatttaa	accatatggt	aaactgcata	ttagttgtgt	tacacaaaaa	aattgcctca	120
gctgatctac	acaagtttca	aagtcattaa	tgcttgatat	aaatttactc	aacattaaat	180
tatcttaa	tattaattaa	aaaaaaaaact	ttctaaggaa	aaataaacaa	atgtagaccg	240
tgattatcaa	aggattatta	aagaatcttt	accaaaaatt	tcaaccctac	aacctaaaac	300
cgcaaatctc	tattttttaa	catcagaaaa	taactcttgg	ttcattactt	atgacccaaa	360
gtttttatct	cactattcaa	tatctgaaaa	gtatca			396

<210> 70

<211> 402

<212> DNA

<213> Homo sapien

<220>

<221> misc\_feature

<222> (1)...(402)

<223> n = A,T,C or G

<400> 70

accannccc	accaggcaa	acagctccga	catgtttngt	aagttagaca	agccagtgc	60
agttttttt	tttttttct	ttttctttt	ttgtctttt	gcttaccttc	ttgcttaatg	120
gaattgttat	ggctaagcac	atagaaggcc	aaaaaaggag	tttttcaaac	ccagcaaata	180
aagtgtcttg	attctgaact	gccaaaagaa	aactgcactt	ccctctttaa	gtaaaacgaa	240
atgagtttct	taggtaaatg	tattcatcag	cccagataaa	aaaaaaacca	gttatgtgag	300
cgttagtcac	tgctcatttc	caggaanata	aaacaaaata	ccagcccagc	cagactcaca	360
tgtgggnata	tatatataaa	gcaagagagc	cacaccacac	ag		402

<210> 71

<211> 385

<212> DNA

<213> Homo sapien

<220>

<221> misc\_feature

<222> (1)...(385)

<223> n = A,T,C or G

<400> 71

accagtagag	agtggccctt	gcaggccact	tataaacagg	aagctctctc	ctgagctcac	60
tgatcaacct	gcccttggca	cagacagaac	ctaccagaaa	agaacaagta	caaaacacta	120
tcattatctg	ttttctcaag	acagtcccaa	atgtccttgt	gcgatcgcca	caaactcagt	180
gattggccca	agtcattccc	gggtgccata	aacagtaact	ggtgtgcanc	attagaacaa	240
ggggacacgg	ccttgattct	cttctgagca	acatgaactg	ggatttctgc	cnccccggat	300

```
ctcggctgcc acctccgaag aagtcgtgac cagccacctc cacagtaaaa gattcctccc 360
gtgagtatga ttggaatgc gncct 385
```

```
<210> 72
<211> 538
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(538)
<223> n = A,T,C or G
```

```
<400> 72
caattaatta acagaggtat aattgtctca ctttcagaag tgatcattta tttttattta 60
gcacagggtca taagaaaaat atatagaaaa ataatcaatt tcatatataa aaggattatt 120
tctccacctt taattattgg cctatcattt gttagtgtta ttgggtcata ttattgaact 180
aatgtattat tccattcaaa gtctttctag atttaaaaa gtatgcaaaa gcttaggatt 240
atatcatgtg taactattat agataacatc ctaaaccctc agtttagata tataattgac 300
tgggtgtaat ctcttttgta atctgntttg acagatttct taaattatgt tagcataatc 360
aaggaagatt taccttgaag cactttccaa attgatactt tcaaacttat tttaaagcag 420
tagaaccttt tctatgaact aagtcacatg caaaactcca acctgtaagt atacataaaa 480
tggacttact tattcctctc accttctcca ggcctaggaa tattcttctc tggagccc 538
```

```
<210> 73
<211> 405
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(405)
<223> n = A,T,C or G
```

```
<400> 73
actttatnna tggaattttc ttctacttgt atccatttnc cggggcttat ggacccattc 60
atactctcca tatttagaat caaagggtcc tttctgaaga gaccttaatt ttaaggtaaa 120
acgtgggtcca agttccctgaa ttcccacttt cttttcactc ctgaatatgt atctgtgaaa 180
tctgaagaat atgtaatccc gttgattgtg gaatgtggca acctgccttc cgataaattg 240
aggattatga ggaaagagag atgcaaacat acgtccaatt gaatgaccca gccgtgttgt 300
aaaattattc agaattattt caggtatgtg ttctgtgggg tccttgccctc ttctcttaat 360
ttctttacga agacgaacac tgctcatttt aaaatgagca gttgg 405
```

```
<210> 74
<211> 498
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1)...(498)
<223> n = A,T,C or G
```

```
<400> 74
tgagccctgc acctgtttcc tgcacccctt gccnactggt tctatggcca caaggagttt 60
taccagtaa aggagtttga ggtgtattat aagctgatgg aaaaataccc atgtgctggt 120
cccttggtgg ttggacctt tacgatgttc ttcagtgtcc atgaccaga ctatgccaaag 180
attctctga aaagacaaga tcccaaaagt gctgttagcc acaaaatcct tgaatcctgg 240
```

```

gttggtcgag gacttgtgac cctggatggt tctaaatgga aaaagcaccg ccagattgtg      300
aaacctggct tcaacatcag cattctgaaa atattcatca ccatgatgtc tgagagtgtt      360
cggatgatgc tgaacaaatg ggaggaacac attgcccaaa actcacgtct ggagctcttt      420
caacatgtct ccctgatgac cctggacagc atcatgaagt gtgccttcag ccaccagggc      480
agcatccagt tggacagt                                     498

```

```

<210> 75
<211> 458
<212> DNA
<213> Homo sapien

```

```

<400> 75
agccttgac atgataactca gattcctcac ccttgcttag gagtaaaaca atatacttta      60
cagggtgata ataatctcca tagttatttg aagtggttg aaaaaggcaa gattgacttt      120
tatgacattg gataaaatct acaaatcagc cctcgagtta ttcaatgata actgacaaac      180
taaattattt ccctagaaaag gaagatgaaa ggagtggagt gtggtttggc agaacaactg      240
catttcacag cttttccagt taaattggag cactgaacgt tcagatgcat accaaattat      300
gcatgggtcc taatcacaca tataaggctg gctaccagct ttgacacagc actgttcatc      360
tggccaaaca actgtgggta aaaacacatg taaaatgctt tttaacagct gatactgtat      420
aagacaaagc caagatgcaa aattaggctt tgattggc                                     458

```

```

<210> 76
<211> 340
<212> DNA
<213> Homo sapien

```

```

<220>
<221> misc_feature
<222> (1)...(340)
<223> n = A,T,C or G

```

```

<400> 76
accttatacc aaaanaatgc ttattccaaa atattttttg tagctagtag ttctttcctt      60
ggaggtaaag aaaatatacc caaactttta attaccagga ttcagaatat ttaagagaac      120
aatTTtagtt aagaatcaaa tatactgaga ttcaaagagg ggaaaaaaag gaaatattat      180
agaagacaaa ggtcaaactg gcattccaga tctggagcaa ttttgtaaag caggaaaaca      240
actatgacaa tctgnagctt cttagatcat tatagtgaat gtncccatTT actataaggg      300
tttttataat ggtgtttcct aaataaagga acataaatgt                                     340

```

```

<210> 77
<211> 405
<212> DNA
<213> Homo sapien

```

```

<400> 77
actccatttg tggaaactcg gtccgagctt ggtaaacagc cgaatgtctt cctcccctac      60
agtttctctt ccttgcattg gagcagtgat gtcttgatta aaggcattaa ttttatctat      120
caggaagaac attttttcat ttctgtcttc cggatgtgac acaccatact tttgtagctc      180
ctctgttatt ctctgggtgag tctccttgat ttgatTTtct aacaggggca gagatttaca      240
gatatgtgtg atgagctcgc tggtaagttt ttctgccagg cagggaaaccg tggcctttcc      300
ttcttcacag agatccctga aatatgggtg gttctcaaag aagatcttct ctctctgcag      360
ggcttcggac aggtctcagct ggtcctggat ctctctgttg ccccg                                     405

```

```

<210> 78
<211> 410
<212> DNA
<213> Homo sapien

```

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (1)...(410)

&lt;223&gt; n = A,T,C or G

&lt;400&gt; 78

acagcagntn	tagatggctg	caacaacctt	cctcctaccc	cagcccagaa	aatatttctg	60
ccccacccca	ggatccggga	ccaaaataaa	gagcaagcag	gcccccttca	ctgaggtgct	120
gggtagggct	cagtgccaca	ttactgtgct	ttgagaaaga	ggaaggggat	ttgtttggca	180
ctttaaaaat	agaggagtaa	gcaggactgg	agaggccaga	gaagatacca	aaattggcag	240
ggagagacca	tttggcgcca	gtcccctagg	agatgggagg	agggagatag	gtatgagggg	300
aggcgctaag	aagagtagga	gggggtccact	ccaagtggca	gggtgctgaa	atgggctagg	360
accaacagga	cactgactct	aggtttatga	cctgtccata	cccgttccac		410

&lt;210&gt; 79

&lt;211&gt; 512

&lt;212&gt; DNA

&lt;213&gt; Homo sapien

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (1)...(512)

&lt;223&gt; n = A,T,C or G

&lt;400&gt; 79

acagtgaaaa	acaaactaat	ataaagcatt	ccagnngata	aaaacctcct	caggcttatg	60
gtttgttttc	caaggaaatt	atgtttcaat	gtaaagtttg	aaatactcca	gacatacatt	120
ccatgtaggt	tttgggtgcc	aatgttaaaa	tttcaaat	ttgcattgcaag	gcttagcaaa	180
gaaacactgg	cagaattcca	gcatttgcaa	aattctaagt	tttgggtgaat	attgtaaata	240
ttacaattgg	tattagaaag	ccatgatgaa	tccagaatta	agagaaaacc	catttcataa	300
atattttgtt	tgattaaaaa	ataccaggct	taccatgttc	taaataacac	aagaaaatat	360
ctttaaaaaa	aaaaggactg	caatttaaca	gtaatctgta	tatcttttagc	tgccattaaa	420
aaaagaaaaa	agaacaacca	aaaacaatga	aaatgttaca	actggtataa	agtnaccna	480
tgatgctccc	cttacgagaa	aacaaaactg	tc			512

&lt;210&gt; 80

&lt;211&gt; 174

&lt;212&gt; DNA

&lt;213&gt; Homo sapien

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (1)...(174)

&lt;223&gt; n = A,T,C or G

&lt;400&gt; 80

tgattcccca	gacctcaaat	gggctaacac	gcttctcttc	tncagcagnc	ttcctgtccg	60
tgaagntncc	ttccagattg	gtacatggaa	ctgaaaacaa	agggagcctc	agctggattg	120
aaatctggag	catgccacaa	agncttgac	tnggcatttt	cnagaagaac	ccat	174

&lt;210&gt; 81

&lt;211&gt; 274

&lt;212&gt; DNA

&lt;213&gt; Homo sapien

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (1)...(274)

<223> n = A,T,C or G

<400> 81

```
ttgcaacaag cacattaaat taaggcctgc tngaatttct tcctcccca tcaggtaaac      60
tttctttgcc aataaagttt gagggagtgg catttgaaaa tctctttaa aaagaagtct      120
tcctctattc acnagaaaac tcaaaaataa ttttcattat caacacaca actaactcaa      180
tctctgcttt aagtttctat tggccaattt ttctgattna tacgagaatt attntcagnt      240
ntagaaaatc ctggtctttg gtcattacaa gntg                                274
```

<210> 82

<211> 101

<212> DNA

<213> Homo sapien

<220>

<221> misc\_feature

<222> (1)...(101)

<223> n = A,T,C or G

<400> 82

```
atggagaaga tcgaacctga gcctnntgag aattgcctgc tacngcctgg cagccctgcc      60
cgagtggccc agcnnccattt cacnagntgg gcatgatttg n                                101
```

<210> 83

<211> 182

<212> DNA

<213> Homo sapien

<400> 83

```
tattatgggg aaagataact gagaataaag ctatcatgca gatatttgca gagataaaag      60
taatgcagat actgagtggg gttttgatca aactatgctt gaaagccact ctaccactag      120
ttacacaaac caataatttc ccttcgcagt ggaagtcagc ttgagttttt tcagggtgtt      180
tt                                                                182
```

<210> 84

<211> 229

<212> DNA

<213> Homo sapien

<220>

<221> misc\_feature

<222> (1)...(229)

<223> n = A,T,C or G

<400> 84

```
actgtttgta gctgcactac aacagattct taccgtctcc acaaaggcca gagattgtaa      60
atggtcaata ctgacttttt ttttatccc ttgactcaag acagctaact tcattttcag      120
aactgtttta aaccttttg tgctggttta taaaataatg tngtaaatcc ttggtgcttt      180
cctgatacca nactgtttcc cgnggttggt tagaatatat tnngttcng                    229
```

<210> 85

<211> 500

<212> DNA

<213> Homo sapien

<220>

<221> misc\_feature

<222> (1)...(500)



<223> n = A,T,C or G

<400> 85

```

ggggagtang tgatttatta aagcaagacg ttgaaacctt tacnttctgc agtgaagatc      60
aggggtgtcat tgaagacag tggaaccag gatgaaagt tttacatgac acacactaca      120
tttcttcaat attttacca ggacttcgc aatgaggctt cgtttctgaa gggacatctg      180
atccgagcat ctcttcaact ctaacttggc tgcaacagct tccagagggg catcaaattt      240
ggcaagactt aacttgaaca gaggttcact aatgaagaag aagtctaaca gctcagaaac      300
aagagctggg cagaactcgg cattggcctg gtagcagcag agggccagcg tgaccagcag      360
gagacacacc gacagcttca tgggtggctt ttttgctgtg agctcagctt tcacaaacaa      420
tgagtgtatt ggactccacc ccaggagcct gtggagctgc agagcccagg gctatttgta      480
cctgccccgg cgngcgtctg

```

<210> 86

<211> 323

<212> DNA

<213> Homo sapien

<220>

<221> misc\_feature

<222> (1) ... (323)

<223> n = A,T,C or G

<400> 86

```

ccgccagtgt gctggaattc gcccttgccg cccgggcagg tactcagaag tcatttgta      60
tttacaattg ggtttgtgtg ggatgggatn tanggcggat gagccagtgc ttttgcaatg      120
aagatgcaat antcattgtc ctctccact gtctctctct tcctcacccc atggcagctn      180
tcatgaccca ttcccaaagg gtccaccgag tcctgaactc agcttcatca ccaacattcc      240
tcgccttcag ttgaattcaa cactgncaan ggagnagang caaagacttg ggtcaggggag      300
agggngggaa acacanaaca aac

```

<210> 87

<211> 230

<212> DNA

<213> Homo sapien

<400> 87

```

gcagcattga gccacccctt tggcaggcga tacggcagct ctgtgccctt ggccagcatg      60
tggagtggag gagatgctgc ccctgtggtt ggaacatcct ggggtgaccc ccgaccagc      120
ctcgctgggc tgtccctgt ccctatctct cactctggac ccagggtga catcctaata      180
aaataactgt tggattagac aaaaaaaaaa aaaaaaaaaa aaaaaaaagg      230

```

<210> 88

<211> 249

<212> DNA

<213> Homo sapien

<220>

<221> misc\_feature

<222> (1) ... (249)

<223> n = A,T,C or G

<400> 88

```

atgtgaccag gtctaggtct ggagtttcag nttggacact gagccaagca gacaagcaaa      60
gcaagccagg acacaccatc ctgccccagg cccagcttct ctctgcctt ccaacgccat      120
ggggagcaat ctcagcccc aactctgct gatgccctt atcttgggct tcttgtctgg      180
aggtgtgacc accactcctt ggtctttggc ccggccccat ggatcctgct ctctggaggg      240
ggtntagat

```

```
<210> 89
<211> 203
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1) ... (203)
<223> n = A,T,C or G

<400> 89
tgtttacact gtcaaggatg acaaggaaaag tgttontatc tntgatacca tcatcccagc      60
tgttcctcct cccactgacc tgcgattcac caacattggt ccagacacca tgcgtgtcac      120
ctgggctcca ccccatcta ttgatttaac taacttcctg gtgcggnact cacctgtgaa      180
aatgangaa gatgttgacag agt                                           203

<210> 90
<211> 455
<212> DNA
<213> Homo sapien

<400> 90
ctctaagggg gctggcaaca tggtcagca ggcttgcccc agagccatgg caaagaatgg      60
acttgtaatt tgcacacctg tgatcacctt actcctggac cagaccacca gccacacatc      120
cagattaaaa gccaggaagc acagcaaacg tcgagtgaag gacaaggatg gagatctgaa      180
gactcaaatt gaaaagctct ggacagaagt caatgccttg aaggaaatc aagccctgca      240
gacagtctgt ctccgaggca ctaaagtcca caagaaatgc taccttgctt cagaagggtt      300
gaagcatttc catgaggcca atgaagactg catttcctaa ggaggaaatc tggttatccc      360
caggaaactcc gacgaaatca acgacctcca agactatggt aaaaggagcc tgccaggtgt      420
caatgacttt tggctgggca tcaatgacat ggtca                               455

<210> 91
<211> 488
<212> DNA
<213> Homo sapien

<400> 91
actttgcttg ctcatatgca tgtagtcact ttataagtca ttgtatgta ttatattccg      60
taggtagatg tgtaacctct tcaccttatt catggctgaa gtcacctctt ggttacagta      120
gcgtagcgtg gccgtgtgca tgtectttgc gcctgtgacc accaccccaa caaacatcc      180
agtgcacaaac catccagtgg aggtttgtcg ggcaccagcc agcgtagcag ggtcgggaaa      240
ggccacctgt cccactccta cgatacgcta ctataaagag aagacgaaat agtgacataa      300
tatattctat ttttatactc ttcctatctt tgtagtgacc tgtttatgag atgctgggtt      360
tctacccaac ggcctcgcag ccagctcacg tccaggttca acccacagct acttggtttg      420
tgttcttctt catattctaa aaccattcca tttccaagca ctttcagtcc aataggtgta      480
ggaaatag                                                                488

<210> 92
<211> 420
<212> DNA
<213> Homo sapien

<220>
<221> misc_feature
<222> (1) ... (420)
<223> n = A,T,C or G
```

<400> 92  
tctccggcag gctctgcccc ggtcgtagcn agnnaaccta taatcctgac cttttttgta 60  
gacaaccttg gtgctgaggt taactccatc cattgtagtg gcctgtatat caatgggacg 120  
attgcatatt tttcctgggt gagctttcca gaggtctgaa attttctccc cacccttagt 180  
ctgagatact ttatcatgat cganccactc cgtccactcc acgtnttgaa cccactcact 240  
ggacaaagaa acattgaaat attcgccatg ctctgtctgg aacaatttga ataccggggc 300  
agcagcagag cctcgatgnc caggatattc aatatggtct tccactgaag atgatggatt 360  
tcctttcaca gntagaaaac ttncnagggg gtctaaatcc aaggtgcagg aagngngngc 420

<210> 93  
<211> 241  
<212> DNA  
<213> Homo sapien  
  
<220>  
<221> misc\_feature  
<222> (1)...(241)  
<223> n = A,T,C or G

<400> 93  
accacgaatt ncaacatcca gatccaccac tatectaattg ggattgtaac tngaaactgt 60  
gcccggctcc tgaaagccga ccaccatgca accaaccggg tgggtgcacct catcgataag 120  
gtcatctcca ccatcaccaa caacatccag cagatcattg agatcganga cacctttgag 180  
acccttcggg ctgctgnggc tgcacacagg ctcaacacga tgcttgaagg naacggncag 240  
t 241

<210> 94  
<211> 395  
<212> DNA  
<213> Homo sapien  
  
<220>  
<221> misc\_feature  
<222> (1)...(395)  
<223> n = A,T,C or G

<400> 94  
actctattnt aattctgcct ttttatactt aattctaaat ttttcccctc taattttacaa 60  
caaattttgt gattttttata agaattctatg cctcccacat tctcagattc ttctcttttc 120  
tccttttatt ctttgcctta attcagtata agctttcttg gtatttttagg cttcatgcac 180  
attcttattc ctaaacacca gcagttcttc agagacctaa aatccagtat aggaataact 240  
gtgttagttc ttgaaaaagc attaaagaca tttttccctg aaacatacag aacatgtcat 300  
gccaaatctc ttgtttacat aataaactgg taataccggg gaattgcaca tacagatttt 360  
atctccaaga tagaataact taaatattaa aacgt 395

<210> 95  
<211> 304  
<212> DNA  
<213> Homo sapien  
  
<220>  
<221> misc\_feature  
<222> (1)...(304)  
<223> n = A,T,C or G

<400> 95  
cgaggtagag tgatngctcc cctggggcaa tacaatacaa gaacngnggg ttttgtcaaa 60  
ttggaacaag gaaacagaac cacagaaata aatacattgg ttaacatcag attagttcag 120

```

gttacttttt tgtaaaagtt aaagtacgag gggacttctg tattatgcta actcaagtan    180
actggaatct cctgttttct tttttttttt taaatnggtt ttaatttttt ttaattggat    240
ctatcttctt ccttaacatt tcagttggag tatgtagcat ttagcaccac tggctnaaac    300
ctgt                                           304

```

```

<210> 96
<211> 506
<212> DNA
<213> Homo sapien

```

```

<400> 96
acactgtcag cagggactgt aaacacagac aggggtcaaag tgttttctct gaacacattg    60
agttggaatc actgttttaga acacacacac ttactttttc tgggtctctac cactgctgat    120
atthttctta ggaaatatac ttttacaagt aacaaaaata aaaactctta taaattttcta    180
tttttatctg agttacagaa atgattactg aggaagatta cttagtaatt tgtttaaaaa    240
gtaataaaat tcaacaaaca ttgctgaat agctactata tgtcaagtgc tgtgcaaggt    300
attacactct gtaattgaat attattcctc aaaaaattgc acatagtaga acgctatctg    360
ggaagctatt tttttcagtt ttgatatttc tagcttatct acttccaaac taatttttat    420
ttttgctgag actaatctta atcattttct ctaatatggc aaccattata accttaattt    480
attattaacc ataccctaag aagtac                                           506

```

```

<210> 97
<211> 241
<212> DNA
<213> Homo sapien

```

```

<220>
<221> misc_feature
<222> (1)...(241)
<223> n = A,T,C or G

```

```

<400> 97
atthttcttt taattacttt agagagctag ggatgcaaag gttttcagtt agaaagcctt    60
tatttacttt tggaaattga acaagaaatg catctgtctt agaaactgga gattatttga    120
tgtaggttaa aacatgtaat tgnntctctg gcaaatttgt atcantnatt ngaaatgag    180
atattangaa aaaccaatc ttcttaaatc tagnncatct ttctttanaa gaacattana    240
t                                           241

```

```

<210> 98
<211> 79
<212> DNA
<213> Homo sapien

```

```

<220>
<221> misc_feature
<222> (1)...(79)
<223> n = A,T,C or G

```

```

<400> 98
ggcaaacana cttatgctgn ancnggggtt tancaaggtt ttcaaagnaa aaanccatt    60
ngactttatg gaaaatatt                                           79

```

```

<210> 99
<211> 316
<212> DNA
<213> Homo sapien

```

```

<220>

```

<221> misc\_feature  
 <222> (1)...(316)  
 <223> n = A,T,C or G

<400> 99  
 ccacatatgt aaaaccacaga aagaccngnt tngcactttc actgagagtt gagtcacatg 60  
 ggctgtcnac aggtgtctga cgtgtaaact tggaaatcaaa ctgacttaca tctctctcag 120  
 attgcaacag aggttttaaag gggggctcca cctttcgagc cagaagttct tcccagttaa 180  
 tgtgtctaaa gaatggatga gcttgaactt ctccagcgct cccaggacca gctcccagac 240  
 gagaagcagc atttcttttc agcagctttt taagcagatc tctggcttct tgngtgaggt 300  
 agggaggcaa attgag 316

<210> 100  
 <211> 425  
 <212> DNA  
 <213> Homo sapien

<220>  
 <221> misc\_feature  
 <222> (1)...(425)  
 <223> n = A,T,C or G

<400> 100  
 accgctttca gaaagtttat atgggttatt cttcagcctc tcttttatgc ctttcgacct 60  
 ctgtttatca accccaaacc aattacgtat ctggaagtta tcaataccgt ggcacaggctc 120  
 acttttgaca ttttaattta ttactttttg ggaattaaat ccttagtcta catgttggca 180  
 gcatctttac ttggcctggg ttgcaacca atttctggac attttatagc tgagcattac 240  
 atgttcttaa agggncatga aacttactca tattatgggc ctctgaattt acttaccttc 300  
 aatgtgggtt atcataatga acatcatgat ttccccaaca ttccctggaaa aagtcttcca 360  
 ctggtgagga aaatagcagc tgaatactat gacaacctgc ctcaactaca tttctggata 420  
 aaagg 425

<210> 101  
 <211> 156  
 <212> DNA  
 <213> Homo sapien

<220>  
 <221> misc\_feature  
 <222> (1)...(156)  
 <223> n = A,T,C or G

<400> 101  
 actgacttgg gaatgtcaaa attctttatt atgatcttcc gagtggtgtc ctgagctttg 60  
 ttggccctca actgcaggca gagaaccagg agcagggtgg cagggtctggc cctgaacagg 120  
 agctggagca agcgcatgct ngagaaaaca gaaggc 156

<210> 102  
 <211> 230  
 <212> DNA  
 <213> Homo sapien

<220>  
 <221> misc\_feature  
 <222> (1)...(230)  
 <223> n = A,T,C or G

<400> 102

```

actccaggcc gggnetcagg ttatcaaaag tgcaggagct ctgatcagca tggaccactt      60
cttccaaaga atttccctgc tggccgtttg taggggttggt ggttaattcta taaccagtaa      120
tgtctggggg ggtgctcctc tcccaggaga ctgtgagcac tccagtgtca gggtttgcct      180
ccagatgcaa gntngtnggt ggagacaatg gtgncaccac tttgtnnaca      230

```

```

<210> 103
<211> 404
<212> DNA
<213> Homo sapien

```

```

<220>
<221> misc_feature
<222> (1)...(404)
<223> n = A,T,C or G

```

```

<400> 103
actgtgaacc ctgnggnttc nangcgacct acctggagct ggccagtgtc gtgaaggagc      60
agtatccggg catcgagatc gagtccgccc tggggggcac aggtgccttt gagatagaga      120
taaattggaca gctggtgttc tccaagctgg agaattggggg ctttccctat gagaaagatc      180
tcattgaggc catccgaaga gccagtaatg gagaaaccct agaaaagatc accaacagcc      240
gtcctccctg cgtcatcctg tgactgcaca ggactctggg ttctctgctc gttctggggg      300
ccaaaccttg gtctcccttt ggtcctgctg ggagctcccc ctgcctcttt cccctactta      360
gtccttagc aaagagaccc tggcctccac tttgcccttt gggg      404

```

```

<210> 104
<211> 404
<212> DNA
<213> Homo sapien

```

```

<220>
<221> misc_feature
<222> (1)...(404)
<223> n = A,T,C or G

```

```

<400> 104
accaggttat ataatagtat aacactgeca aggagcggat tatctcatct tcctcctgta      60
attccagtgt ttgtcacgtg gttgttgaat aaatgaataa agaattgagaa aaccagaagc      120
tctgatacat aatcataatg ataattatct caatgcacaa ctacgggttg tgctgaacta      180
gaatctatat tttctgaaac tggctcctct aggatctact aatgatttaa atctaaaaga      240
tgaagttagt aaagcatcag aaaaaaaagt gggatttctc acaagtcagg acattctacg      300
tgactataat ataatctcac agaaatttaa cattaatacn ttctaagatt taattcttag      360
antctnggta aacaaagtag ctctgtgga natgattggc atca      404

```

```

<210> 105
<211> 325
<212> DNA
<213> Homo sapien

```

```

<220>
<221> misc_feature
<222> (1)...(325)
<223> n = A,T,C or G

```

```

<400> 105
acagcagaag ccagtctang atgggtgat tcaatttctg cctctagtat ttctttgtct      60
tgtttttcct tcaatttaga agtgagcatt gtgttctcag ctatcagaac tttaagctgc      120
ccactatatt gagatgccct ttagctaat gattcctctt tcagttttag ggtcatctga      180
agttcagcat tcttttcttt taaaatctta atgtcctcaa agtatttatt ttccttttcc      240

```

tggtattggn gtttcagngt ggctatttcc agtttttagca tggcaattnc ctttttcaac 300  
atgcaatttt catgtaagag ataata 325

<210> 106  
<211> 444  
<212> DNA  
<213> Homo sapien

<220>  
<221> misc\_feature  
<222> (1)...(444)  
<223> n = A,T,C or G

<400> 106  
actgtcttca atnctatgcg tgcaggtgtc taccacaggc aaacagtttt ctccccattt 60  
tgtagtaatg tgatttttct attagcaaaa agaggtcacc agcccctgta gacttaaggg 120  
actcaagtca caggatgggg atttcctctt aatatttttt atttngttgt ttgaactctt 180  
gatgcaacat tgtagagcag ggtgttcagg acctgctgtg cccaaggagc tgataaagga 240  
aaaagctcta ttatttcttt ttgtgatttg atgcacagat gaaaaactta acacacaata 300  
acagaagttg gncgttaata aatcacatcc taggctttca gcgcttnogt aagcagacga 360  
catcttcagt ttcttagctc ttgnagnttc aacacngnaa catcaatgat gcatatgtnc 420  
agaatcagtt acaaagacca tccg 444

<210> 107  
<211> 287  
<212> DNA  
<213> Homo sapien

<220>  
<221> misc\_feature  
<222> (1)...(287)  
<223> n = A,T,C or G

<400> 107  
acctgcactc gnacntcagg cantaggect ccacgtcatg gccaggcact ggcatgggct 60  
ccaccacgtg caggcagttg cagtccttct gggatacatt ctggttgtaa atgtgcccac 120  
tgatgtttct ataagtgagg acagatgcat ttgcaccgga tatcttcana actcttgttg 180  
gctncagctg ggggcaccaa caaacacccg accacagcca ccaaagataa nagcttcatg 240  
cttatcangc ttgctgggoc agnaaagccg gacacctaca agccnc 287

<210> 108  
<211> 478  
<212> DNA  
<213> Homo sapien

<400> 108  
acatgtgcaa gaatttgaa aagcagggca ttttccctca tctctcctag agggaaatc 60  
acagcatctg tctctactgg tccacactgg actgcagaca atgtcaaaac tctggatttg 120  
gaatgaggct gatttccctt cccctttaag gagttttcca agaatttcat aaccatcagt 180  
tgttatattt ccagcttcc tcatagcagt caatgtaaat 240  
cttaacactt tttgaggtca ctacaatatg aacctgtga aaacttccat aaaataatgt 300  
ctttacttct tctgtgtcaa atgtaacagt ttgcacctcg cctcttgat ccttgtaaaa 360  
gaatgataac gtcttgctag aaggatctgc aatcactoca acttggtggt tgtagtctct 420  
gtctgtgatt tgccaaattg caaaagggc actgggaggt tctgggagaa gtctgaat 478

<210> 109  
<211> 361  
<212> DNA

<213> Homo sapien

<220>

<221> misc\_feature

<222> (1)...(361)

<223> n = A,T,C or G

<400> 109

gaatttttct tctanaataa gtattctgtt gacacagact attggttaaga ttttcaacat	60
aaggtaatgc taggactggc ctcttagcat gagttgtgag taaagatctg gtctgttggt	120
tctccaaaag aagnttctta ctgcttgctc ctcatgagtt ttctgtttct gctttctctt	180
tttcatattg atatatacgg ntttttaaag ggtnattgta attaaatatc tcctcatttt	240
tctcttttag gagatgatgt tgcattttcc tctcaagaaa atgaatatca attgttatct	300
tgcttttgnt gncagcttcc ttatgtgcat gaactaattg ctgttgaagc cacatatttt	360
t	361

<210> 110

<211> 305

<212> DNA

<213> Homo sapien

<220>

<221> misc\_feature

<222> (1)...(305)

<223> n = A,T,C or G

<400> 110

acataatgac tnncanagtg aagctgattg gctgcggttc tggagtaaata ataagctctc	60
cgttccctggg aatccgcact acttgagtca cgtgcctggc ctaccaaata cttgccaaaa	120
ctatgtgcct tatccaccc tnnaatctgn ctctcatttt ntcagctgtt ggatcagaca	180
atgacattcc tntagatntg gcgatcaagc attccanacc tgngccaaact gcaaacgggtg	240
cctncaagga gaaaacgaag gcncacacaa atgnaaaaaa tgaangnccc ttgaatgtac	300
taaaa	305

<210> 111

<211> 371

<212> DNA

<213> Homo sapien

<220>

<221> misc\_feature

<222> (1)...(371)

<223> n = A,T,C or G

<400> 111

cgggggcccag ccgggggtat tcagccatcg atcaaaactca aaacctggaa tgatatccac	60
tctctttttc ttaagctcag ggaaatatcc caagtagaag tccagaaaagt catcggctaa	120
gatgcttcgg aatttgaatt catgcacata ggccttgaga aaactgtcaa actgatcctg	180
atcacccacc aagtgggcca ggtatgagac aaagcagaaa cttttctcgt aggggggtctc	240
attatagggtg tcgtccgggt caacgcctgg ttcaatcttc acgcggagct tgttgagtg	300
gttttctctc ccagtgatgt ccatgtgctg acgcagcaga ncccgccccg ttgcagcctc	360
caagcagng t	371

<210> 112

<211> 460

<212> DNA

<213> Homo sapien



<220>  
 <221> misc\_feature  
 <222> (1)...(460)  
 <223> n = A,T,C or G

<400> 112  
 acatcttagg ttttnttcc tttantgtga agaggcggtt ccaccaaccc acagctctgc 60  
 gtcgagtttt tactagattg ctgcaaattt catggaatct ttgctgttgt tcagtgggtcc 120  
 atttattgga gccaaaaatt ctagggcgct agaattggaa caaggtagtc agccaagcac 180  
 aaaaacataa caaaacagga aacgccggac agaacagatg gatctagata gtagataatc 240  
 agaaacacca aagaaaccac acccatgatg gcagggtgaa accaggctct ttctcatcgg 300  
 aggactttat cagccatcag catcaattct ccccatcctt gcagctgttc ttccagactt 360  
 gcagtctctg cagccagcag gttgggtgct gcgattacct ccctccgcca tcgtctcggg 420  
 gatgcagtct ctacaagcgc aggccacctc cccaacgagt 460

<210> 113  
 <211> 204  
 <212> DNA  
 <213> Homo sapien

<400> 113  
 gagaagacag cagagctgct ttccgcctct ttgagaccaa gatcacccaa gtccctgcact 60  
 tcaccaagga tgtcaaggcc gctgctaatac agatgcgcaa cttcctgggt cgagcctcct 120  
 gccgccttag cttggaacct gggaaagaat atttgatcat ggggtctagat ggggccacct 180  
 atgacctcga gggacacccc cagt 204

<210> 114  
 <211> 137  
 <212> DNA  
 <213> Homo sapien

<220>  
 <221> misc\_feature  
 <222> (1)...(137)  
 <223> n = A,T,C or G

<400> 114  
 accgcaagaa atgggacagc aacgtcattg agacttttga catcgncgc tngacagtca 60  
 acgctgacgt gggctattac tcttgagggt gtcccaagcc cctgaagaac cgtgatgtca 120  
 tcaccctcgc ntccctg 137

<210> 115  
 <211> 278  
 <212> DNA  
 <213> Homo sapien

<220>  
 <221> misc\_feature  
 <222> (1)...(278)  
 <223> n = A,T,C or G

<400> 115  
 gcggggcggt ttntggactc gctcatttac agagcatgcg tgggtcttcac ccttggcatg 60  
 ttctccgcgc gcctctcgga cctcaggcac atgcgaatga cccggagtgt ggacaacgtc 120  
 cagntcctgc cctttctcac cacggangtc aacaacctgg gctggctgan ttatggggct 180  
 ttgaaggag acgggatcct catcgtcanc aacacagtgg gtgctgcgct tcanacctg 240  
 tatatctttg gcatactgc attactgcc tcggaage 278

<210> 116  
<211> 178  
<212> DNA  
<213> Homo sapien

<220>  
<221> misc\_feature  
<222> (1)...(178)  
<223> n = A,T,C or G

<400> 116  
acacgcgtcat angtcaaaaag tncagtgcgtg gccatcttgc atcaaatggt cttaaggcag 60  
tgactggcta tcaaccacag nttctgtctc cccagntgca aacacaggat ccatgcaaca 120  
gttctgagac catacactta gaaaccacng ggagatgcgg atcanatgca naactnnc 178

<210> 117  
<211> 360  
<212> DNA  
<213> Homo sapien

<220>  
<221> misc\_feature  
<222> (1)...(360)  
<223> n = A,T,C or G

<400> 117  
actccccaat ggnggattta ttactattaa agaaaccagg gaaaatatta attttaatat 60  
tataacaacc tgaaaataat ggaaaagagg tttttgaatt ttttttttaa ataaacacct 120  
tcttaagtgc atgagatggg ttgatgggtt gctgcattaa aggtatttgg gcaaacaata 180  
ttggagggca agtgactgca gttttgagaa tcagttttga ccttgatgat tttttgtttc 240  
cactgtggaa ataatgttt gtaaataagt gtaataaaaa tccctttgca ttctttctgg 300  
accttaaatg gtagaggaaa aggctcgtga gccatttgtt tcttttgctg gttatagttg 360

<210> 118  
<211> 125  
<212> DNA  
<213> Homo sapien

<220>  
<221> misc\_feature  
<222> (1)...(125)  
<223> n = A,T,C or G

<400> 118  
gcgtcgtgct atgaccggac ttngtcttga aaggggatga cagcatggga ggcaatggnt 60  
ncacatgtaa accccacact gaaagacaag gcactctctc cacagcagcc ccaacaacta 120  
gcctt 125

<210> 119  
<211> 490  
<212> DNA  
<213> Homo sapien

<220>  
<221> misc\_feature  
<222> (1)...(490)  
<223> n = A,T,C or G

<400> 119  
nacaaagaaa agcaaaaaga atttacgaag attgtgatct cttattaaat caattgttac 60  
tgatcatgaa tgtagttag aaaatgtag gttttaactt aaanaaaatn gtattngat 120  
tttcaatntt atgttgaaat cngngtaata tcctgangtt nttttcccc cagaagataa 180  
agaggataga caacctctta aaatattttt acaatttaat ganaaaaaagn ttaaaattct 240  
caatacnaat caaacaattt aaatatttta agaaaaaagg aaaagtagat agtgatactg 300  
agggtaaaaa aaaattgatt caattttatg gtaaaggaaa cccatgcaat ttacacctaga 360  
cagccttaaa tatgtctggt ttccatctg ctacgatttc agacatttta tggtcctctt 420  
actcaattga taccaacaga aatatcaact tctggagtct attanatgtg ttgtcacctt 480  
tctnaagctt 490

<210> 120  
<211> 361  
<212> DNA  
<213> Homo sapien

<220>  
<221> misc\_feature  
<222> (1)...(361)  
<223> n = A,T,C or G

<400> 120  
caggtagagt aaaattaaca ctcccggttac aggaaatgta tgacgcaaat aatataaaat 60  
taaaagggtga aaaaaagggtg acactggttt cctaagatac aatttactct ttacaaccag 120  
gggtccacagg tccaggctgc anagcgggca tcaggaagca gagcctncca cctgcttctg 180  
ggggacctgg taataaaaaat cagcccatga tggcgctatg gcctctcaga caccacacgc 240  
tgcctaaaca cctagagctc tggaaatagt caacaggaga gtgatttcca tgggggaaat 300  
tttaanaaag atgcacatgg gacaggcaat agaaagttag ccaaggntaa atttggtacc 360  
t 361

<210> 121  
<211> 405  
<212> DNA  
<213> Homo sapien

<220>  
<221> misc\_feature  
<222> (1)...(405)  
<223> n = A,T,C or G

<400> 121  
acacaaaacc ttttnacata ttgggggctt accgctccaa attgctactg atcctttaag 60  
ttcacaaat agaatctt caccaattaa gtaataaccc tcattacaaa taaagtgcac 120  
ctgataacca aactcgtaag tccatttgc agggactgct tggccattta aaggatcccg 180  
tatatatgga catgtttctc tataacaggc gtcactctgag acaggtagcc atgtatgatt 240  
ccgatcacia atagtatggg tggcaagagg aggtatatag aagtatcctt ttttacactt 300  
ataatctact cgttcaccaa tctcatagta ggggtttggt ttaccaatga gctccatan 360  
cttcaaatgt tgggtggctn ctacaggca tcnggcanaa ngagt 405

<210> 122  
<211> 152  
<212> DNA  
<213> Homo sapien

<220>  
<221> misc\_feature  
<222> (1)...(152)  
<223> n = A,T,C or G

<400> 122  
accccgtcc gttgncacag atcgtgtctt gccactcca tcggccattc acttggcagg 60  
tgcgattggc agagccccgg agagtgtaac cgtcatagca gtggaaagag atctcatcac 120  
tcacattgta gtagggagac cggggccaan ta 152

<210> 123  
<211> 336  
<212> DNA  
<213> Homo sapien

<400> 123  
acatctgaca tatttatata gcacataaat tagggagtgc tctgaccctt gcccgaggag 60  
cccaagcact gagcagggag gtgaacgcca gtccagaaaag aagggtctgg agcccctgct 120  
ctgtctcttc catcacgggg ctccccctagg gcctccccag gcctccttgg ctcagtccag 180  
gtgtctgcag gaggaagggt ttgtctgcat ttagtgtctg agactgggtt tgaggaggca 240  
ccagataaaa ggagatacac ttgcagctat aaagttagct tcaaaccoca gggcttgtaa 300  
ttccaagagg aggggtgggga ggcgaggcca tagtct 336

<210> 124  
<211> 253  
<212> DNA  
<213> Homo sapien

<220>  
<221> misc\_feature  
<222> (1)...(253)  
<223> n = A,T,C or G

<400> 124  
ctgcaagagc ccagatcacc cattccgggt tcaactcccc cctccccaag tcagcagtc 60  
tagcccaaaa ccagcccaga gcaggggtctc tctaaagggg acttgagggc ctgagcagga 120  
aagactggcc ctctagcttc taccctttgt ccctgtagcc tatacagttt agaataattta 180  
tttgtaatt ttattaaaaa gctttaaaaa aacaaaaaaa aaaaaaaaaa aaaaaaaaaa 240  
aaaaaagntt gtn 253

<210> 125  
<211> 522  
<212> DNA  
<213> Homo sapien

<400> 125  
acaactgcaa gtctaagata atgttcattc attcccatca taaatgtaac attctaata 60  
ggtgtcttct gatgtcatct gtcagaattt cttttaaaact ttttcttcat cttcaacatt 120  
atcaaagttc atccttattc ctcttgctt gatttcggag agtttccaat ttttcaactta 180  
ttaaggcagc gattgtcttt gcatctctgg tatttatctg ctcttcttga aaatttctct 240  
ttgtcttttc gtagaaataa aacttaacag ttggatagga cctgatccca gctttctggc 300  
atgtctgagc ataagcctga cagtctactt ttccagcttt cacttttctt ttaatcatcc 360  
tagccaagag ctcaaattct ggagcaaaat tetggcaagg tccacaccaa ggagcataga 420  
aatcaatcac ccaatgatth ttcccttgta gaacttttct actgaaagtc tgagggtgta 480  
gatctgtgga tacttgaggt aaaaatccta gacccagat tc 522

<210> 126  
<211> 374  
<212> DNA  
<213> Homo sapien

<220>

<221> misc\_feature  
 <222> (1)...(374)  
 <223> n = A,T,C or G

<400> 126  
 tttttaagat attaacttta cctttataaa tctttgtgtg aaatgaaaaa aaaaatcaag 60  
 gcatacaaat ttcattgtgt tctacatttt taaataccat cctttgtctc cgttaaaaga 120  
 ttttcatcca tttattcaaa aaccttttaa gttcaactgt ccaatttaag acagagtga 180  
 gacatttttg agtatctgaa ctaagcattg tcttgactga aacgaagtaa gaactcaatg 240  
 agagtccctg tgggcctccc aggcattgct tcccgtagat agggaaacttc atctttgttg 300  
 gncatcacgc ctgctatgtc taaatgtgcc cacttaggat gagttacgaa ttctttcagg 360  
 aatgctgcag ctgt 374

<210> 127  
 <211> 130  
 <212> DNA  
 <213> Homo sapien

<220>  
 <221> misc\_feature  
 <222> (1)...(130)  
 <223> n = A,T,C or G

<400> 127  
 aaagccaaga cngccattgg cactgctatg gtaaggncac agggcancca gggccttctg 60  
 gcaaaaggng atacnaccag cactatnaac agacaggaca tggttgagag gnagnctaca 120  
 caantcctaa 130

<210> 128  
 <211> 350  
 <212> DNA  
 <213> Homo sapien

<220>  
 <221> misc\_feature  
 <222> (1)...(350)  
 <223> n = A,T,C or G

<400> 128  
 aacttgattt cegntnaaaa gaancatcat ctttaccttg acttttcagg gaattactga 60  
 actttcttct cagaagatag ggcacagcca ttgccttggc ctcacttgaa gggctctgat 120  
 ttgggtcctc tggctctctg ccaagnttcc cagccactcg agggagaaat atcgggaggt 180  
 ttgacttcct ccggggcttt ccgagggtct tcaccgtgag ccctgcggcc ctcagggtcg 240  
 caatcctgga ttcaatgtct gaaacctcgc tctctgcctg ctggacttct gaggccgtca 300  
 ctgccactct gtctccagc tctgacagct cctcatctgt ggctgttga 350

<210> 129  
 <211> 505  
 <212> DNA  
 <213> Homo sapien

<220>  
 <221> misc\_feature  
 <222> (1)...(505)  
 <223> n = A,T,C or G

<400> 129  
 acaataccaa agcttcataa tgctaaagaa aacccaaaca aaagacaatg gtttacacag 60

```

ggaataacc ctaaggcaat atgaaaacag tcataattta ttactgataa agagtaaagg      120
catccttccc atagaggggg ggaattcaca gggaacacta attatatcag atgaaccacg      180
gggatagaaa ataggcccat ttttaaaatt cattgagaaa ttattacttt ttctccacaa      240
ctgtgattct atacaaaata taaaccttgc aaaccttatg tgctacctga cagataaaaag      300
tagcaggagc cagactcttg aagcacttga gactgatttc tacaaagtcc aggaagagca      360
atgattccag tgtgcagtgc tgatgcatgt gtgagcctaa catgttattc agctctgggt      420
gcagcccat ctacatgggg cccagttagt ttttagggag tcacagatta ngcaggcaac      480
cgaggggcat gatttaaaaa gcaca                                           505

```

&lt;210&gt; 130

&lt;211&gt; 526

&lt;212&gt; DNA

&lt;213&gt; Homo sapien

&lt;400&gt; 130

```

acaaaagagc ctgattcttt ttaattccac aaatacctag catctcaaag taacatgtaa      60
acaaacttct atgctgctca atgaatcctt ccaatttcga taataaacta aatagtattg      120
gatctagtat atgactttca tgtgtaagtt atggttctat ccattacttt aacaatatta      180
ctgatgtaac agagaaaaat tttcaactat tgtacttatt taaaacaaac tgacaagtgc      240
aagcacctgt cttcagaaaa gccagcagca tttttttttt ttttaacatac tcaaagtaag      300
atgtggccta agcccttaat acctttctga acagccatgc aactaaacac cctcaggaga      360
tgttacataa gggagagaag aacatggagc aatttgact tttccccta gataatatta      420
acaaggtaaa gcaaatccag atctttatga atgaatggct gtcattgtta atacacttgg      480
agctctataa aactagagcc actatcatat atgtttatat agatat                                           526

```

&lt;210&gt; 131

&lt;211&gt; 477

&lt;212&gt; DNA

&lt;213&gt; Homo sapien

&lt;400&gt; 131

```

ctcagttttc ccagcaacag atgctcctga gcaatttatt agtcaagtga cgggtgctgaa      60
atacttttct cattacatgg aggagaacct catggatggg ggagatctgc ctagtgttac      120
tgatattcga agacctcggc tctacctcct tcagtggcta aaatctgata aggccctaatt      180
gatgctcttt aatgatggca cctttcaggt gaatttctac catgatcata caaaaatcat      240
catctgtagc caaaatgaag aataccttct cacctacatc aatgaggata ggatatctac      300
aactttcagg ctgacaactc tgctgatgtc tggctgttca tcagaattaa aaaattgaat      360
ggaatatgcc ctgaacatgc tcttacaaag atgtaactga aagacttttc gaatggaccc      420
tatgggactc ctcttttcca ctgtgagatc tacaggggaa ccaaaagaat gatctag      477

```

&lt;210&gt; 132

&lt;211&gt; 404

&lt;212&gt; DNA

&lt;213&gt; Homo sapien

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (1)...(404)

&lt;223&gt; n = A,T,C or G

&lt;400&gt; 132

```

accacacgan cgggnatcnt ttgnacatag tgagaccggg ctgattccca tacatgaatc      60
cattcatgga gtgcatttta ttagatncct gaaagtcttc atcttcctta tccacctgat      120
caggngcagt tgtaaacatn cctaataatta tcttcaggga gtaaactctc attctcatca      180
aatactgtag gaaacaaata gaattccttg tctacatctt tctgtctccc atttgcatat      240
aaacttctct tcttgcatat tttcattggc ccaataagcc cagtgaatat atcttttagtg      300
ggatccacag cagaataata catcttagct agacacacag ggatctgcat tacnggggtc      360
ctacttcttt ggggacagcc cttcatacgn gaatgtttnt gtgg                                           404

```

<210> 133  
 <211> 552  
 <212> DNA  
 <213> Homo sapien  
  
 <220>  
 <221> misc\_feature  
 <222> (1)...(552)  
 <223> n = A,T,C or G

<400> 133  
 accccaaatt atctctctcc tgaagtcctc aacaaacaag gacatggctg tgaatcagac 60  
 atttgggccc tgggctgtgt aatgtataca atgttactag ggaggccccc atttgaaact 120  
 acaaatctca aagaaactta taggtgcata agggaaagcaa ggtatacaat gccgtcctca 180  
 ttgctggctc ctgccaaagca cttaattgct agtatgttgt ccaaaaaccc agaggatcgt 240  
 ccagtttgg atgacatcat tcgacatgac ttttttttgc agggcttcac tccggacaga 300  
 ctgtcttcta gctgttgtca tacagttcca gatttccact tatcaagccc agctaagaat 360  
 ttctttaaga aagcagctgc tgctcttttt ggtggcaaaa aagacaaagc aagatatatt 420  
 gacacacata atagagtgtc taaagaagat gaagacatct acaagcttag gcatgatttg 480  
 aaaaagactt caataactca gcaacccagc aaacacaggg acagatgang agctccacca 540  
 cctaccacca ca 552

<210> 134  
 <211> 496  
 <212> DNA  
 <213> Homo sapien

<400> 134  
 acattgatgg gctggagagc aggggtggcag cctgttctgc acagaaccaa gaattacaga 60  
 aaaaagtcca ggagctggag aggcacaaca tctccttggg agctcagctc cgccagctgc 120  
 agacgcta at tgctcaaact tccaacaaag ctgcccagac cagcacttgt gttttgattc 180  
 ttcttttttc cctggctctc atcctcctgc ccagcttcag tccattccag agtcgaccag 240  
 aagctgggct tgaggattac cagcctcacg gagtgaactc cagaaatc ctgaccacca 300  
 aggaagtaac agaaaatctg gagacccaag tggtagagtc cagactgacg gagccacctg 360  
 gagccaagga tgcaaatggc tcaacaagga cactgcttga gaagatggga ggggaagccaa 420  
 gacccagtg ggcgcatccg tccgtgctgc atgcagatga gatgtgagct ggaacagacc 480  
 ttttctgggc cacttt 496

<210> 135  
 <211> 560  
 <212> DNA  
 <213> Homo sapien

<400> 135  
 actgggagtg atcactaaca ccatagtaat gtctaataat cacaggcaga tctgcttggg 60  
 gaagctagtt atgtgaaaag caaatagagt catcacagtag ctcaaaaaggc aaccataatt 120  
 ctctttggg caggtcttgg gagcgtgac tagattacac tgcaccattc ccaagttaat 180  
 cccctgaaaa ctactctca actggagcaa atgaactttg gtcccaata tccatctttt 240  
 cagtagcgtt aattatgctc tgtttccaac tgcatttccct ttccaattga attaaagtgt 300  
 ggctcgttt ttagtcattt aaaattgttt tctaagtaat tgctgcctct attatggcac 360  
 ttcaattttg cactgtcttt tgagattcaa gaaaaatttc tattcttttt ttgcatcca 420  
 attgtgcctg aactttttaa atatgtaa at gctgccatgt tccaaaccca tegtcaagt 480  
 tgtgtgttta gagctgtgca ccctagaaac aacatattgc ccatgagcag gtgcctgaac 540  
 acagaccctt ttgcattcac 560

<210> 136  
 <211> 424

<212> DNA  
 <213> Homo sapien

<220>  
 <221> misc\_feature  
 <222> (1)...(424)  
 <223> n = A,T,C or G

<400> 136  
 accagcaaat ctccattagc atttctcagg ttctcatgac cttttcagat atgttggttg 60  
 attttatgta tatattgctt agaaacaaaa atccacctga tattaataca aaccacaaaa 120  
 aatcataaaa gcaagcaaat gaacacaaaa ccctagtgtt gttgtgcttt tctttcacat 180  
 ttctacacag gagatttcta tatctcagat actttcaaaa tctaataagg aagtaaaatt 240  
 agtgcccttaa ccaaacagta agatacacia gaatcctcca tcacaagtta ctgaatcaaa 300  
 cttctcatga catttgccgt atattcagat ttgaagattt tttaaattta gaatttaaaa 360  
 caaactttag actgctgatt ttccatattt caaagactgt agctgtntgc agcatataaa 420  
 tgga 424

<210> 137  
 <211> 392  
 <212> DNA  
 <213> Homo sapien

<220>  
 <221> misc\_feature  
 <222> (1)...(392)  
 <223> n = A,T,C or G

<400> 137  
 tgcgggntg aaggctagca aaccgagcga tcatgtcgca caaacaaatt tactattcgg 60  
 acaaaatcga cgacgaggag tttagatcgc gacatgtcat gctgcccag gacatagcca 120  
 agctgggccc taaaacccat ctgatgtctg aatctgaatg gaggaatcct ggcatcagc 180  
 anagtcaggg atgggtccat tatatgatcc atgaaccaga acctcacatc ttgctgttcc 240  
 ggcgcccact acccaagaaa ccaaagaaat gaagctggca agctactttt cancccaag 300  
 ctttacacag ctgnccttac ttctaacat ctttctgata acattattat gctgccttcc 360  
 tgttctcact ctganatnta aaagatgttc aa 392

<210> 138  
 <211> 284  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <222> (1)...(284)  
 <223> n = A,T,C or G

<400> 138  
 tgccctgtgca cctctttgct tgaaatatgg caagacttgg aaaaatgttt gcccttagaa 60  
 tctatctcac tacttttagtt agttgtctcc ttggggcctg ggcacagttc tggccctgat 120  
 ctggaacaga ctcccttttc taaaactgaa cttgaccaca tcaaaagntt gnaaaacaat 180  
 ctccatggta attaaacttg cattcaacac catatggnaa cagaagatgg caggaggata 240  
 anatncagat cttatgatct ttccangnan ggcattgtac atga 284

<210> 139  
 <211> 249  
 <212> DNA



<213> Homo sapiens

<220>

<221> misc\_feature

<222> (1)...(249)

<223> n = A,T,C or G

<400> 139

```
gaggaagggg ggactgaatc tancacntg acngaactag agacagccat gggcatgac 60
atagacnct ttacccgata ntcgggcagc gagggcagca cgcagaccct gaccaagggg 120
gagctcaagg ggctgatgga gaaggagcta ccagggttcc ngcagagngg aaaaanacaag 180
gangccgtgg ataaattgct caaggaccta gacgccnatg gaggatgcc aggtggactc 240
cagcgagnt 249
```

<210> 140

<211> 390

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<222> (1)...(390)

<223> n = A,T,C or G

<400> 140

```
tcataatggg tggggcagct ataatnnact acaanaatca natgtttcac atctagacct 60
cgggcagcaa cagaggtagc cacaagaagt ttgcangtcc cattcttaaa gtcatttatg 120
atgctatctc tgtcatattg atcaatgcct ccatgaagag acatgcaagg ataagatgct 180
ctcattaaat ccttaagaag accatcagca tgttcctgct tatccacaaa tataatgaca 240
gatcctgact cttgataatg gcctagaagc tcaagtaact tcaagaattt cttttcttct 300
tcaatcacia tcacttgtn gctccacatct gagcaaacca cactcctgcc tccaacttgt 360
acctgccccg ggcgggcgct caagggcgaa 390
```

<210> 141

<211> 420

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<222> (1)...(420)

<223> n = A,T,C or G

<400> 141

```
gacactcagg gaaaagcatn ngncaaaanag agcttaaaat gcatcgccaa cggggtcacc 60
tccaaggtct tcctcgccat tcggaggtgc tccactttcc aaaggatgat tgctgaggtg 120
caggaagagt gctacagcaa gctgaatgtg cgcancatcg ccaagcggaa ccngaagcc 180
atcactgagg tcgtgcagct gcccaatcac ttctccaaca natactataa cagacttggn 240
cgaagcctgc tggaatgnga tgaanacaca gggcagcaca atcaggagac agcctgatgg 300
anaaaantgg gcctancatg gccaggcctc ttccacatcc tngcangaca gaccactgtg 360
cccaaacaca ccnctgagc tgacttnnac aggagacgca cnaaggagcc cggcagangc 420
```

<210> 142

<211> 371

<212> DNA

<213> Homo sapiens

<400> 142

```
gggttcgaca atgctgatcc gcaattagaa gacactggta agctgtgtta cactgggctt 60
cattgaaatc ttcaaggata tagccagctc ctgctcgaag ctgggattct gtatactgct 120
tggtgaaagg aggaatttcc aaaaattcct cctcttcttc actgcttcct gtaggacat 180
ctggcagttt ggagcggctg gccaaactgt cactggtgtt ggccatggta aggagaaatg 240
cgtagcccag aaacaaggtc ttgttgagag gcaaaggccc tctctgctct tccagggcag 300
agggttcacc ggtgttgtct ccactctcac aggggctcac aaactctcct gccctactt 360
gcaccagggt t 371
```

<210> 143  
<211> 270  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(270)  
<223> n = A,T,C or G

```
<400> 143
gggtggctgtg atnacctttn ttagtttaca aataaaaaag ntaaaaagaa atactgtgtt 60
tagggtaagg taacannttc atctaatacag aggagagtga agangaggcn ctgccttcta 120
ggngctgtga ccttctcctt ttcgngattc ttcnccacct tgggnaacat cttccccgct 180
atgctggaan tacttcggng ttctgcggtg gccatgntga acatctgatg aactgaaant 240
ncatccnaat gcacacgaag anatagncna 270
```

<210> 144  
<211> 259  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(259)  
<223> n = A,T,C or G

```
<400> 144
ttctctttgc tttttataat tttaaagnaa ataacacatt taactgtatt taagtctgtg 60
caaataatcc ttcagaagaa atatccaaga ttctgtttgc agaggtcatt ttgtctctca 120
aagatgatta aatgagtttg tcttcagata aagtgtcctt gtccagnaga actcaaaagg 180
ccttcaagct gttagtaag tgtaggttca gataagactc cgncatacga attccagctt 240
cccgtgcccc ctgtacctc 259
```

<210> 145  
<211> 433  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(433)  
<223> n = A,T,C or G

```
<400> 145
accacatnta ccatagtgtg attagtttta attttcacat gaatcaaagg ttctctttca 60
tgtctattta cagtccaatt gtgccaaact cttacttgtg tgetgactaa caaggcattt 120
agggtgtcag catcctagag tgctccaggg cagtgtcagc gttctcggga gtaaaagggtg 180
ccacttggtg gcaatgatat tccagaatta aatggggttt tgttgccatg gagactgcat 240
ttatataaat gtagcctgta gcttaagtta actaaaccta atgctgctgt taaaaacagt 300
```

```

ttattttaat attaaaaatac agttgattag caacagcggg gctgtatttt aagagacact 360
ttatttgaag tgcaatcata gttatttggt ttacaaattt tacagngcat tctaattact 420
gatgggtgca att 433

```

```

<210> 146
<211> 576
<212> DNA
<213> Homo sapiens

```

```

<400> 146
acctcaggcc tgtgcacctc tttgcttgaa atatggcaag acttggaaaa atgtttgccc 60
ttagaatcta tctcactact ttagttagtt gtctcctttg ggcctgggca cagttctggc 120
cctgatctgg aacagactcc cttttctaaa actggacctt gaccacatca aaagtttgta 180
aaacaatctc catggtaatt aaacttgcac tcaacacccat atggtaacag aagatggcaa 240
aggataagat tcagatctta gatctttcca agtagggcat gttagatgat agaaggatta 300
gttgcaagct ggatctgagc tcaggcttgg gcatgaagga aactgtctcc catgtgggtt 360
ggaagagtta ggggctccct gagctctatt gtgaactata cgggtttcat ccaaggaatg 420
gtatgatgtg ggcataaaaac cattcttcag acaactgaag atgggtccct tctgtagcca 480
gaaacactag ctgtctgca ttgccatttc ctttacccca ggcggcctgc agaaggaaa 540
gccataatta attaaaaggc ttaatgaagt tttgga 576

```

```

<210> 147
<211> 300
<212> DNA
<213> Homo sapiens

```

```

<400> 147
ccagccccc ggaggaaggt gggctctgaat ctaccacat gacggaacta gagacagcca 60
tgggcatgat catagacgtc ttaccctgat attcgggcag cgagggcagc acgcagaccc 120
tgaccaaggg ggagctcaag gtgcttatgg agaaaggagc taccaggctt ctgcagagt 180
gaaaagacaa ggatgccgtg gataaattgc tcaaggacct agacgccaat ggagatgcc 240
agggtggaatt cagtgaagtc atcgtgttcg tggctgcaat cacgtctgcc tgtcacaagt 300

```

```

<210> 148
<211> 371
<212> DNA
<213> Homo sapiens

```

```

<400> 148
acataatcct cataatgggt ggggcagcta taatttacta caagaatcag atgtttcaca 60
tctagacctc gggcagcaac agaggtagcc acaagaagtt tgcaggctcc attcttaaag 120
tcattttatga tgctatctct gtcataattga tcaaatggcc tccatgaaga gacatgcaag 180
gataagatgc tctcattaaa tccttaagaa gaccatcagc atgttctctgc ttatccacaa 240
atataatgac agatcctgac tcttgataat ggcctagaag ctcaagtaac ttcaagaatt 300
tctttttctt ttcaatcaca atcacttggt gctccacatc tgagcaaaacc acactcctgc 360
ctccaacttg t 371

```

```

<210> 149
<211> 585
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<222> (1)...(585)
<223> n=A,T,C or G

```

&lt;400&gt; 149

```
cgagggtacan cactgctaaa tttgacactn anggaaaagc attcgtcaaa gagagcttaa 60
aatgcatcgc caacgggggc acctccaagg tcttctcgc cattcggagg tgctccactt 120
tccaaaggat gattgctgag gtgcaggaag agtgctacag caagctgaat gtgtgcagca 180
tcgccaagcg gaaccttgaa gccatcactg aggtcgtcca gctgccaat cacttctcca 240
acagatacta taacagactt gtccgaagcc tgctggaatg tgatgaagac acagtcagca 300
caatcagaga cagcctgatg gagaaaattg ggcctaacat ggccagcctc ttccacatcc 360
tgcagacaga ccactgtgcc caaacacacc cacgagctga cttcaacagg agacgcacca 420
atgagccgca gaagctgaaa gtctctctca ggaacctccg aggtgaggag gactctccct 480
cccacatcaa acgcacatcc catgagagtg cataaccagg gagaggntat tcacaacctc 540
ccaaactagt atcatttttag ggggngttga cacaccagtt ttgag 585
```

&lt;210&gt; 150

&lt;211&gt; 642

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (1) ... (642)

&lt;223&gt; n=A,T,C or G

&lt;400&gt; 150

```
acttncgggt tcgacaatgc tgatccgcaa ttagaagaca ctggttaagct gtgttacact 60
gggcttcatt gaaatcttca aggatatagc cagctcctgc tcgaagctgg gattctgtat 120
actgcttggt gaaaggagga atttccaaaa attcctctc ttcttctactg cttctctgtat 180
gaccatctgg cagtttggag cggctggcca acttgctact ggttggtggc atggttaagga 240
gaaatgcgta gcccagaaac aaggtcttgt tgagaggcaa aggccctctc tgctcttcca 300
gggcagaggg ttcaccggtg ttgtctccac tctcacaggg gctcacaaac tctctctgcc 360
ctactgcacc aggttttact gtggcagact tgcgacctcg cttggcaggg gaccgttctt 420
cttcagaagt gataagtttt cttttgcctg agagaactcc catggaggca cgaggacttt 480
ctgtgatctt tcgggtaggg gttgtgctgc tactggaggc agtanggggtg gctggggagc 540
tgacgttact gcgcggttcc cgcttctctc caccaaattg ctaagctgat atctgctgcc 600
tttghtaagaa gnggtactgc ttcatanggg ccaagcccat ac 642
```

&lt;210&gt; 151

&lt;211&gt; 322

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (1) ... (322)

&lt;223&gt; n=A,T,C or G

&lt;400&gt; 151

```
nttgacaac atcttccccg ctatgctgga attacttcgg tgttctgcgg tggccatggt 60
gaacatctga tgaactgaaa ttccatcgga atgcacagga agatatagtt gatcttcaaa 120
aatgtccttt ccaggaccac catactgggg aagttcttcc ggggtgcctgc naatgggctg 180
caccctgggg ctggggccga gctctagctc tgtcatgcca tcgccactga aatcggtttt 240
cagatgatta gtctcttcat gccccgtcca tttttcggtt tttctccagt gttcagaaat 300
tcaaatgatt aacttctggg aa 322
```

&lt;210&gt; 152

<211> 262  
<212> DNA  
<213> Homo sapiens

<400> 152  
acaaagtctt ctctttgctt tttataattt taaagcaa ataacacattta actgtattta 60  
agtctgtgca aataatcctt cagaagaa atccaagatt ctgtttgcag aggtcatttt 120  
gtctctcaaa gatgattaaa tgagtttgct tttagaataa agtgctcctg tccagcagaa 180  
ctcaaaaggc cttcaagctg ttcagtaagt gtagttcaga taagactccg tcatacgaat 240  
tccagcttcc cgtgccact gt 262

<210> 153  
<211> 284  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)  
<223> n=A,T,C or G

<400> 153  
ctcgggagta aaagggtgcc cttggtagca atgatattcc agaattaaat gggtttttgt 60  
tgccatggag actgcattta tataaatgta gcctgtagct taagttaact aaacctaatg 120  
ctgctgttaa aaacagttta ttttaattt aaaatacagt tgattagcaa cagcgggtgct 180  
gtattttaag agacacttta ttggaagtgc aatcatagtt atttggtttc acaattttac 240  
ngtgcattct aattactgat gggngcaatt acttttaate gngg 284

<210> 154  
<211> 531  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(531)  
<223> n=A,T,C or G

<400> 154  
accacccta aatttgaact cttatcaaga ggctgatgaa tctgaccatc aaataggata 60  
ggatggacct ttttttgagt tcattgtata aacaaatttt ctgatttgga cttaattccc 120  
aaaggattag gtctactcct gctcattcac tctttcaaag ctctgtccac tctaactttt 180  
ctccagtgtc atagataggg aattgctcac tgcgtgccta gtctttcttc acttacctgg 240  
cctctgatag aaacagttgc cctctcatt tcataaggtc gaggacttgt gaccctggat 300  
ggttctaaat ggaaaaagca cggccagatt gtgaaacctg gcttcaacat cagcattctg 360  
aaaatattca tcaccatgat gtctgagagt gtteggatga tgctgaacaa atgggaggaa 420  
cacattgccc aaaactcacg tctggagctc tttcaacatg tctccctgat gaccctggac 480  
agcatcatga agtgtgcctt cagccaccag ggcagcatcc agttngacag t 531

<210> 155  
<211> 353  
<212> DNA  
<213> Homo sapiens

<220>

<221> misc\_feature  
<222> (1)...(353)  
<223> n=A,T,C or G

<400> 155  
tcttgacaag actgagagag ttacatgttg ggaaaaaaaa agaagcatta acttagtaga 60  
actgaaccag gagcattaag ttctgaaatt ttgaatcatc tctgaaatga agcagggtga 120  
gcctgccctc tcatcaatcc gtctgggtgc cagaactcaa gggtcagtgg acacatcccc 180  
ctgttagaga cctcatggg ctaggacttt tcatctagga tagattcaag acctttacct 240  
canaattatg taaactgtga ttgtgtttta gaaaaattat tatttgctaa aaccatttaa 300  
gtctttgtat atgtgtaaat gatcacaaaa atgtatttta taaaatgttc tgt 353

<210> 156  
<211> 169  
<212> DNA  
<213> Homo sapiens

<400> 156  
agtttgttct actacatttg tgggccacta gttcactttg ctgtgttgat aagcggtacc 60  
accaattgca ctttctatag cctcttttac aatgttgctc acttcatcaa caacaaaagc 120  
agtctctccc gcagcctggg agtcttccat ctttctccc gcgcgtccc 169

<210> 157  
<211> 402  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(402)  
<223> n=A,T,C or G

<400> 157  
gttaactacc cgctccgaga cgggattgat gacgagtcct atgaggccat tttcaagccg 60  
gtcatgtcca aagtaatgga gatgttcag cctagtgcgg tgggtcttaca gtgtggctca 120  
gactccctat ctggggatcg gttaggntgc tttaatctac tatcaaagga cacgccaagt 180  
gtgtggaatt tgtcaagagc tttaacctgc ctatgctgat gctgggaggc ggtggttaca 240  
ccattcgtaa cgttgcccgg tgctggacat atgagacagc tgtggccctg gatacggaga 300  
tccctaataga gcttccatac aatgactact ttgaatactt tggaccagat ttcaagctcc 360  
acatcagtc tccaacatg actaaccaga acacgaatga gt 402

<210> 158  
<211> 546  
<212> DNA  
<213> Homo sapiens

<400> 158  
actttgggct ccagacttca ctgtccttag gcattgaaac catcacctgg tttgcattct 60  
tcatgactga ggttaactta aaacaaaaat ggtaggaaag ctttcttatg cttegggtaa 120  
gagacaaatt tgcttttgta gaattggtgg ctgagaaagg cagacagggc ctgattaaag 180  
aagacatttg tcaccactag ccaccaagt aagttgtgga acccaaagg gacggccatg 240  
gaaacgtaga tcatcagctc tgctaagtag ttaggggaag aaacatattc aaaccagtct 300  
ccaaatggat cctgtgggta cagtgaatga cactcctgc tttatttttc ctgagattgc 360  
cgagaataac atggcactta tactgatggg cagatgacca gatgaacatc atcatccaa 420  
gaatatggaa ccaccgtgct tgcataata gatttttccc tggtatgtag gcattcctgc 480  
catccattgg cacttggctc agcacagtta ggccaacaag gacataatag acaagtccaa 540

aacagt

546

&lt;210&gt; 159

&lt;211&gt; 145

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (1)...(145)

&lt;223&gt; n=A,T,C or G

&lt;400&gt; 159

acttttgcta taagtttcct aaaaatattt aatacttttt tttttcaatt taaattaaat 60  
ctnttgatga acaggggggg gntggcaaaa tttccaagcn ctggactgga attttganan 120  
aggcatttac ngaccctnat aactt 145

&lt;210&gt; 160

&lt;211&gt; 405

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 160

tgtaaatcgc tgtttgatt tcttgatttt ataacagggc ggctgggttaa tatctcacac 60  
agtttaaaaa atcagcccct aatttctcca tgtttacact tcaatctgca ggcttcttaa 120  
agtgcagta tcctttaacc tgccaccagt gtccccctc cggccccctg cttgtaaaaa 180  
ggggaggaga attagccaaa cactgtaagc ttttaagaaa aacaaagttt taaacgaaat 240  
actgctctgt ccagaggctt taaaactggt gcaattacag caaaaaggga ttctgtagct 300  
ttaacttgta aaccacatct tttttgcact tttttataa gcaaaaacgt gccgtttaaa 360  
ccactggatc tatctaaatg ccgatttgag ttcgcgacac tatgt 405

&lt;210&gt; 161

&lt;211&gt; 443

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (1)...(443)

&lt;223&gt; n=A,T,C or G

&lt;400&gt; 161

tttgctttta atgaaggaca agggattaag acncatagag actggccana caaatgggaa 60  
accgaccaga ccagcccatg accaaaatat cacaggcaga ccaccacaa atgcagaggc 120  
ctcagagtcc acagtgggcg gttggaaccc agggccccag ggaatctttc agctgcattc 180  
cggctgtgat cggcgggcaa caggtagagg tgctggaggg ggctgagtcg tgattttcgg 240  
tgtctgtcat attcgatcaa gtgtgtcata gagcttctg tttcatctcc cagttattca 300  
aggagaggct ggtggctcca cttcccagg aactgtgctg tgaagatctg aagacaggca 360  
cgggctcagg caccgcttgt ctggaatgtc aatttgaaac ttaaaaagca gcgaccatcc 420  
agtcatttat ttccctccat tcc 443

&lt;210&gt; 162

&lt;211&gt; 228

&lt;212&gt; DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<222> (1)...(228)

<223> n=A,T,C or G

<400> 162

```
tcgttatcaa aatggaagac accaaacccat tactggcttc taagctgaca gaaaaggagg 60
aagaaatcgt ggactagtgg agtaaatttt atgcttnctc aggggaacat gaaaaatgcg 120
gacagtatat tcagaaaggc tattccnagc tcaagatata tnattgtgaa ctanaaaata 180
tagcanaatt tgagggcctg acagacttct canatacntt caagttgt 228
```

<210> 163

<211> 580

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<222> (1)...(580)

<223> n=A,T,C or G

<400> 163

```
acccaaggct acacatcctt ctgtgaaaca gtctcacgga gactctcaga atcccaagaa 60
ttttcttcaa ctttcttttg ttttgattct gaagggaaca tctgatctgc tctcaatggt 120
tggtcattct tcaattccaa ggcttttatt ggaacagact ttgcatttca atggcaggct 180
cgaaggcaga tggcttctcg ggaggctctg ctttgaaagt ttgcntgtcc atcaattcta 240
aggctttagn tggaatagaa actttcattc tgcagggagc cttcagaaaa ccatcattat 300
caggagactc ttctaatttt ccatttattt tatctatttc tttttgatgc gcagccttgg 360
gtanacacac atccttctgt gaaacagtct cacagagact ctcagaatcc caagaacttt 420
cttcatagtc cttttgtttg gattctgatg ggagtatctc atctgctctc aatgtttgtt 480
cattcttcaa ttccaaggct ttatttgga cagacttttg catttcaatg gcaggctcga 540
aggcagatgg cttctcggga ggctctgctt tgaaaagtgt 580
```

210> 164

<211> 140

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<222> (1)...(140)

<223> n=A,T,C or G

<400> 164

```
acttatatct tttggncttg ggcttctcaa agttcacgac agacataggc actctcacag 60
tatcaagccc atttaccgnc acctcacacc aatactcgcc ccaccgngng ataggntctg 120
ctggnaactt taatgnatgn 140
```

<210> 165

<211> 370

<212> DNA

<213> Homo sapiens



<220>  
<221> misc\_feature  
<222> (1)...(370)  
<223> n=A,T,C or G

<400> 165  
acatggagcc actgccacca gtggtgatgg aaagcactgc cttcttactc cggaagggtc 60  
ctttgtcata catggcagcg taagtgttaag caaactctcc tatgaacact cgctcaaacc 120  
agcctttcag aatggcaggg actccaaacc actgcnnngg ggaactggaa tatcacaagg 180  
tctgaggctt ccagcttctt ttgttcagcc acaatatctg ggctcanatg gnetttctta 240  
taagccagaa cagactcggg aggatactga aagttcgcag ggccttcan tttacctgng 300  
atgncctttn tggaaatgat gggattgaag ntcatggnat aaaggncga ctncaccacc 360  
tccattcttt 370

<210> 166  
<211> 258  
<212> DNA  
<213> Homo sapiens

<400> 166  
gtcaaaagtc atgattttta tcttagttct tcattactgc attgaaaagg aaaacctgtc 60  
tgagaaaatg cctgacagtt taatttaaaa ctatggtgta agtctttgac aagaaaaaaa 120  
aacaacaaaa cacttcttct catcagtaac actggcaatc ttctgtttaa ccactctcct 180  
tagggatggg atctgaaaca acaatggtca ccctcttgag attcgtttta agtgaattc 240  
cataatgagc agagggtgt 258

<210> 167  
<211> 345  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(345)  
<223> n=A,T,C or G

<400> 167  
ggtcagccaa acaccagga tctctgtaaa actgaagaac aggncaatgc caccaacaaa 60  
tctcaaaacc tctccagcat attctectat gattggagca catggngagc acnantgggtc 120  
acttttaaca canctagcca gacaggngnc atttgggtta acacttcgga acccacagca 180  
ntttanantt ctctggatgt catttcgagc acttgtatct attggtcann tttctgtatc 240  
tngcgcttgg ttagccctga accaggagca acagggncag cttctggagg ntggttggaa 300  
caatacggca agtgntngaa atgacatcca acctncngaa atgac 345

<210> 168  
<211> 61  
<212> DNA  
<213> Homo sapiens

<400> 168  
gatatgtgtg tttatggact gaggtcaaaa tctaagaagt ttgcagacc tgacatccag 60  
t 61

<210> 169

<211> 344  
<212> DNA  
<213> Homo sapiens

<400> 169  
acattggtgc tataaatata aatgctactt atgaagcatg aaattaagct tcttttttct 60  
tcaagttttt tctcttgtct agcaatctgt taggcttctg aaccaagacc aaatgtttac 120  
gttcctctgc tgcataccaa cgttactcca aacaataaaa aatctatcat ttctgctctg 180  
tgctgaggaa tggaaaatga aacccccacc ccctgacccc taggactata cagtggaaac 240  
tgttcattgc tgatgaatgc agcagtcacc aaaaaatata cccaatcttc cagataacct 300  
cagtgcactt taggaaatca aaaattacct ggaagcaatt tagt 344

<210> 170  
<211> 114  
<212> DNA  
<213> Homo sapiens

<400> 170  
agcagtggtg cctccatgaa taaacaggag ttctggaggc ccatcttctg catcttctgc 60  
tgattgttct tccccaattt tacttaaatc ccacacattc aggcggcggt cagt 114

<210> 171  
<211> 150  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(150)  
<223> n=A,T,C or G

<400> 171  
actgagagca ttataatct gaccaaattc ataggcatta ttaggcttgg ctatcggaag 60  
ttctcagggg tctctggng acctgctgct ttgctctccc ttctcanaag caaggcatcc 120  
catggagacc tcccctgcag ggcttccagg 150

<210> 172  
<211> 435  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(435)  
<223> n=A,T,C or G

<400> 172  
atttgtttcc cactgcctca cactagtgag ctgtgccaa tagtagtgtg acacctgtgt 60  
tgtcatttcc cacatcacgt aagagcttcc aaggaaagcc aaatcccaga tgagtctcag 120  
agagggatca atatgtccat gattatcttc tggtttaggt ctacagtcaa tgtgatgggt 180  
gtctttgtct cccagtctgc cagaatatct ttgtgcttct ctaatcattg gctttaaaag 240  
taatcaatgt gttggcagca tctctgtcac tcttgtttaa cacgtgaaga aatcaggtag 300  
atctttttct gtggcattgt ttctggacct aaaatcagggt atgctgacta tttccaaggg 360  
gtttttcagt tgcttcattt gcttgtaaag cagggaatcc tcttgntgct tttcttttcc 420  
tcgatgagcc cgtgt 435

<210> 173

<211> 622  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(622)  
<223> n=A,T,C or G

<400> 173  
actgntttcc cccaagtcca tgacatgtat acataattaa tggtttgcct ccttgattgt 60  
tttctccaac atccagacat agaggtgac caacgctttt aatgtatcca gatataacag 120  
gattaagggtc tggcacatac acctctggat aaatgttggt cagataccat gtaaaatttt 180  
tacctgaag gcggtgtttt atttcaaata tttttgaaag atcaccaaata gctttttgtt 240  
taacaatttt tgctgcatct gtatttctcc tataaaatat ttcttgtat tcatccatcc 300  
agacttctgc aaggcgaaact tggtttctag caatcacctg agtgcctttt ggaaagctat 360  
gagggctttt gctgcgaaaa acatgtccaa caacagagca aggcataatc tccaactgcc 420  
caccacattg ccatactctg aaagacattt ctatattttc acctccccag atttccattt 480  
cttcatcata gcttccaata tactcaaaat attcttttga tatggaaaaa agtcctcctg 540  
caaaagtggg tgttttaatt gggtaggggt catctttcct tctttgcttc tcatgatcag 600  
gaagcgactt ccaccaatg aa 622

<210> 174  
<211> 362  
<212> DNA  
<213> Homo sapiens

<400> 174  
acgggtgcagt tgaccactg ttggtctctc ttgcagttcc tgatatgtca tctttagcat 60  
gtggctactt acgtaatctt acctggacac tttctaactt ttgccgcaac aagaatcctg 120  
caccctccgat agatgctgtt gacgagattc ttcctacctt agttcagctc ctgcatcatg 180  
atgatccaga agtggttagca gatacctgct gggctatttc ctacctact gatgggtccaa 240  
atgaacgaat tggcatgggt gtgaaaacag gagttgtgct ccaacttgtg aagcttctag 300  
gagcttctga attgccaat gtgactcctg ccctaagagc catagggaat attgtcactg 360  
gt 362

<210> 175  
<211> 486  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(486)  
<223> n=A,T,C or G

<400> 175  
acagntnctc tactacactc agcctcttat gtgccagtt tttctttaag caatgagaaa 60  
ttgctcatgt tcttcatctt ctcaaatac cagaggccga agaaaaaac tttggctgtg 120  
tctaaaaact gacacagtca atagaatgaa gaaaattaga gtagttatgt gattatttca 180  
gctcttgacc tgtccctctt ggctgcctct gactctgaat ctcccaaaga gagaaaccaa 240  
tttctaagag gactggattg cagaagactc ggggacaaca tttgatccaa gatcttaaat 300  
gttatattga taaccatgct cagcaatgag ctattagatt cattttggga aatctccata 360  
atttcaattt gtaaaccttg ttaagacctg tctacattgt tatatgtgtg tgacttgagt 420  
aatgttatca acgtttttgt aaatatttac tatgttttct tattagctaa attccaacaa 480  
ttttgt 486

<210> 176  
<211> 461  
<212> DNA  
<213> Homo sapiens

<400> 176  
accctggcca ctcccttctt tttggctggc caatgtctcc tctgtaggct ccagaaggct 60  
ctcagggatg caggcggcct cctgcagggt tgagttgcaa tgggaacaaa gacagctgtg 120  
gtcccatagc accctcatct ggtgacatcc tgctactgac agtcaaaaaga agccttccca 180  
gatgaaattt tagtcctctg cgcagccatg ctcttcttcc agcaaaaagag ccatgtgcag 240  
tcgggtctgc tcccatggg ggctttgatg tgggcccagc agtggatcag ccttccagac 300  
acgctcaact ctgcacactc ttcttgccgc ctcaggcttt ccaggaccct cccgagcctt 360  
atcagagtcc ttaccctcag ggctactgat accttgctgg gtgaccttgg acagattcac 420  
ttacctggac tcagtttcat aatatgaaaa tgatagggtt g 461

<210> 177  
<211> 234  
<212> DNA  
<213> Homo sapiens

<400> 177  
acacattttg taattacctt ttttgttgtt ttgtagcaac catttgtaaa acattccaaa 60  
taattccaca gtctgaagc agcaatcgaa tccctttctc acttttggaa ggtgactttt 120  
caccttaatg catattcccc tctccataga ggagaggaaa aggtgtaggc ctgcccttacc 180  
gagagccaaa cagagcccag ggagactccg ctgtgggaaa cctcattgtt ctgt 234

<210> 178  
<211> 657  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1) ... (657)  
<223> n=A,T,C or G

<400> 178  
gagcteggan ccctagtaac ggccgccagg gtgctggnat gngcccttgc gagecngncg 60  
cccgggcagg nacttttntc cccctcctac ttctgttagc tcatttgtnt ctctcatttt 120  
ttggcatatt tttcaagtca cacttaaaaa ctcttccatg tattcacttc tcatcacttg 180  
gtctacatgc cgaacctaag gtcaggattc caaaaagatg agtatcctct caaacgcctc 240  
ctaagcctct ggtatacatg actttggctg tgcacttcat ttagacttca cctttttgtt 300  
tgctgttgtt ttttactacta gattcctttg tcttcattaa agataatgaa agattcacat 360  
cacagtgcag ctcttcgctt tgtcctttcg taagtccgta gcaactgccg agagttcttg 420  
tctgctaggc atgtgtgaaa tccgctttgt ggctctctgt gatttgttcc gcttaacggt 480  
tttatttgtc ttattttacac atgccaaggt ggcaacgtga aaaatgtctc tgacgctatt 540  
ttccgactgt aaagctgagc attcgatata agtagctgct ccaatctgtt tggccatact 600  
tgccccctgg tcataggaca ctggcgtctg cctgtgattg gagagctcta ctaatgt 657

<210> 179  
<211> 182  
<212> DNA  
<213> Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (1)...(182)

&lt;223&gt; n=A,T,C or G

&lt;400&gt; 179

```

acaaaanctt ttaaatttta tattattttg aaactttgct ttgggtttgt ggcaccctgg 60
ccaccccatc ttgctgtgac agcctctgca gtccgtgggc tggcagtttg ttgatctttt 120
aagtttcctt ccctaccag tccccatttt ctggtaaggt ttctaggagg tctgttaggt 180
gt 182

```

&lt;210&gt; 180

&lt;211&gt; 525

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 180

```

acacgctttt ggccccgacc aatgaggcct tcgagaagat ccctagttag actttgaacc 60
gtatcctggg cgacccagaa gccctgagag acctgctgaa caaccacatc ttgaagttag 120
ctatgtgtgc tgaagccatc gttgcggggc tgtctgtaga gaccctggag ggcattgacac 180
tggagggtggg ctgcagcggg gacatgctca ctatcaacgg gaaggcgatc atctccaata 240
aagacatcct agccaccaac ggggtgatcc actacattga tgagctactc atcccagact 300
cagccaagac actatttgaa ttggctgcag agtctgatgt gtccacagcc attgaccttt 360
tcagacaagc cggcctcggc aatcatctct ctggaagtga gcggttgacc ctccctggctc 420
ccctgaattc tgtattcaaa gatggaaccc ctccaattga tgcccataca aggaatttgc 480
tcggaacca cataattaaa gaccagctgg cctctaagta tctgt 525

```

&lt;210&gt; 181

&lt;211&gt; 444

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 181

```

acaccacaat gtgcatcaag gagacgtgcc gattgattcc tgcagtcctg tccatttcca 60
gagatctcag caagccactt accttcccag atggatgcac attgcctgca gggatcaccc 120
tggttcttag tatttggggt cttcaccaca atcctgctgt ctggaaaaac ccaaagggtc 180
ctgacccctt gaggttctct caggagaatt ctgatcagag acacccctat gcctacttac 240
cattctcagc tggatcaagg aactgcattg ggcaggagtt tgccatgatt gagttaaaag 300
taaccattgc cttgattctg ctccacttca gactgactcc agacccacc aggctcttta 360
ctttcccaaa ccattttatc ctcaagccca agaatgggat gtatttgcac ctgaagaaac 420
tctctgaatg ttgatctca ggggt 444

```

&lt;210&gt; 182

&lt;211&gt; 441

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 182

```

acaaccttta ttgcttctcc agcattttcc agaagaatgg tgtcattaga gggccacagg 60
ggatggggga gtaaaaaata acataaacga actgaacaga aatgcaggag ggtggcaaga 120
ggggccgaga ttgggtgttc agggcagaga ggtggaagac caggggcagt cagtgttct 180
tagctttcag ccaccagagt ggagaattcg tcaaccccaa ttttgccgtc cccatctttg 240
tctccagcag ccatacagcat cttggtttct ttagcagaca ggtctctggc atctggggag 300
aagcctttta ggatgaatcc cagctcatcc tcctcgatga agccactttg tccttggtca 360
gcattgtgaa cactttcttc acatcatccg cactcttttt cttcaggccg accatttggg 420
agaacttttt gtggtcgaag g 441

```

<210> 183  
<211> 339  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(339)  
<223> n=A,T,C or G

<400> 183  
tgtntcatcn taaggggatt gggctctaga tctgtcgacg gcgcattgag gatttgcnat 60  
cgggttangtg gtccgcgagt catgaatttt tgctctggag cgttattgtt tgtgaagttt 120  
atccaggaga gaactatgat tgtgtcgatg cgtttactgc aggaagantc acggtctcag 180  
tcacggaggt gtaaggggtg actgactgan tgagacaagg gatattntngt tnttatannc 240  
ttgtgatgaa cctgcctacc gtttatgtct ctttgcataat gggctctcng tncgtgnatt 300  
cncncaagct gcgggggctt ccncggttct gggctctga 339

<210> 184  
<211> 490  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(490)  
<223> n=A,T,C or G

<400> 184  
atatagcaag cttgtacgac cgacacatac ggcgcatgtg gctggattgc ttatcttgtc 60  
gcgcgacgtc tatataancg anactacata gtctcggaat tccactcant ttcaagttcc 120  
caaaaanacng ganaaaaaacc catgccttat ttaactaanc atcagctcgc ttctccttct 180  
gtaaccgcgc ttntngctcc cagcctatag aagggtaaaa ccacactcgc tgcgncagtc 240  
atcnnataac tgattcgccc gggactgccc gggcggcgct cganaccaat tngcanaatt 300  
cacacattgc ggcgctcnan aagctctaga aggccaatcg ccatattgat ctatacatta 360  
tggccgtcgt tnacacgtcg tgacgggana ncctggngta ccattaatcg ctgcacantc 420  
ccttcgcagc tggggntnac aaaagccgcc catcctcca cgttgcgncg gatggcaagg 480  
acnccctnat 490

<210> 185  
<211> 368  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(368)  
<223> n=A,T,C or G

<400> 185  
ctnnanatag cangettgtg cgaccgacac aatacggcca ntgtgctgga ttcgcttcag 60  
cgccgcccgg gcagtaaccg cgctcatcta tcngatgatg gcgcaccaat gtggggtttt 120  
aaccttttta tatggctggg gacanaaagc gcggttacnn aaccnataac gagctgatgg 180  
tcatttaaaa atgcttgggg ttttcccggt cttttgggga attgaaactg agtgggactt 240

canaaactgt gctacttttcg cttatctaag tactcgggcg caacacctag ccgaatccgc 300  
 anatatcatc acnctggggcg gcgtcancat gcntctaaag ggccaattcn cctanatgag 360  
 tcttatac 368

<210> 186  
 <211> 214  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <222> (1)...(214)  
 <223> n=A,T,C or G

<400> 186  
 ngggagatcg cagcttgtag gactcgatcat ataacggnca atgtgctgga tcgcttcanc 60  
 gccgcggcg gtctaactcg gtccggattn tgtgtgtntt gtctntntta canggtgcta 120  
 tccccctctt cctcctctc tgccatctc atcctttatc tcctttttgg acaagtgtca 180  
 nancagacag angcagggtg gtggcaccgt tgaa 214

<210> 187  
 <211> 630  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <222> (1)...(630)  
 <223> n=A,T,C or G

<400> 187  
 cagctgggac gagtcgatca tatacggcg atgtgttgna tcgctatcgt gtccggcgag 60  
 tanttattan attactgtta tttctgctcc tactggatat gatctcttga nggcangtct 120  
 gtgtcgtctg gtcacaccat gttctcaggc tgggcaaata ccttctata atagtttatg 180  
 gataatgaat gacgactang tctanaaana cgctagctaa ataacacact cagggaaga 240  
 gtcttaaata ttgtgaagggt gtttttanta tacaacnttt gtttacataa taggaaataa 300  
 tttttagact tttaaacaga cacttgagcc agatttgta atgttaccat ctatagtgtc 360  
 ttgaaaatat tcctcttagt ttccaatatg aatgaatcta aaatccatct tttcaattat 420  
 gcccgagccc gtggtcaatg cncctcnac acttcattaa cggattatac cttgggaaac 480  
 cataatctgg cntaggacga atgcctggc ncangctaan aactgcctg tattgagggg 540  
 ttatnnctga ttgcnaggt gcctctccag gtccccaaag ggtcgtactg ttgaanctgg 600  
 ctctaanttt ntcttgctn acaggctctc 630

<210> 188  
 <211> 441  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <222> (1)...(441)  
 <223> n=A,T,C or G

<400> 188  
 cnngeaanac anggtcggat tccngtgagg naanaattcc ctnatagggc tcgcccccta 60

```

ttcacaaac caancgaaa ctcttgcggt caaatctaag ctatnncaca accccactct 120
gnagggtatg cgccccgccc ctgcaatgaa atcaatanca tatttggaga cagagagata 180
gagagagaga ggttccctggc cttnnctatt ctgctcttac ttggnagatn tcaganatag 240
aaaaacctat cctaggtccn nccaatgatn gcggcttncg aatccccgng tggccantcc 300
ccggatcgga ctaaatcaaa gaagatcctc cgtctcctg ttctccaca ctggagtccc 360
attgtatgca tgggtntttc actggctnat cataccnag gatctgtcca ccttnaactc 420
ttctctngga antccctncc c 441

```

```

<210> 189
<211> 637
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<222> (1)...(637)
<223> n=A,T,C or G

```

```

<400> 189
agggngtata taccacttg tacnaactga tcatanacgc gcatntctga atcgcttntc 60
ggccgcgatg tactgtgggc acttaagcac tgagtactgt ttgcgtcatg ccnggtcana 120
agatgctgct gcaaaggagc tccaacnaaa tacactgtct tcaacaggag ttaacacctc 180
acacttgggt ganaanagaa ctcaactggtg gtgatgcaca cgactgnatc catcaagtgc 240
gtttgcctgt tgactgctaa ccaaggtctt ggcagtacct gcccgggcgg cgctcgaaac 300
caaatctgca aatatcatca cactggcggn cgtcagcat catctanaag gccatcgct 360
atagtgaatc tatacatcat ggccgcnttt acactcctac tggaaaaacct gcgtaccact 420
taatcgcttc acacatcccc ttctcgngtn gcttatannc aaaagccac gatgcctcca 480
cattgcncnc tgatggcatg ancccttac gcgcatancc gcggtntgtg taccncangt 540
accgtntcgc acgctacnnc tcttctctct cctcttcccc ttcccgttcc tcaccattcg 600
gggccttagg tenatatctc gnccacccaa atntagg 637

```

```

<210> 190
<211> 653
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<222> (1)...(653)
<223> n=A,T,C or G

```

```

<400> 190
agggggtata taccacttg tacgactgna tcatatacgc gcatgtctgg aatcgcttnc 60
gtggctgcca tgtattgaca ctacttctaa gaactacaaa agtgatactg angatacatt 120
acacagaang gctnacattc tcncagatcc tcatttntca tgatatgtgg acatcangan 180
cacgtggata agtgtatcta aanaatggct ttcaaaatat ttccacttta ttaaggtttg 240
acatganatt cataaaatgt cttaatacta tttctnaaaa taacatctaa tcggaaacta 300
tgctnaact gcacnttttn tgtgtanata atcntanttg tacgcccggc ggcgccaaag 360
ccnaatctgc gattcctcac ctggcgccgc tcaacatcat ctaaaaggcca atcgctata 420
ntantctata catcctggcc gcgttttacac gtctaattgg aaaccggcgt accacttatc 480
gcttgagca ctccccctcc cactgggtta tacnaaagcc gcncgatgcc tcccacattc 540
canctgatgc aatgaccctt gttcgctta ncccgcggtt tgtgtaccca ntnaccacnt 600
cagcgtgcn cntctctntt ctctctctct gccntnctg tccctcactc nng 653

```

```

<210> 191

```



<211> 663  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(663)  
<223> n=A,T,C or G

<400> 191  
angnggtata taccactgt ncgactcgat catatacgcg catgtcggat cggctccanc 60  
gcgcggcat gtactatata tacatcaact gtattatcat ttanatattg atnaaagaca 120  
aaatcatact tccatctgct cactgatgat aattactatg atacatgatc atgtaaactg 180  
atcaatataa caatggaaga tccctctgac tatgcaagcc taattttcca atcncatgca 240  
ctctcatagc tcaaanatnt cacngacatc ctgatgaaac tatnatacan tttccacaca 300  
aatcacttcg ctttagatct ctccattatt cttgcttttc cccctaaca actacaaatc 360  
ctcntgggat gggaagaata tatatcatct actaaaaata atatataatc ccctgcanat 420  
ttgtggnaaa tcnggtgtct caanagccac aggagnacaa gggggnacca actaggactt 480  
ttgtatgctt atctctgtac tcgcgcacac ctaagcgatt ctgcnattct ccctggcggc 540  
gtcacanctc tanaggccat cncnatatga tctatacatc ntggcgtctt tacactctga 600  
cggaaaccgg gtnccantta ccctggacca tcccttcgcn ctgntataca aagccccga 660  
ncc 663

<210> 192  
<211> 361  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(361)  
<223> n=A,T,C or G

<400> 192  
antttttata taccactgg tacaactcga nccatacgg cgcanttneg gaatcanctt 60  
cancggcgcc ggcatgtacc ggtnatcatc atcngatgat ggcgctcnaa tgtgggtttt 120  
acctnttata cggctgagat canatcgct acataacaaa nncaactgat ggtnaatnta 180  
aatncggttg ggttctccn ntctgttggg gaacttgana ctgagtngna cntccatana 240  
cgtgctattn tcggctancn antcctcagc gnacacctat ngnagtgcgc naattcatcc 300  
atgntggcct cgactnttc aaaangccnt ncgcccacnt gntcgnana cantctcggc 360  
c 361

<210> 193  
<211> 314  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(314)  
<223> n=A,T,C or G

<400> 193  
aggngnata taccaactgg tncgactcga tccatacgc gcatttcgga ttcgcttcaa 60  
cggcgccggc atgtacaaa cctcaatccc aaccgtctca nttngacggg ctcagttctg 120  
tcacagccac cccacatttc ttttgtttt tctgccactt caaagaatt ccaataaga 180

attctgtctgc agctccgtac aaggatatgg gcagcacagc acacacagag tngtgcctct 240  
cacactttctc tggnaatgtc tcgtgaatat ctcaacagtc angaagtggg gcgttatcaa 300  
aaacaatcag gcc 314

<210> 194  
<211> 550  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1) ... (550)  
<223> n=A,T,C or G

<400> 194  
aggngngata taccactgg tncgactcga tcctatacgc gcatgtcggg ncgctatgtg 60  
gtcncgcaag tacctcttct gcagtgatgg tctgtntcct ctatgatnag tgategaata 120  
atcatcgaat tcancgaaag ttattcgagt gatatntgtg gctttagaa tctatgtctc 180  
atgggtgtgg cactgtcaag attaacacag aatggaagan ncngcactgc ataaaagatg 240  
ttgtcaaatt ggggtgcgtg atcngatagc tcntcccaag aggtcantgg tgttcaggat 300  
tncnacataa gatnttggat caccngacga ccagangata ccngtgcaaa ctgtgaancn 360  
ngtaatctgc ctatncctgc cctctcggan gatccctcgg ggacgacgag atcattcttg 420  
aaacagcnan tgatagtcca gtnnangatt gatgancgac ganacgcntg atanatgtct 480  
gacgtgagat tnggatgtga atcttccnt gtgtgacctg cncntaccn aanggtgcgn 540  
ctccactcnn 550

<210> 195  
<211> 452  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1) ... (452)  
<223> n=A,T,C or G

<400> 195  
nngcgggnat gataccaact ggtacgaact cganctctat nacggcgctn tttcnngatc 60  
tgctatgtgg tctcggcaat gtacattata acngggcana catataatct acntctgtct 120  
ttntctcccc cngagagcgc aancatctcc aaatcgggtt ctgggtcatc caatgggtctc 180  
cantaatcac acaactcata tatatttatg gaangtgtct gtcacgtcc ccacgangga 240  
agtnncgtcg ctgtntgtct gtcactagg tngtactctc cagtacttga aanctggtna 300  
nggctgtctg tngtactggc cggcgccctc gaaancgaat ctgtnnatat catcacatng 360  
cgncgcccga ncatcactna gggncanttc gcctatactg atcgtntgcg annctgcgn 420  
cncttacacg tcgnacggga naccggcctt cc 452

<210> 196  
<211> 429  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1) ... (429)  
<223> n=A,T,C or G

```

<400> 196
gcggggnnat gataccagct ngtagcactc gatcctataa cggcgcatgt gngtatcggc 60
tacgtgtctc ggcgatgtac atataacggg gcaacatata atnatacant ctgtcttttt 120
ctcccccgga aacgggaacc atctccaata tcggtctggg tctccaatgg tctccaacta 180
aatcacacaa gtcaaatata nttanggaaa gtgtctgtct cntccccaga aggagtancg 240
ttagtgttg tctgtcatta ggttggtacc tccagtnaca tgaaaactgg tgagggtgtc 300
cttgtagaag ctctgcctca ccagatccta tactattagg gggccacgg ttatctatct 360
taagggtctn aaaacctgga cttcatctgc tccggcggan gaatgtccg cttacttacg 420
ntgttccac 429

```

```

<210> 197
<211> 471
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<222> (1)...(471)
<223> n=A,T,C or G

```

```

<400> 197
atgatacgca gctngtacga gccgtcacta tnacggcnca ttgtgtggat tcnngctntga 60
tcggcgcccg ggcatgtcca tcnagagcgc atcatgggan tgnactcccc atatnntgac 120
caangttcgc gcaaggagcc naganccgat actacctgag ctgtcgtctn gttatacacg 180
tttctggcca angancaact ccacatncaa caagttggtg ttgaaatggt gtttatnagt 240
ccaccaaccg gccgctctgt cccttcccca tgatccgaag ataaagcttcc tgtccggaan 300
acgaacggcg tgggtgtgngg acatantgat atgtgcgggt caggaagtac tcgncgcaac 360
ncgcaagcna atctgcnata tcatcacctg gcggcgctcg agctgccana ngcccnttcg 420
cctatatgag tctatacatt cctggcgcgc tnttacactc ngacgggaaa c 471

```

```

<210> 198
<211> 643
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<222> (1)...(643)
<223> n=A,T,C or G

```

```

<400> 198
tngtncgacc gtcactatac gcccatgtgt ggatccgntc cacggcgccg ggcangtacg 60
anactatatt gatectctga tattgaaagt tgggtctanca ataaccttta angcaaatca 120
ctcantgagt tttgaccaga agtcaccaca tcatgaatca cagtctatgg caaatgatac 180
cagtgtctct aagtcctatg ctcaaggtaa gagcatgcta ttccggttta catttactgg 240
aatttactgt tcattcatna ttaaaatctc tagttttcat cctcaactgt ctaanaccag 300
tgtgcacaga cttaagactc tgttctctc attttctcca acagaaacat tctcagtgtc 360
tactgttcta aaagggaatt tccgaggtgg cacttctcgg aatatcgacc ctcnngctct 420
atcaggcggt acttcnngca ctgcgtcatt gggcttggtc anttgtctta tctgtccagt 480
cacttcattt taagaaaaca attgatcgct ggtcacatgt nattcattgg cagccgggtg 540
gactgctgag tctcgcgcac acnctagcaa tcgnnattct ccatggngcg tcaactctcta 600
naggccatcc cctatatgat ctataatctg gcgtctttac act 643

```

```

<210> 199

```

<211> 292  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(292)  
<223> n=A,T,C or G

<400> 199  
nccggcnggag ttccgcagttg nacgaccgat cctatacgnc gcattttctga tccgctacnt 60  
gtccggcgag tctatgctat ttatttntga ttaaatacaat atttttcttc tgaatattaa 120  
tcttatctnt acttttatac tattgaccta gctatatgta ttganctttt tgaactccta 180  
tcagnttttt tcatgctatc gtatattttc cacttggtac ctntngctga ntctagata 240  
tcgtaaaaca tctctnnatc ntcacacnga gncagggt ctgtatngaa tt 292

<210> 200  
<211> 275  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(275)  
<223> n=A,T,C or G

<400> 200  
atacgcaagc ttggtaccga gctnggatcc ctattaaccg gccgcaatat tctggaattc 60  
tgcttanccgt ggtcncggcc gaagtactat gctatnttac ttttttgga tataaatca 120  
atatatttct tctnaagta tataaatctt atcncgtat cnttcnatac ctntctgaca 180  
ntaagcttat angtatntga tctntgttga actcctatca agtgntttcn catgctatcg 240  
tganntcttc cacnttggtg ccttttacgc tgaat 275

<210> 201  
<211> 284  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(284)  
<223> n=A,T,C or G

<400> 201  
cgnnnatcca gtgtanaccg tcnttacgcg cattctgac gtacacgcc gcgtctttat 60  
atctatctcg actgattcac ctgtcattgt aaanaattcg tgtagctgt ctaccnctta 120  
nacatcatct aatcnaacta nctgataaa tttcttcaat agggatanac ntntagtaca 180  
tacgnttcca ttgagntacn tccgaggacc cncatcgcaa acnncatgcg gtcagtcnna 240  
gcatectcta tettaatccg tccttaacct ntgaacgctc cact 284

<210> 202  
<211> 448  
<212> DNA  
<213> Homo sapiens

<220>

<221> misc\_feature

<222> (1)...(448)

<223> n=A,T,C or G

<400> 202

```
atgatacgca agcttgtagc actcggatca tataacggcc gcaatgtgct ggaattccgc 60
ttcgacggac gccgggcatg tacttttata atnctactcc tcagaccttg catctcnacc 120
gctnggtcca gtttgtaaaa acnnacttcc gtngtgcagc cctggttctg ancantctct 180
atcaenctct atcctcncat ccncaanact anatcgctg aattcatatt tattcatttt 240
ccataatgat gggggaanga ctatcnctna tnatgcttan cacnctnget gcanttcgnc 300
natctcgchna ngcntgaaac gattactctg tcgcgaaccc tctangntga attctgchna 360
atatctntna cnctggcngg cgctcnangn atgcctctcg anggccaatc cgccnngcat 420
gattctaatt anatecntng gtcccntt 448
```

<210> 203

<211> 321

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<222> (1)...(321)

<223> n=A,T,C or G

<400> 203

```
gggtgcnaga tcgcagtngt acgaatcngt catatacggc gcatgtgntg antcgctacg 60
tgtccggcga ngtaccatat aatcgaanta ncatagttct ggangcccnc tcattttcaa 120
tttcccaaaa nacgggaaaa ccnaagcctt atttaactaa ctatctgctc gcttctcgct 180
tctgtaccgc gctatntgct nccagcctat aanaagggtg aaacccacac tcggtgcgtc 240
agtctccnat atantgagtc nccgggtact ggccgggcgg tcgttcnaaa ncaattcncg 300
aanttcacta ctggcggcgc c 321
```

<210> 204

<211> 369

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<222> (1)...(369)

<223> n=A,T,C or G

<400> 204

```
ntgtngtatg tacccagtgg tacgactcga tctagtagc gcgcagtgtg ctgaatcggt 60
acttgtcgcg gccaaagtatc tataaagcaa actatcacag ttctgaaagt ccatctcant 120
ttcagttccc aaaagancgg gaaaacccaa gccttattaa actaacaatc agtcgctctc 180
gcttctgtac cgcgcttttg gccccagcc tataaaaggg taaaacccac actcgggtgcg 240
ccagtcacgc ataactgaat cgcccggtac tgcccgggcg gcgctcnann ccaaactctg 300
agatatcaca cactggcggc gctcancatg ctctagaagg ccaattcncc tatantgatt 360
ctattacaa 369
```

<210> 205

<211> 2996

<212> DNA

<213> Homo sapien

&lt;400&gt; 205

cagccaccgg	agtggatgcc	atctgcaccc	accgccctga	ccccacaggc	cctgggctgg	60
acagagagca	gctgtatttg	gagctgagcc	agctgaccca	cagcatcact	gagctggggc	120
cctacaccct	ggacagggac	agtctctatg	tcaatggttt	cacacagcgg	agctctgtgc	180
ccaccactag	cattcctggg	acccccacag	tggacctggg	aacatctggg	actccagttt	240
ctaaaccctg	tccctcggtc	gccagccctc	tccctgggtc	attcactctc	aacttcacca	300
tcaccaacct	gcggtatgag	gagaacatgc	agcaccctgg	ctccaggaag	ttcaacacca	360
cggagagggt	ccttcagggc	ctggtccctg	ttcaagagca	ccagtgttgg	ccctctgtac	420
tctggctgca	gactgacttt	gctcaggcct	gaaaaggatg	ggacagccac	tggagtggat	480
gccatctgca	cccaccaccc	tgaccccaaa	agccctaggg	tggacagaga	gcagctgtat	540
tgggagctga	gccagctgac	ccacaatatc	actgagctgg	gcccctatgc	cctggacaac	600
gacagcctct	ttgtcaatgg	tttcaatcat	cggagctctg	tgtccaccac	cagcactcct	660
gggaccccca	cagtgtatct	gggagcatct	aagactccag	cctcgatatt	tggcccttca	720
gctgccagcc	atctcctgat	actattcacc	ctcaacttca	ccatcactaa	cctgcggtat	780
gaggagaaca	tgtggcctgg	ctccaggaag	ttcaacacta	cagagagggt	ccttcagggc	840
ctgctaaggc	ccttgttcaa	gaacaccagt	ggtggccctc	tgtactctgg	ctgcaggctg	900
accttgctca	ggccagagaa	agatggggaa	gccaccggag	tggatgccat	ctgcaccac	960
cgccctgacc	ccacaggccc	tgggctggac	agagagcagc	tgtatttggg	gctgagccag	1020
ctgaccaca	gcactactga	gctgggcccc	tacacactgg	acagggacag	tctctatgtc	1080
aatgggttca	cccctcggag	ctctgtaccc	accaccagca	ccgggggtgt	cagcgaggag	1140
ccattcacac	tgaacttcac	catcaacaac	ctgcgctaca	tggcggacat	gggccaaccc	1200
ggctccctca	agttcaacat	cacagacaac	gtcatgaagc	acctgctcag	tcctttgttc	1260
cagaggagca	gcctgggtgc	acgggtacaca	ggctgcaggg	tcctgcact	aaggtctgtg	1320
aagaacggtg	ctgagacacg	ggtggacctc	ctctgcacct	acctgcagcc	cctcagcggc	1380
ccaggctctg	ctatcaagca	ggtgttccat	gagctgagcc	agcagaccca	tggcatcacc	1440
cggctggggc	cctactctct	ggacaaagac	agcctctacc	ttaacggtta	caatgaacct	1500
ggtccagatg	agcctcctac	aactcccaag	ccagccaaca	cattcctgcc	tcctctgtca	1560
gaagccacaa	cagccatggg	gtaccacctg	aagaccctca	cactcaactt	caccatctcc	1620
aatctccagt	attcaccaga	tatgggcaag	ggctcagcta	cattcaactc	caccgagggg	1680
gtccttcagc	acctgctcag	acccttgttc	cagaagagca	gcattggggc	cttctacttg	1740
ggttgccaac	tgtctccct	caggcctgag	aaggatgggg	cagccactgg	tgtggacacc	1800
acctgcacct	accaccctga	ccctgtgggc	cccggtctgg	acatacagca	gctttactgg	1860
gagctgagtc	agctgaccca	tgggtgcacc	caactgggct	tctatgtcct	ggacagggat	1920
agcctcttca	tcaatggcta	tgcaccccag	aattttatca	tccggggcga	gtaccagata	1980
aatttccaca	ttgtcaactg	gaacctcagt	aatccagacc	ccacatcctc	agagtacatc	2040
accctgctga	gggacatcca	ggacaaggtc	accacactct	acaaaggcag	tcaactacat	2100
gacacattcc	gcttctgcct	ggtcaccaac	ttgacgatgg	actccgtgtt	ggtcactgtc	2160
aaggcattgt	tctcctccaa	tttggacccc	agcctgggtg	agcaagtctt	tctagataag	2220
accctgaatg	cctcattcca	ttggctgggc	tccacctacc	agttggtgga	catccatgtg	2280
acagaaatgg	agtcatcagt	ttatcaacca	acaagcagct	ccagcaccca	gcacttctac	2340
ctgaatttca	ccatcaccaa	cctaccatat	tcccaggaca	aagcccagcc	aggcaccacc	2400
aattaccaga	ggaacaaaag	gaatattgag	gatgcgctca	accaactctt	ccgaaacagc	2460
agcatcaaga	gttatttttc	tgaactgtcaa	gtttcaacat	tcaggctctgt	ccccaacagg	2520
caccacaccg	gggtggactc	cctgtgtaac	ttctcgccac	tggctcggag	agtagacaga	2580
ggtgccatct	atgaggaatt	tctgcggatg	acccggaatg	gtaccagct	gcagaacttc	2640
accctggaca	ggagcagtg	ccttgtggat	gggtattttc	ccaacagaaa	tgagccctta	2700
actgggaatt	ctgaccttcc	cttctgggct	gtcatcctca	tcggtctggc	aggactcctg	2760
ggactcatca	catgcctgat	ctgcggtgtc	ctggtgacca	cccgcggcg	gaagaaggaa	2820
ggagaatata	acgtccagca	acagtgcaca	ggctactacc	agtcacacct	agacctggag	2880
gatctgcaat	gactggaact	tgcggtgcc	tggggtgcct	ttccccagc	caggggtccaa	2940
agaagcttgg	ctggggcaga	aataaacat	attggtcgga	cacaaaaaaa	aaaaaa	2996

&lt;210&gt; 206

&lt;211&gt; 914

&lt;212&gt; PRT

&lt;213&gt; Homo sapien

<400> 206

Met	Ser	Met	Val	Ser	His	Ser	Gly	Ala	Leu	Cys	Pro	Pro	Leu	Ala	Phe
1				5					10					15	
Leu	Gly	Pro	Pro	Gln	Trp	Thr	Trp	Glu	His	Leu	Gly	Leu	Gln	Phe	Leu
			20					25					30		
Asn	Leu	Val	Pro	Arg	Leu	Pro	Ala	Leu	Ser	Trp	Cys	Tyr	Ser	Leu	Ser
		35					40					45			
Thr	Ser	Pro	Ser	Pro	Thr	Cys	Gly	Met	Arg	Arg	Thr	Cys	Ser	Thr	Leu
	50					55					60				
Ala	Pro	Gly	Ser	Ser	Thr	Pro	Arg	Arg	Gly	Ser	Phe	Arg	Ala	Trp	Ser
65					70					75					80
Leu	Phe	Lys	Ser	Thr	Ser	Val	Gly	Pro	Leu	Tyr	Ser	Gly	Cys	Arg	Leu
			85					90					95		
Thr	Leu	Leu	Arg	Pro	Glu	Lys	Asp	Gly	Thr	Ala	Thr	Gly	Val	Asp	Ala
			100					105					110		
Ile	Cys	Thr	His	His	Pro	Asp	Pro	Lys	Ser	Pro	Arg	Leu	Asp	Arg	Glu
		115				120						125			
Gln	Leu	Tyr	Trp	Glu	Leu	Ser	Gln	Leu	Thr	His	Asn	Ile	Thr	Glu	Leu
	130					135					140				
Gly	Pro	Tyr	Ala	Leu	Asp	Asn	Asp	Ser	Leu	Phe	Val	Asn	Gly	Phe	Thr
145					150					155					160
His	Arg	Ser	Ser	Val	Ser	Thr	Thr	Ser	Thr	Pro	Gly	Thr	Pro	Thr	Val
				165				170						175	
Tyr	Leu	Gly	Ala	Ser	Lys	Thr	Pro	Ala	Ser	Ile	Phe	Gly	Pro	Ser	Ala
			180					185					190		
Ala	Ser	His	Leu	Leu	Ile	Leu	Phe	Thr	Leu	Asn	Phe	Thr	Ile	Thr	Asn
		195					200					205			
Leu	Arg	Tyr	Glu	Glu	Asn	Met	Trp	Pro	Gly	Ser	Arg	Lys	Phe	Asn	Thr
	210					215					220				
Thr	Glu	Arg	Val	Leu	Gln	Gly	Leu	Leu	Arg	Pro	Leu	Phe	Lys	Asn	Thr
225					230					235					240
Ser	Val	Gly	Pro	Leu	Tyr	Ser	Gly	Cys	Arg	Leu	Thr	Leu	Leu	Arg	Pro
				245					250					255	
Glu	Lys	Asp	Gly	Glu	Ala	Thr	Gly	Val	Asp	Ala	Ile	Cys	Thr	His	Arg
			260					265						270	
Pro	Asp	Pro	Thr	Gly	Pro	Gly	Leu	Asp	Arg	Glu	Gln	Leu	Tyr	Leu	Glu
		275					280					285			
Leu	Ser	Gln	Leu	Thr	His	Ser	Ile	Thr	Glu	Leu	Gly	Pro	Tyr	Thr	Leu
	290					295					300				
Asp	Arg	Asp	Ser	Leu	Tyr	Val	Asn	Gly	Phe	Thr	His	Arg	Ser	Ser	Val
305					310					315					320
Pro	Thr	Thr	Ser	Thr	Gly	Val	Val	Ser	Glu	Pro	Phe	Thr	Leu	Asn	
				325					330					335	
Phe	Thr	Ile	Asn	Asn	Leu	Arg	Tyr	Met	Ala	Asp	Met	Gly	Gln	Pro	Gly
			340					345					350		
Ser	Leu	Lys	Phe	Asn	Ile	Thr	Asp	Asn	Val	Met	Lys	His	Leu	Leu	Ser
		355					360					365			
Pro	Leu	Phe	Gln	Arg	Ser	Ser	Leu	Gly	Ala	Arg	Tyr	Thr	Gly	Cys	Arg
	370					375					380				
Val	Ile	Ala	Leu	Arg	Ser	Val	Lys	Asn	Gly	Ala	Glu	Thr	Arg	Val	Asp
385					390				395						400
Leu	Leu	Cys	Thr	Tyr	Leu	Gln	Pro	Leu	Ser	Gly	Pro	Gly	Leu	Pro	Ile
				405				410					415		
Lys	Gln	Val	Phe	His	Glu	Leu	Ser	Gln	Gln	Thr	His	Gly	Ile	Thr	Arg
			420					425					430		
Leu	Gly	Pro	Tyr	Ser	Leu	Asp	Lys	Asp	Ser	Leu	Tyr	Leu	Asn	Gly	Tyr
	435						440					445			
Asn	Glu	Pro	Gly	Pro	Asp	Glu	Pro	Pro	Thr	Thr	Pro	Lys	Pro	Ala	Thr

450		455		460
Thr Phe Leu Pro Pro Leu Ser Glu Ala Thr Thr Ala Met Gly Tyr His				
465		470		480
Leu Lys Thr Leu Thr Leu Asn Phe Thr Ile Ser Asn Leu Gln Tyr Ser				
	485		490	495
Pro Asp Met Gly Lys Gly Ser Ala Thr Phe Asn Ser Thr Glu Gly Val				
	500		505	510
Leu Gln His Leu Leu Arg Pro Leu Phe Gln Lys Ser Ser Met Gly Pro				
	515		520	525
Phe Tyr Leu Gly Cys Gln Leu Ile Ser Leu Arg Pro Glu Lys Asp Gly				
	530		535	540
Ala Ala Thr Gly Val Asp Thr Thr Cys Thr Tyr His Pro Asp Pro Val				
545		550		560
Gly Pro Gly Leu Asp Ile Gln Gln Leu Tyr Trp Glu Leu Ser Gln Leu				
	565		570	575
Thr His Gly Val Thr Gln Leu Gly Phe Tyr Val Leu Asp Arg Asp Ser				
	580		585	590
Leu Phe Ile Asn Gly Tyr Ala Pro Gln Asn Leu Ser Ile Arg Gly Glu				
	595		600	605
Tyr Gln Ile Asn Phe His Ile Val Asn Trp Asn Leu Ser Asn Pro Asp				
	610		615	620
Pro Thr Ser Ser Glu Tyr Ile Thr Leu Leu Arg Asp Ile Gln Asp Lys				
625		630		640
Val Thr Thr Leu Tyr Lys Gly Ser Gln Leu His Asp Thr Phe Arg Phe				
	645		650	655
Cys Leu Val Thr Asn Leu Thr Met Asp Ser Val Leu Val Thr Val Lys				
	660		665	670
Ala Leu Phe Ser Ser Asn Leu Asp Pro Ser Leu Val Glu Gln Val Phe				
	675		680	685
Leu Asp Lys Thr Leu Asn Ala Ser Phe His Trp Leu Gly Ser Thr Tyr				
	690		695	700
Gln Leu Val Asp Ile His Val Thr Glu Met Glu Ser Ser Val Tyr Gln				
705		710		720
Pro Thr Ser Ser Ser Thr Gln His Phe Tyr Leu Asn Phe Thr Ile				
	725		730	735
Thr Asn Leu Pro Tyr Ser Gln Asp Lys Ala Gln Pro Gly Thr Thr Asn				
	740		745	750
Tyr Gln Arg Asn Lys Arg Asn Ile Glu Asp Ala Leu Asn Gln Leu Phe				
	755		760	765
Arg Asn Ser Ser Ile Lys Ser Tyr Phe Ser Asp Cys Gln Val Ser Thr				
	770		775	780
Phe Arg Ser Val Pro Asn Arg His His Thr Gly Val Asp Ser Leu Cys				
785		790		800
Asn Phe Ser Pro Leu Ala Arg Arg Val Asp Arg Val Ala Ile Tyr Glu				
	805		810	815
Glu Phe Leu Arg Met Thr Arg Asn Gly Thr Gln Leu Gln Asn Phe Thr				
	820		825	830
Leu Asp Arg Ser Ser Val Leu Val Asp Gly Tyr Phe Pro Asn Arg Asn				
	835		840	845
Glu Pro Leu Thr Gly Asn Ser Asp Leu Pro Phe Trp Ala Val Ile Leu				
	850		855	860
Ile Gly Leu Ala Gly Leu Leu Gly Leu Ile Thr Cys Leu Ile Cys Gly				
865		870		880
Val Leu Val Thr Thr Arg Arg Lys Lys Glu Gly Glu Tyr Asn Val				
	885		890	895
Gln Gln Gln Cys Pro Gly Tyr Tyr Gln Ser His Leu Asp Leu Glu Asp				
	900		905	910
Leu Gln				



<210> 207  
 <211> 2627  
 <212> DNA  
 <213> Homo sapiens

<400> 207  
 ccacgcgtcc gccacgcgt ccggaaggca gggcagctc cactcagcca gtaccagat 60  
 acgctgggaa ccttcccag ccatggcttc cctggggcag atcctcttct ggagcataat 120  
 tagcatcatc attattcttg ctggagcaat tgcactcatc attggctttg gtatttcagg 180  
 gagacactcc atcacagtca ctactgtcgc ctacagctggg aacattgggg aggatggaat 240  
 cctgagctgc acttttgaac ctgacatcaa actttctgat atcgtgatac aatggctgaa 300  
 ggaaggtggt ttaggcttgg tccatgagtt caaagaaggc aaagatgagc tgtcggagca 360  
 ggatgaaatg ttcagaggcc ggacagcagt gtttgcgtgat caagtgatag ttggcaatgc 420  
 ctctttgcgg ctgaaaaacg tgcaactcac agatgctggc acctacaaat gttatatcat 480  
 cactttctaa ggcaagggga atgctaacct tgagtataaa actggagcct tcagcatgcc 540  
 ggaagtgaat gtggactata atgccagctc agagaccttg cgggtgtgagg ctccccgatg 600  
 gttccccag cccacagtgg tctgggcac ccaagttgac caggagacca acttctcgga 660  
 agtctccaat accagctttg agctgaactc tgagaatgtg accatgaagg ttgtgtctgt 720  
 gctctacaat gttacgatca acaacacata ctctgtatg attgaaaatg acattgcaa 780  
 agcaacaggg gatatacaag tgacagaatc ggagatcaaa aggcggagtc acctacagct 840  
 gctaaactca aaggcttctc tgtgtgtctc ttctttcttt gccatcagct gggcacttct 900  
 gcctctcagc ccttacctga tgctaaaata atgtgccttg gccacaaaaa agcatgcaaa 960  
 gtcattgtta caacagggat ctacagaact atttcaccac cagatatgac ctagttttat 1020  
 atttctggga ggaaatgaat tcatatctag aagtctggag tgagcaaaaca agagcaagaa 1080  
 acaaaaagaa gccaaaagca gaaggctcca atatgaacaa gataaatcta tcttcaaaga 1140  
 catattagaa gttgggaaaa taattcatgt gaactagaca agtgtgttaa gagtataag 1200  
 taaaaatcac gtggagacaa gtgcattccc agatctcagg gacctcccc tgctgtcac 1260  
 ctggggagtg agaggacagg atagtgcatt ttctttgtct ctgaattttt agttatatgt 1320  
 gctgtaattg tgccttgagg aagcccttg aaagtctatc ccaacatctc cacatcttat 1380  
 attccacaaa ttaagctgta gtatgtacct taagacgctg ctaattgact gccacttcgc 1440  
 aactcagggg cggtgcatt ttagtaattg gtcaaatgat tcacttttta tgatgcttcc 1500  
 aaaggtgcct tggcttctct tcccaactga caaatgccaa agttgagaaa aatgatcata 1560  
 atttttagcat aaacagagca gtcggcgaca ccgattttat aaataaactg agcaccttct 1620  
 ttttaaaaca acaaatgcgg gtttatttct cagatgatgt tcatccgtga atgggccagg 1680  
 gaaggacctt tcaccttgac tatatggcat tatgtcatca caagctctga ggcttctcct 1740  
 ttccatcctg cgtggacagc taagacctca gttttcaata gcatctagag cagtgggact 1800  
 cagctggggg gatttcgccc cccatctccg ggggaatgtc tgaagacaat tttgggtacc 1860  
 tcaatgaggg agtggaggag gatacagtcg tactaccaac tagtgataa aggccaggga 1920  
 tgctgtcaa cctcctacca tgtacaggac gtctcccat tacaactacc caatccgaag 1980  
 tgtcaactgt gtcaggacta agaaacctcg gttttgagta gaaaagggcc tggaaagagg 2040  
 ggagccaaca aatctgtctg ctctctcaaa ttagtcattg gcaaataagc attctgtctc 2100  
 tttggctgct gcctcagcac agagagccag aactctatcg ggcaccagga taacatctct 2160  
 cagtgaacag agttgacaag gcctatggga aatgcctgat gggattatct tcagcttgtt 2220  
 gagcttctaa gtttctttcc ctctattcta cctgcaagc caagttctgt aagagaaatg 2280  
 cctgagttct agctcaggtt ttcttactct gaatttagat ctccagaccc ttctggcca 2340  
 caattcaaat taaggcaaca aacatatacc ttccatgaag cacacacaga cttttgaaa 2400  
 caaggacaat gactgcttga attgaggcct tgaggaaatga agctttgaa gaaaagaata 2460  
 ctttgtttcc agcccccttc ccacactctt catgtgttaa ccactgcctt cctggacctt 2520  
 ggagccacgg tgactgtatt acatgttgtt atagaaaact gatttttagag ttctgatcgt 2580  
 tcaagagaat gattaaatat acatttccta caccacaaaa aaaaaaa 2627

<210> 208  
 <211> 282  
 <212> PRT  
 <213> Homo sapiens

&lt;400&gt; 208

```

Met Ala Ser Leu Gly Gln Ile Leu Phe Trp Ser Ile Ile Ser Ile Ile
      5              10              15

Ile Ile Leu Ala Gly Ala Ile Ala Leu Ile Ile Gly Phe Gly Ile Ser
      20              25              30

Gly Arg His Ser Ile Thr Val Thr Thr Val Ala Ser Ala Gly Asn Ile
      35              40              45

Gly Glu Asp Gly Ile Leu Ser Cys Thr Phe Glu Pro Asp Ile Lys Leu
      50              55              60

Ser Asp Ile Val Ile Gln Trp Leu Lys Glu Gly Val Leu Gly Leu Val
      65              70              75              80

His Glu Phe Lys Glu Gly Lys Asp Glu Leu Ser Glu Gln Asp Glu Met
      85              90              95

Phe Arg Gly Arg Thr Ala Val Phe Ala Asp Gln Val Ile Val Gly Asn
      100             105             110

Ala Ser Leu Arg Leu Lys Asn Val Gln Leu Thr Asp Ala Gly Thr Tyr
      115             120             125

Lys Cys Tyr Ile Ile Thr Ser Lys Gly Lys Gly Asn Ala Asn Leu Glu
      130             135             140

Tyr Lys Thr Gly Ala Phe Ser Met Pro Glu Val Asn Val Asp Tyr Asn
      145             150             155             160

Ala Ser Ser Glu Thr Leu Arg Cys Glu Ala Pro Arg Trp Phe Pro Gln
      165             170             175

Pro Thr Val Val Trp Ala Ser Gln Val Asp Gln Gly Ala Asn Phe Ser
      180             185             190

Glu Val Ser Asn Thr Ser Phe Glu Leu Asn Ser Glu Asn Val Thr Met
      195             200             205

Lys Val Val Ser Val Leu Tyr Asn Val Thr Ile Asn Asn Thr Tyr Ser
      210             215             220

Cys Met Ile Glu Asn Asp Ile Ala Lys Ala Thr Gly Asp Ile Lys Val
      225             230             235             240

Thr Glu Ser Glu Ile Lys Arg Arg Ser His Leu Gln Leu Leu Asn Ser
      245             250             255

Lys Ala Ser Leu Cys Val Ser Ser Phe Phe Ala Ile Ser Trp Ala Leu
      260             265             270

Leu Pro Leu Ser Pro Tyr Leu Met Leu Lys
      275             280

```

&lt;210&gt; 209

&lt;211&gt; 309

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 209

```

His Ala Ser Ala His Ala Ser Gly Arg Gln Arg Gln Leu His Ser Ala
      5              10              15

Ser Thr Gln Ile Arg Trp Glu Pro Ser Pro Ala Met Ala Ser Leu Gly
      20              25              30

Gln Ile Leu Phe Trp Ser Ile Ile Ser Ile Ile Ile Ile Leu Ala Gly
      35              40              45

Ala Ile Ala Leu Ile Ile Gly Phe Gly Ile Ser Gly Arg His Ser Ile
      50              55              60

Thr Val Thr Thr Val Ala Ser Ala Gly Asn Ile Gly Glu Asp Gly Ile
      65              70              75              80

Leu Ser Cys Thr Phe Glu Pro Asp Ile Lys Leu Ser Asp Ile Val Ile
      85              90              95

Gln Trp Leu Lys Glu Gly Val Leu Gly Leu Val His Glu Phe Lys Glu
      100             105             110

Gly Lys Asp Glu Leu Ser Glu Gln Asp Glu Met Phe Arg Gly Arg Thr
      115             120             125

Ala Val Phe Ala Asp Gln Val Ile Val Gly Asn Ala Ser Leu Arg Leu
      130             135             140

Lys Asn Val Gln Leu Thr Asp Ala Gly Thr Tyr Lys Cys Tyr Ile Ile
      145             150             155             160

Thr Ser Lys Gly Lys Gly Asn Ala Asn Leu Glu Tyr Lys Thr Gly Ala
      165             170             175

Phe Ser Met Pro Glu Val Asn Val Asp Tyr Asn Ala Ser Ser Glu Thr
      180             185             190

Leu Arg Cys Glu Ala Pro Arg Trp Phe Pro Gln Pro Thr Val Val Trp
      195             200             205

Ala Ser Gln Val Asp Gln Gly Ala Asn Phe Ser Glu Val Ser Asn Thr
      210             215             220

Ser Phe Glu Leu Asn Ser Glu Asn Val Thr Met Lys Val Val Ser Val
      225             230             235             240

Leu Tyr Asn Val Thr Ile Asn Asn Thr Tyr Ser Cys Met Ile Glu Asn
      245             250             255

Asp Ile Ala Lys Ala Thr Gly Asp Ile Lys Val Thr Glu Ser Glu Ile
      260             265             270

Lys Arg Arg Ser His Leu Gln Leu Leu Asn Ser Lys Ala Ser Leu Cys
      275             280             285

```

Val Ser Ser Phe Phe Ala Ile Ser Trp Ala Leu Leu Pro Leu Ser Pro  
 290 295 300

Tyr Leu Met Leu Lys  
 305

<210> 210  
 <211> 742  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <222> (1)...(742)  
 <223> n=A,T,C or G

<400> 210  
 cattgggtac gggcccccctc gagtcgacgt atcgataagc ttgatatcga attcggcacg 60  
 aggcccgacc gctccctgag agccagcaac gggcagtgat gtttagcccc gaggaaaaat 120  
 tacatgcgga atggaaagca ggcgctcagg gtggctcctg ctggaatgag agctggagtg 180  
 caggetccgt gggtcctggg catgcgggtg tggctcagtt ctcaccttgc agatggagtg 240  
 ggactgttga cccaggccag cctggggact gcctcctcac ctccctgcgc aggctgacct 300  
 tgtcaccttg cctccttgagc ttgcctctct cctgcccaga ngctccttga gcaaaatgga 360  
 ggtcgagagg catttggcac tcacgcctca ccacggacac tgggtgcattc ttgggtacct 420  
 cttggscctca atctattgct gggggganga ngactganc ccattgctgg ggccctgaat 480  
 gcagggactg taaccaccca tccccttctc agggcacctc tcccctctcca gcacncttgc 540  
 tttgctatta atgctaccta atttctact gangtggtct agaagctcct ccgccattgc 600  
 ccttgccgcc agcaaat tttt tatccctagg gttaagataa cagaaggcan ccttgggcct 660  
 tgctgcac attctcaggt ntncactgaa gcacagtatc tatttctcca aaaatagggg 720  
 ctgtnaactt gttactaccc cc 742

<210> 211  
 <211> 946  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <222> (1)...(946)  
 <223> n=A,T,C or G

<400> 211  
 ggcacgaggc acatcgctgg atttctcatt gccaaagctct attaatcat tctttttcat 60  
 aacctcttat tcttatttca tggatgcaac attttctttg tctctcaggg aataataatt 120  
 attcctactt ttaaaggctct aatttcttta ttactttatt tctctgggag tgagtttttc 180  
 ctaaagggat aatgagatgg aaaatgaaaa aacaaagtgt agacatggag ataccttctg 240  
 aaactcaagc attcctctac gtggatgtgc cagagggaaa gaacagaaca aaggagggtta 300  
 gacactattt aaataaaaa atataagaat attacataac aaacaaaaaa gcccaaatcc 360  
 tcagggttgaa aaggaggaga aaatgtcaag caagacaaaa acagatgaag caacaaaaaa 420  
 agtgacatag ctggtcacct atattgaaat ttcagaacat gactgataaa ggactcccag 480  
 aaaaaaacaa aacccaaact aaaaaacaga aaaaaaggac tttaccaccn aaaacttgan 540  
 gaatcaggaa gactcagctc ctcatlaaga aaantgctat aggggatggg ggcaaggcct 600  
 tcaaagtngc aggggatacc aataacctct ctgaagtgtt ggaacttcat actccaaaat 660  
 ngaatttttg tttgaatagc cccggttagg ggccaatttt aggacttaga aaggaccnng 720  
 gnaaatcatt ccnncttgc ccccccgaa agaaattaat agaaggggtt tattcccgcc 780  
 attannaanaa aaggaatcca ggaattnccg nttttttcca gtgttangnt gggngtgan 840

aaactgaggg cttagcaagg gcggnattaa ccaccnnggg tcccacccca aaantggnng 900  
gggtgggccc caaatcggg nttntncct ttaangcgtt aaacce 946

<210> 212  
<211> 610  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(610)  
<223> n=A,T,C or G

<400> 212  
ggcacgaggt ttctggctgg agcctcggac actggctcac tgcagttggt ggtgtcgaca 60  
gtggtangag ggcaaccagt aacgggagct tctcctgccca ggcaggaaga cgagtagaag 120  
ggagcggcat gctggaggct ggagcctgag cccctggggc tcgccttgct gtgtttggtg 180  
gtgacgtggg acactgcagc tcggccagag tggtaaaaaa tgtcctggtg tacgcttttc 240  
tggctttgcc cgtctatctg ctccaagcca ggctgganga ngagganaag gaatcacctg 300  
tggtagcgtg gagcctgcat gtggcgtgac tctgcaactc gcctcgtgtg actgatggca 360  
gccacggaga ctgcagctcg acagggagtg aggccttctca ntggcttgaa agctcagctg 420  
actcccacga aatttgccgg aaactcaagg ctgtcagtga cnttcgtggc gccaaagactt 480  
aancangcgc gttgcatgca tccggccagt gtctgtgcc a cgtgccctga cnccaccttg 540  
anataancac ccggaacgcg cnnccgcgag gccgcgcgca cacgnccggg cancaacttg 600  
gctggcttcc 610

<210> 213  
<211> 438  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(438)  
<223> n=A,T,C or G

<400> 213  
ccganagcgg tttaaacggg ccctctagac tcgagcggcc gccctttttt tttttttttg 60  
aaataaattt ctgattattt tattacataa gcagaccact gaaacattta ttcaaaagta 120  
ttccattgag agtcaaaaac atattgatat gattattatt ggtctgttaa agaaaacaaa 180  
ataaaaagaa caaactggga attatcaata aacaaatcaa aacttagatg taattataac 240  
ctaaagggct cacagggcaa atgtgaagca agcttctgtc tcagagcctg catatggaag 300  
acatgtagta cttagctttg gcattcttct ttctcctct tggttgagtt taagtattaa 360  
taaaagggtg actgagaaaa ccttttttta caatcttatg gggatttttt agtggaaacg 420  
ttttagaagt aggaatat 438

<210> 214  
<211> 906  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(906)  
<223> n=A,T,C or G

```

<400> 214
gccctctaga tcgngcggcc gccctttttt tttttttttt gaaataaaatt tctagattat 60
ttattacata agcagaccac tgaaacattt attcaaaagt attccattga gagtcaaaaa 120
catattgata tgattattat tgggtctgtta aagaaaacaa aataaaaaga acaaactggg 180
aattatcaat aaacaaatca aaacttagat gtaattataa cctaaagggc tcacagggca 240
aatgtgaagc aagcttctgt ctcagagcct gcatatggaa gacatgtagt acttagcttt 300
gncatctttc ttctctctc ttgnttgagt ttagtattaa taaaagttgg actgagaaaa 360
ccttttttta caatcttatg gggtattttt agtggaaacg tttagaagta gaataacat 420
attaaaactg cncagaacaa atgnggtgca tctcaaattg nggtccattt tcaaaatatg 480
aacacatatg ggcagcantt ttttttttaa aaagtcagaa ggggcctnct catgccctt 540
tccacttctt cactcattgg nccttcaacc caagcttaac tactntcctg acctccaaca 600
tcataaacta gtttccnagc tttgaaactt tttccaatg agtcntaccg gaatagatgn 660
tcacagaanc ctcttaaaaa ttttgacccc tgcccgggnt ntaaaaaggg tgcaataaac 720
ccaccaacat cttggctggg ggggcagggg ccaaaagaan ttcccaaac cgtttttgat 780
naaaaaaggg gacttttgaa aaaaaaatta aaatttttgc cagnaaagca tgggnccccc 840
cccttgaana aacccccctgc atnaaaccaa cnttntggga ntttttngg tanggtttt 900
ctggct 906

```

```

<210> 215
<211> 312
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<222> (1)...(312)
<223> n=A,T,C or G

```

```

<400> 215
ggcacgagga aaccagggtg gctgggtttt ggggtgtaaac ttaaaaatga caatcagcat 60
gagctggccg tgggctgttg ggggtgtagg ggcattcttg taagggaacc ctgctcagt 120
ccctctctgt tctgggtggg aggacaagga gggccaatag gggccaatag ggaggctgct 180
gctaggangg ttctctaaaa gaacagggtg agggctaggg ctggttctta gttcagggtg 240
ctctgggcag tgatttatat ccacacacct ttctgcaaaag tgcctaagg aganggcagg 300
gataggagtg tc 312

```

```

<210> 216
<211> 341
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<222> (1)...(341)
<223> n=A,T,C or G

```

```

<400> 216
taagcctntc gaanataatg aatgagtcn ggagaggctn atgangaaat nccaaacacc 60
tgactaatng gtgccacatg attncaatgg nctanacatg ggtagatct cntcngnga 120
atgagcaata acacnttaa antcntcaat tgacctagac acttcacact tgaaanatca 180
tcacttttna ngaccacgaa tgatgcttaa gaatcacatt ttgtgnngaa ntggantctg 240
gctacttaca cgaacagatt cttattcctg ttcattgagcc agtagaccgc gaanaagact 300
taagagcttc tganctttct cttagctcca nngcttgaan g 341

```

```

<210> 217

```

<211> 273  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <222> (1)...(273)  
 <223> n=A,T,C or G

<400> 217  
 nnccttcncc ccttnacnga catgaacaaa acagcngtct ngaaatttta ttaacattnn 60  
 aagggttacn ctccctnctt ntgttttccg ntaaannta nacctgcgcg gggcgggccg 120  
 atncagccct atagtgaagaa gcctaattnc agcacactgg cggccgttac tanngnatcc 180  
 cgactcggta ncaanttttg gngtaaagat ggacatanct ctatccnnga gnactcgtca 240  
 nccnttctct atnttacatg cnctaacgna gac 273

<210> 218  
 <211> 687  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <222> (1)...(687)  
 <223> n=A,T,C or G

<400> 218  
 ttttcagtgc tgttttggtc tcaattttga tgtcaaaatc tctgggttct tctaancnng 60  
 ttatgttctt ccancaaatc ctccagttt ttgtaatttt tttctatata agaagcgccct 120  
 gancceaatg cccaattnat acaccggtct tctccggaac gcttggtcna aagggtntag 180  
 tcnattnggc tcctggaagc atctnaaatg ctccaggtta ctcccangnc cctggannac 240  
 ttcaantgtc tanacgaatc ctggttttgc agcgggtcct gatatcgcaa ggaaatacgg 300  
 taataattat ccaagctctc ttcccaactna gganttcgga tctcatcagc cgggttaaagg 360  
 aaaactctc angaagtgtg ggcttccctt ccgggtctacc ggctaattgt aggaattact 420  
 tctggtctct ttccgataca tctctctctc aaagtnaaga aggttaaaag aatnttaacn 480  
 tctccagtg gctaattggtc aaacaccatc ctcatnagtc agactggggt ttcgaaagga 540  
 ggatataacc tccttgcnag tttnaattaa aagggattaa ccanatggac tanccctcnc 600  
 cccgggattt nctctctcac aggagaaggg gtctcncnc ttggtctatc cgaagcatag 660  
 gcaaaccnccn gggaattttc agaaacc 687

<210> 219  
 <211> 247  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <222> (1)...(247)  
 <223> n=A,T,C or G

<400> 219  
 gggcccttcn cctttnaatc gagagatcca aggttcaagg catgaaatac cagnctataa 60  
 aatgtctcaa gacntaaata atacggatng ngatagagag gttgaataat aaatgaanaa 120  
 anatgaaagn nattatgngg gaatacnaaa aaancngact aanggcggca ctgctgggca 180  
 tggnaaatc ggattaattc ctcataggac agccnaacc cttaaaatct cantttccgt 240  
 nacccga 247

<210> 220  
<211> 937  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(937)  
<223> n=A,T,C or G

<400> 220  
cgggctcgag tgcggccgca agcttttttt actatagacc aatattaaag tcagttaagt 60  
tccaaataca ganttggaag actaaagtaa aatatttaaat gggagaatat ctgcatctga 120  
atatgtcaac tgtttgctat ttttcagcta ttttaactct ctacctgtat ctcaaaaaca 180  
aatttaaaaa ttaatagatt tgacagcaaa atcattcagc actttactta ctccatcagc 240  
aagggtattta tgtagtcatt tccatccatg tggccaaact gaaaatccct aaccaccacc 300  
aaccaaaaat aaataaataa aaggagaggg ggtgggggga gagagagaga gaaagctcat 360  
taaatagtaa aaaagtaaat aaaacaatga agttaaatcc aggcctcagt aggccagaa 420  
actgtaaaaca tttcacatgt aaatcatata caataaacac tgctaaaagt gtaaattcta 480  
ctggcttctg agatacaaat acacgagtag aggaattctt aagacatttc tacttggttt 540  
atgcatattt aaaattcagg gaaatatcag ctattctacc tgaaatatgt ttaagaaaaa 600  
ttcctatttt ctctaaaaaa aggaataatc agaagacgct acatactatg taagaaaact 660  
atacaatgac ccatcattag aagattcaga ataggaaaga aataataatt cactaataaa 720  
atatatttat attgactgtc tttttttatg atagcaacaa tgattcagca taaagtaaaa 780  
atatatgtat ttccgatgcc attttttatt cagttattct tttgagtttc tgttagaata 840  
attatctgcc tatctctgac ttctgancag tcatttatgt ccaattataa gtacatgtgc 900  
atattttatt acctaaacg cctctcaaat cctttca 937

<210> 221  
<211> 353  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(353)  
<223> n=A,T,C or G

<400> 221  
ggctatnnna tnnntntaan atcntgncnn ccttgacgct gttantaaan aaaaacaaac 60  
gaatatacctt tttttgctcc cccctgtncg gataactaat tcacactaat acttacagta 120  
taactnttcc tttcaactac caatattaag ttccaagcca cctgggctta agtatcccaa 180  
caacttaggt aatttggtgc taaccacat actatatgct aattataaca ctctaagccc 240  
caaggaattt ttgttcagat ttcttatant ttccacttat aaatatnatt ccncctctat 300  
gggtatatnn nncctctagn cccatatnnc ccacngggat ttgttgaggg ggc 353

<210> 222  
<211> 813  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(813)



<223> n=A,T,C or G

<400> 222

```
ggcacgaggc tttactaagg ccagactcac tatccccgct tctgttctgt ggtacactgt 60
tactcctca gtccatccta acctgacttc ctggccactg cagctcttcc gataagggtc 120
agcagtggct tagttattgc taaataataa gcgcacatgc actccctctt tcctgaaaca 180
ttgtccctcc ttggtttctg ttcttctcta ggtctcctat cactcctcct tagtcttctg 240
tgccgacttc tgttcttctt gccctttaaa agtttggtatt ttccaggatt ctgtcctagg 300
cccacttact tctcattctg cacgttcttg ttggatgatt ctatcacatc cctaacttct 360
gctgcccagt atgcacttaa aattcccaaa tctgtatatc tggatctggc ctgtgtctct 420
agcctagaag tgtgctttat ccagaaagca cctcaaacac tgcactttgg aaattaagct 480
tactgagtct cgagtctcaa gtcccaaact gacttctttt tctctatttt ggtagtgac 540
aacactattt attcagtcac gcaaaccaga gccctgagaa ccactttaca ttctctttct 600
ccctttactc agttcttget tctgttcttt ctcctcncc tctcctgcct gtgggcctag 660
ngncattaa ctgggtggca ctgctttact ttcnattttt ttggctganc taaccnaag 720
ancctnttgt aggggccttt ctntcaggcn tnacttctnn caagancccc cgaaaccaga 780
tccnggggan tgctatggnn tggaaatatt ttg 813
```

<210> 223

<211> 882

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<222> (1)...(882)

<223> n=A,T,C or G

<400> 223

```
tcacactact gagaagcagg gaaacccact gaaagggcac gtttcttaac ctcagaatgg 60
ggctactagc ctctaaagca ggaattgcgt tttgtttagt atttccatgg tctgctgcaa 120
ggcgtggcct ttaccaaatg gataaatgcg tacaaggctc ttgtgagcag tcaagtttct 180
cgaggtttac agttgaaggg aagtgggatt gtttctctgc gcatttaaat gaaggtagg 240
gggtgatcac ctttctttaa atgtgtgaag ggatgagata aagagatagg catcttaatt 300
gccactgatg gccttcagggt gaggacaggc atgagccaac tgaagctttg acaattgtgc 360
tgaacccaaa acttcaaaaa caagaaaaaa catagactgg ctgaaatgat ctaagtcaac 420
agagcatggc cagcgcttca tacaaggcag gaccacaggg gaacactgac agcccaggag 480
gcactgagac agaggcagtg ggaagaagtg acagacccca gggactcccc accaacagca 540
gctgctgttg attaggaacc ccagtagac tgtcaggcac ctggtagtgg agaggctacc 600
aaggcccgga ctggagagga gccaaaggaa gaaacagtgc agtgcttaga cccctctggg 660
tctgcccggtg tccatacccc tagggagatt ccattccaga agtgagcata ttcccacaga 720
gtgctctggg ctactcatc acagctgccc ctncatgaag gcattctcac tgcagcetta 780
ncagggaaca gggtcatttg cattaggcan cttgctgtcc tagaaggcnt cggngtccc 840
tacactgccc atgttcccaa ngnggttcaa nctcnaaan tn 882
```

<210> 224

<211> 660

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<222> (1)...(660)

<223> n=A,T,C or G

<400> 224

```
gattaaactc aatcattcac ccgggctcga gtgcggccgc aagctttttt tttttttttt 60
tttttttttt ttttggncct ctgggcttgt gcccggaaagg ggantgctgg gccacntggg 120
tgtccgtgtt tgattttctg ggacctgccc ccccgnttcc cgccccggnt gccgcgtctc 180
actccccgcc gcggtgcnag ggccccctg tgccgcgcac ccttcaccc gtgttttgct 240
gtttttttga cnttgggcgt cccaggggtg cancgccgt ggggccctgg tttgctttca 300
cctcttcac tgctcactgg ccgcnantgn gtcttnttca aacaaacgn tgaaggnaaa 360
nccctgggct cctgtgaacc cggccgtctt tgccggcaaan tctgaggctc cttegttatt 420
ctggatccgg cctntggctg gangcgtgct ctgcaggcac tgcctccatt gctggcancc 480
ttttctcccc gtggccgcc ggccgcccat naaaggcgtt gcaaaccgcc gccctcgcca 540
gcgcaaagtc aaacnccggt ggccgcgga cccccggcg gncgggaaca cccancagg 600
cgggcaccac aanaagcgcg gncctccgc gtctaaaact nccatgtggc nccccccgn 660
```

```
<210> 225
<211> 438
<212> DNA
<213> Homo sapiens
```

```
<220>
<221> misc_feature
<222> (1) ... (438)
<223> n=A,T,C or G
```

```
<400> 225
aaaaaaaaag gaaaagtacc cagtgtcttc agcttctgag cctcctctac agccctgttg 60
gnttttaaac ctgtgccctg tgtctgtgtc cccacttaat atatatagta cacagctgga 120
gagatggctc agccaggaga gggaccata ggtctgtgaa ttccagagga naggcaggna 180
tttatagggt gntctgtcag gtgaaatcng aggagccaaa gctattgtat gtgcataatg 240
cagccgggct ctgtgggagg tgggtgaaga cctatggnat gggacangtg tncacgctgg 300
gatctctggc cgttccgaa aagtgaggat caggtagtgg gtggctgatt gcacaagttt 360
anaaccagg attagggaca cacaggtcag cacctgcttc tcagcatcct gactgggtgt 420
gatgggcata ctcaaggc 438
```

```
<210> 226
<211> 480
<212> DNA
<213> Homo sapiens
```

```
<220>
<221> misc_feature
<222> (1) ... (480)
<223> n=A,T,C or G
```

```
<400> 226
aaaattaaaa caaaaggat cttagaggtc ctttacttca gtggttctca atgtcagagg 60
atgttatgat acctaatcaa aatctccagg ggaactgttt tgaactcaac agactctctc 120
ctgttctgag agactctggc aaagttggga gagctgccag gtactgtcca catgacctg 180
actgcccatt attcaattac cttgaatggc ttatccagtc caataccttc atttcttaca 240
tgaggaaact gaagcacgta tcacatagtg atacaatgaa aacttggcct taatcgattt 300
tcagtgtctg cagtacaatg tcttgagcat atcaatttct tccaaccctt gacaacataa 360
ggtaacacca tcaaattttt tatttctgct aatttattag accaaaaaaa aagggnatct 420
cnccattgt tttacaggga tgattttatt ncagaggatt tcactntggn gctgattcnt 480
```

```
<210> 227
<211> 423
<212> DNA
```

<213> Homo sapiens

<220>

<221> misc\_feature

<222> (1)...(423)

<223> n=A,T,C or G

<400> 227

```
cattgtgttg ggctctgctt agcacatcac atcggagcac agaggtgacc tgttctgcc 60
cagggatgtt caccttagtc acctgattga ttcctcttca ctttggtcac gtgattcctc 120
caggaggatg ttcaccttgg tcgcttgatt cctccaggag gatgttcacc ttggtcgcct 180
gaccacacag gcatctatca ggctttctca ctgcagccac tatgtcccca taatggatga 240
gtgtcttgtg gagagatagt ccaaagaca ctgatacctt ttgcctcata cggcctcacc 300
ccccacaat cnaccactaa tgactgcctc atagcagttt ttccatttcc acagttcctt 360
ctatatgtat taattgtcat tctactataa agaanacttt ttctttttaa aaaaaaaaaa 420
aag 423
```

<210> 228

<211> 249

<212> DNA

<213> Homo sapiens

<400> 228

```
cattgtgttg ggctgtagta aaatatgtgt ctggaagat atgtgaagaa ataaaaaag 60
atcaattaaa tctggcccat tgaatgacac attaatgtta tattaatatg taatgttaa 120
gatattagga gatggtggga cattatggca aactaaattt gggaggagggt tgaattgtat 180
aatttatgaa atcctaaagt ctagtacatt aacactctct actgtcaact tttcaaagca 240
gtgagaaac 249
```

<210> 229

<211> 436

<212> DNA

<213> Homo sapiens

<400> 229

```
cattgtgttg ggatgttatc tgaccatcac aatatgattt ataatatgga ggcatgaagt 60
catttctcat tggggcagga gtgtggcaag ggggaagaag agctttacca attaaactcaa 120
gattatttgg tgacatttct cttacctttt aggtgaggag aaagagacag aggatggaga 180
attggtgctt ttagtatgct gatacattaa gctgcctgga agcagatgct aaatcctatt 240
gaaaaataatt ttatttgcgt tttgcttagg gcattgttta gcaaaatact acacaaaaag 300
tcttgacctg tgtgtttgaa atggcagatg ttcacagtga ggactgagcc ttggggcaac 360
atcaatcttc acaattctgc acctatttgc tcaataactg gcttggttgg aaaaaaagg 420
aaaaaaaaa aaaaag 436
```

<210> 230

<211> 760

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<222> (1)...(760)

<223> n=A,T,C or G

<400> 230  
cattgtgttg ggnngtgga gaaaaanttt gaggaatga agctaaacat aaaagaggaa 60  
aagcanatgt tacctcaatg accacaatct acaaagtcca aatanaaaac ctgggagtat 120  
gataggatga aactataacc tccagcaaag agcttaacag caattaaaat aaagacaaat 180  
ttctgggatg gatnagacaa agtagcatat attacaaagg aaaatanact agtatcatnt 240  
acgttttgatt aagtaactgc tttcaaataa ttgaatcata aacaatgatt tctgcgggtt 300  
taagtcatt attttggttc cctgggttct cctaggatgc agtatagaat ctccatgcct 360  
gatgtttatg taccaacaga agctgctgct tctttcttct attatttctt ttttaagtga 420  
aagttaatac cttttatatg ttacagagaa gaggcagaaa aagccacact cccactatgc 480  
tattaaatgc cctgaggatc aactgaggga tgattatacn catggctgaa tacagtntat 540  
tcatttgttt ctttggattg tanataacaa aaggtggtat tctgtaacat cttgtgncaa 600  
ttanccaaat gttaaggcga aaatggaatc tttcaaacaa gtgtntaaa caggttttga 660  
ttttccaaaa tttantatta gaacnnttct aattctggaa gttncccaat ttccangttg 720  
tgttttctct tccaattctt ctttctttg naaatcccc 760

<210> 231  
<211> 692  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)... (692)  
<223> n=A,T,C or G

<400> 231  
cattgtgttg gggggtgctn tggggagaac acgcttatgt tganatnggg ctccccgaga 60  
aagcctcatt gacacnttcg aataaggacc cntngggaaa ttcangtgag ttgtggacat 120  
nntagataa natcaaaggc cttgangaag tccgcctggc accttcnctg ctgcgaggag 180  
gttgatacca aatgctaagg ggtccagntg cantgtanta tctgagatc agagtgatgg 240  
gcagggtgtg gcatgcgggc cctcaanang aagtgccag gatgactcag acttatgcct 300  
atateccatt antcctgttc attattttta ncnttcctc naaggacccc caatttnaac 360  
catttgttat tcanggttat acttataaaa gtcatttgtt ttnagtctgg gtgatattaa 420  
aaccatttgg acgccangca tgggtggctn nggcctataa tcctntccac cttgggggaa 480  
ccgaagctgg ttnaatccct naaggtcngg aatttgaaaa ccatcctggg ncaacattgg 540  
gnгааaccct gtctctactn caaaaaacan aaaattttct ggggcctngg ttngcaggtg 600  
gcctgaaaat ttccancnt tactccggga aggcgaatg ccntaaaaaa mnnaccttta 660  
acccccccga angggcgga agtttccatt tn 692

<210> 232  
<211> 518  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)... (518)  
<223> n=A,T,C or G

<400> 232  
actcaaatgn ccncttgaag gtcaccaga ctcanaangt gtcaagcttt ggggtgggtn 60  
gtaatnaata nctcggntc ctgattagtn ctcctagctc gatcnctggc tgagatnngt 120  
tcgagacccc ttcctttgat cccgtcaaac nccnggnaaa agcngcctgc gtatcncct 180  
nagccgaatc tgntttcccg acaccctccg ctcggtcggc tgccctggtg aagcngcntc 240  
ctnaaanaaa aaagngaagt ctccccngtc tcnccnctant cctngggaaa acngcctgaa 300  
ccaatatgnt cccccaaggc cccccaggg cactaaccg gttaggaggg cccccnctg 360

```
gcgttttggc cnaagcccn gcccngnaa taaccccnct anaaccacgn aaaaatgcaa 420
agtcccaaag ggtaaagaat ctcccnaccc cccggttccc tcgcaanctt cccctnngna 480
cttgtgttcc gggaaaaccc ttancccgan cctttcca 518
```

<210> 233  
<211> 698  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(698)  
<223> n=A,T,C or G

```
<400> 233
gcacgagttt ctgtctgtct gtctctctct ctctctctct ctctctctgt ctctctctca 60
cagttagaat ttggtctggt tctttattca ataccccaat atatgttcat tagggttata 120
ctgtatacac tacacataac agttttgttt tttgttttgg atattatttg ataataagaa 180
ttttaccaca tcattaaaaa aagtttcccc aagctataat ttttgataat tgcactcttc 240
cactattcaa atgtttattt aactctttct ctctctggag aggtttacat tccatttttag 300
ctatgatact gctttaagag aaattgtttt aagataaatt tccatagaca ggtcaaagga 360
ggtgaatata tgtaagcttt tcgatgcctg ttactgaatc tcattctgga aaacataact 420
gtcaatgcc tctttttctc atggtaaaaa aatacataac aaaatttacc atcttaatcg 480
tttttaaatg ttacagtacg atagtgttna ctgtatgtac cttgtgcaac agattctctg 540
aaaacttttt catttttcaa aatgaaaact ctgtactcat tgaacaggca gcttcccaac 600
ttccccatc ctcccnccc ctaccctggg ttaanagtct nacaaaaccc gggaatttta 660
tgaaatttga aacactttta naataccnct tattaggg 698
```

<210> 234  
<211> 773  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(773)  
<223> n=A,T,C or G

```
<400> 234
ggcacgagcg cagcttttgc aaagctgtaa tttgttttgt atcaaaagtc ctgcagtata 60
ttagtctcat tgcattttta agagtttcca agtgatcagt gatggttgc tgttttttag 120
tattacggtc ttatgtaatg ttcgaaaact agtcagtttg gtgctgtcgt acggggcgga 180
aagatcaggc caggcaaagt actctggccg ccaaagtaaa tgcttaaggc cgccaacgga 240
ttatgtcctg gggttcgatg agggccgtaa ttaggttgag ctggtgtang ctaacctcgc 300
agccatgtcg gagagagatg agagacataa nattttaaag taggggcgta ttttacgaag 360
ttctgancca ttctctttgt tatcgggtccc ggcaaaagca actgagataa atgtgttaaa 420
agactcgatg attttttcga cttcagcaac gtactcagcc ttgggttctc gtagtttttc 480
aaaggcagct atttgctgag attcatgaaa agtttgactt ganctgcttg tcaatttctg 540
cagcncgggc ttcaactgtt attgaatttg tttgattaag cncaatacgt tgcnggtcac 600
caagggtttc catgttttga ctncacctgg tcgaaccaat ttgaattatg tntttttgcc 660
tgnccgtgtc ccccnccctt aaatccatct cttttttnga aacctttgng nggttgaatt 720
cngccgcccc gttcccaacn tttgtttcna ccttggaata aaanatgggt agt 773
```

<210> 235  
<211> 849

<212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <222> (1)...(849)  
 <223> n=A,T,C or G

```
<400> 235
attgggtacg ggccccctc gagcagcctc cactgcaatg ccgctgaatc aagagacttt 60
tcaatacgtt ttatcagtga aaatgatgtg atctgaagag tcctatcttg agcactttgc 120
atgacatcca acgttaatgt ccacaacgtt cttagctgcc caacccttt atcggaagc 180
tccaaaggtg tgtgcaaacg ttctacggcg tcatgaaaag ctgaaaaatg ctgtgtcaac 240
actgcaccgc tgcgcattct caaaagcagc gcccttatag tctccgcatt cgaagacgat 300
aaccgcgta gaatagcctc ataatacatt ttgtagaagt caatcagagc tgtgttagga 360
acctttccat ccaaaacata cgactgtgag accacgtctg caaaagcaga cgtcacatta 420
tgcataatgcc ctcttaccgt cagccgatca tcctcactca tagcgacgcg agaaagctct 480
tggtccagct cgtgcacggt atccaattca gtaatectac gcaacgccgt ctgaatcgtg 540
ttcataagtt cagtttttaa gctcaaaact tcgtctctta ntttaccctt tgtgactttc 600
aaactgggag antcttcacc attttattaa tcgtctcttt gangganggc ccagcggttag 660
atctgcatcg ccagcggaat cgttactccc tcccatcctt cctccgggta acgcanntag 720
tttctccgaa gccttataat tagccgggga aagggaantt atttgcccca acaanggnat 780
cgcggncccg gtgggttaaaa ggaactgaaa taaaattaaa nccncttgg gggaaangcc 840
cgcatactg                                     849
```

<210> 236  
 <211> 310  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <222> (1)...(310)  
 <223> n=A,T,C or G

```
<400> 236
ggggtggggt gcttccgaaa nccggggccc ggccaacttg ttggcttggg aatattcttg 60
caagaaaatt tccagggcgg cgccaatttn atcaagcccg ggcgccctta aaccgaaaac 120
tctggcaggg tcaacccttt tcatgggcn ttgaaagctt gaagcgcccc aagttactcc 180
caagcttggt gcnnttgccg ttggggggcg gggaaaagt gaaaacacgg gcgntttgtt 240
gcccgccccg cgggcggttt nttacgccat cctgggaaaa ctttcagggt tggctgctta 300
cnaaaacggg                                     310
```

<210> 237  
 <211> 315  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <222> (1)...(315)  
 <223> n=A,T,C or G

```
<400> 237
gcacgagtn tttgtattta natnttgctt tgtttaangg aagaacacaa naatgcctg 60
ctaaagggat tctgtttggg tgcangctgc naggcgggaa aaaatcnaa tgtatnttgc 120
```

```

acaacangat tttttagaan tcagaactat gacatgaagt canncagggc actctacgac 180
tgaatttgcn gtgctgcctt cacangctcc ttntctgctc tntnctggca nengtgactc 240
ntacacgtcc tgganantan cctccctana aggaacgact ccgacacccc cccnntaccc 300
ctnaangttc atcng 315

```

```

<210> 238
<211> 510
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<222> (1)...(510)
<223> n=A,T,C or G

```

```

<400> 238
ngcacgagtn tttgttattt atatattgct ttgtttaaag gaagaacaca aaaatgccct 60
gctaaagggg ttctgttttg ttgcaggctg cnngegggga aaaaatcaaa gtgtattttg 120
cagaaaaatga ttttttanaa gtcagaacta tgacatgaag tcaagcaggg cactctagga 180
ctgaatttgc tgtgctgcct tcatatgctc cttgctcgct cttttctggc agctgtgact 240
cncacaggtc atggaganta tcattcccta aaaggaacaa cnccgatatt catctttatc 300
cattaagtnc atctgtccca ttctatgtng tggatgctaa cttttgatca ttgatngtga 360
tnccatggac atntancatc anctttcana ncctnggac tttgacnagt cttattantn 420
agantccaac tantacgatg ccganttana aatgctggnt ntccaattcc tactcaaata 480
nccnecatga acttccantc cccttgenna 510

```

```

<210> 239
<211> 209
<212> DNA
<213> Homo sapiens

```

```

<400> 239
ggtgcttttc ccttctactc gtcttctctc ctggcaggag aagctccgc tactggttgc 60
ccttctacca ctgtcgacac caccaactgc agtgagccag tgtccgaggc tccagccaga 120
aacaggtagc agccatgccg gataccaaac gccacacatt aagagcctga aatgacctga 180
cgccacctcc gcctgcttta cctactgag 209

```

```

<210> 240
<211> 610
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<222> (1)...(610)
<223> n=A,T,C or G

```

```

<400> 240
ggcacgaggt ttctggctgg agcctcggac actggctcac tgcagttggt ggtgtcgaca 60
gtggtangag ggcaaccagt aacgggagct tctcctgccg ggcaggaaga cgagtagaag 120
ggagcggcat gctggaggct ggagcctgag ccctcggggc tcgccttgct gtgtttggtg 180
gtgacgtggg acactgcagc tcggccagag tggtaaaaaa tgtcctggtg tacgttttc 240
tggctttgcc cgtctatctg ctccaagcca ggctgganga ngagganaag gaatcacctg 300
tggtagctg gagcctgcat gtggcgtgac tctgcaactc gcctcgtgtg actgatggca 360
gccacggaga ctgcagctcg acaggagtg aggtttctca ntggcttgaa agctcagctg 420

```

```
actccccagca aatttgccgg aaactcaagg ctgtcagtga cnttcgtggc gccaaagactt 480
aancangcgc gttgcatgca tccggccagt gtctgtgcca cgtgccctga cnccaccttg 540
anataancac ccggaacgcg cnnccgcgag gccgcgcgca cacgnccggg cancaacttg 600
gctggcttcc                                     610
```

```
<210> 241
<211> 474
<212> DNA
<213> Homo sapiens
```

```
<220>
<221> misc_feature
<222> (1)...(474)
<223> n=A,T,C or G
```

```
<400> 241
ggcacgaggt ttctggctgg agcctcggac actggetcac tgcagttggg ggtgtcgaca 60
gtggtangag ggcaaccaat aacgggagct tctcctgccca ggcaggaaga cgantagaan 120
ggancggcat gctggangct ggancctgan cccctggggc tcccttgctg tgtttggtgg 180
tgacgtggga cactgcagct cggccagant ggtaaaaatg tccctggtgta cgcttttctg 240
gctttgcccg tctatctgct ccaagccacg ctggaagang agganaagga ntcacctgtg 300
gtacgcccgga gccctgcatgt gggngtgact ctgcaactcg cctcgtgtga ctgatggcac 360
ccacggacac tgccactcta cagngaataa ggcttctccn tggactngaa agctcanctt 420
nactcccncc aagtttgncc gaactcaagg ctntcactna atttcgtggc gcca 474
```

```
<210> 242
<211> 415
<212> DNA
<213> Homo sapiens
```

```
<220>
<221> misc_feature
<222> (1)...(415)
<223> n=A,T,C or G
```

```
<400> 242
ngcgggggnt tccaccagct cgtgtgcaca agtngcgcca cacaacatg cgcaggcact 60
gcatgtcatc natgtgcttc gccgtggttc tggaaacagcg agtagaagat ggcgttcggg 120
tcgcgaccaa attcgacgtc ntggatgttc ttgcgcaaga angtcacgta cgggatcggc 180
ccgatggatc cgctnaagcg ccgaaaggcc ctgacttgca aaccgcggct cacagaaccg 240
gcaccaccgg cgccctccgc cnacaaaagt cgagcggcct ccgacacaca ctccctcaca 300
tccccgtcnc gcaattcggc ngtttctagc tccgccacgg ttgtcagcgg caccgcgggc 360
gccnagctgc cggcggcatc cgttgcacac agcacacacg gatccgctct cgtgc 415
```

```
<210> 243
<211> 841
<212> DNA
<213> Homo sapiens
```

```
<220>
<221> misc_feature
<222> (1)...(841)
<223> n=A,T,C or G
```

```
<400> 243
```



```

aacgaggtgt cgatgagcgc gaacaatcgc cctccttcat ctctacctga tggatgaactt 60
cgctcctaca gccgagccaa tgaagacgaa tggctgctgc cgaggatggg agtctcacta 120
gagcacgcgc cgctggacaa ctcatcgact tgtacgcttc cggtagctta gccattcag 180
ctccactgac gacagagacg gagctggcca ctgccatctc gacgcagcgc gacaaggagc 240
agcttcgggc gccgtatgca tcaactgaag agaaccagga gcagccggaa gcaggangcg 300
ctgcacggta caggcacttt cggcgcttca gcggatccat cgggcccgat ccgtacgtca 360
ccttcttgcg caagaacatc caggacgtcg aattcggtcg cgaaccgaat gccatcttct 420
actcgctctt ccaggaccgc gcgaagcaca ttgatgacat gcagtgcctt gcgcagtgtt 480
gtgcggcgct accttggcgc acacgaacga nggcaaccaa cccgcccag gtgcgcgtct 540
atgcattcct gttctgttcc ggtgtgcatg gccggatgtg gaccgtganc ttggtgaatc 600
ggctgggtgca tgaagactta ccgctctcnt caaggcgcaa cgcncctcan ttcgganaag 660
gaacaaaacc ccccnnaag aacggcantt gcancntttt ccccgctgc cggctcttct 720
ccattcgggn attctctntc tcnaaaant ccgnaaatc ttcttctcgt ttctccctg 780
tttttatttg ccttccgcgc cacttgggtt gttttacatc ctacaancct ttttttctc 840
c

```

841

```

<210> 244
<211> 761
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<222> (1)...(761)
<223> n=A,T,C or G

```

```

<400> 244
aacgaggtgt cgatgagcgc gaacaatcgc cctccttcat ctctacctga tggatgaactt 60
cgctcctaca gccgagccaa tgaagacgaa gtggctgctg ccgaggatgg gagtctcact 120
agagcacgcg gcgctggaca actcatcgac ttgtacgctt ccggtagctt agccattca 180
gctccactga cgacagagac ggagctggcc actgccatct cgacgcagcg ggacaaggag 240
cacttcggg cgccgtatgc atcaactcga gagaaaccagg agcagccgga agcaggagcg 300
gctgcacggc acaggcactt tcggcgcttc agcggatcca tcgggcccga cccgtacgtc 360
accttcttgc gcaagaaaca tcaggacgt cgaattcggc cgcgaccga atgccatctt 420
ctactcgctc ttccaggacc cggcgaagca catttgatga actgcagtgc ctgcgcagt 480
ttgttgccgc gctacctggt tgcacncgan cganggcaac aaccgcgcc angttgccgc 540
tctatgcatt cctgtctgt ccggtgttgc atggccggat gtgganctg ancttgtaa 600
tccgctgggt gcatgaagga cttaccgctc tcgtcaagg cgaacgcgcc atcaattccg 660
gaaaaggaa naaaaccccc cccaangac ggnaatttgc ancttttccc ncncctgccg 720
gctcttctcc antnccggct tctctttctc anaaaattcc c

```

761

```

<210> 245
<211> 710
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<222> (1)...(710)
<223> n=A,T,C or G

```

```

<400> 245
aacgaggtgt cgatgagcgc gaacaatcgc cctccttcat ctctacctga tggatgaactt 60
cgctcctaca gccgagccaa tgaagacgaa gtggctgctg ccgaggatgg gagtctcact 120
agagcacgcg gcgctggaca actcatcgac ttgtacgctt ccggtagctt agccattca 180
gctccactga cgacagagac ggagctggcc actgccatct cgacgcagcg ggacaaggag 240

```

```

cagcttcggg cgccgtatgc atcactcgaa gagaaccagg agcagccgga agcaggaggc 300
gctgcacggt acaggcaactt tcggcgcttc agcggatcca tcgggcccgat cccgtacgtc 360
accttcttgc gcaagaacat ccaggacgtc aaattcggtc gcgaccgaat gccatcttct 420
actcgctctt ccaggaaccg gcgaagcaca ttgataacat catgcctgcc catgtttgtt 480
gcggccctcc tggttgcnc aagaancgaag ggcaacaaac ccgcgccagg tngccgctct 540
tatgcattcc ttgtctgttc cggtnntgca tggcccggan nttggaaccg tnancttgg 600
mnaatcggt ggtgcattga aggaacttac cgtctcgtc aagggccgaa cgcnccttc 660
agttcggana aaggancgaa aaccccccn naaggaacgg cnttgcnnng 710

```

&lt;210&gt; 246

&lt;211&gt; 704

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc feature

&lt;222&gt; (1)...(704)

&lt;223&gt; n=A,T,C or G

&lt;400&gt; 246

```

aacgaggtgt cgatgagcgc gaacaatcgc cctccttcat ctctacctga tggatgaactt 60
cgctcctaca gccgagccaa tgaanacgaa ntggctgctg ccgaggatgg gactctcact 120
aaagcacgcg gcgctggaca actcatcgac ttgtacgctt ccggtagctt agccattca 180
gctccactga cgacaganac ggagctggcc actgccatct cgacgcagcg ggacaagggg 240
gcagcttcgg gcgcccgtatg catcactcga agagaacagg agcagccgga agcaggaggc 300
gctgcccggg acaggcaactt tcggcgcttc ancgatcca tcgggcccgat cccgtacgtc 360
accttcttgc gcaanaacat ccaggacgtc gaattcggtc gcgaccgaa ttgccatctt 420
ctactcgctc ttccagggac cggcgaagca cattgatnaa attgcattgc ctgcgcattg 480
ttgtgcgggg ctctctgggtg ccccgancga agggcnacaa ccccgcgcca ggggtccnct 540
ctatgcattc ctntctgttc cgggtgttgc tgggcgggat ttgaaccgtg aancttgggtg 600
aatccgnttg gtgcattaag aacntaaccg ttentcgtca ggggcennacc ggncccttnc 660
aatttcggaa aaangaacca aaanccccc cnccaagga aacn 704

```

&lt;210&gt; 247

&lt;211&gt; 618

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc feature

&lt;222&gt; (1)...(618)

&lt;223&gt; n=A,T,C or G

&lt;400&gt; 247

```

ggccgccagt gtgatggata tcgaattcaa cgaggtgtcg atgagcgcga acaatcgccc 60
tccttcatct ctacctgatg gtgaacttcg ctctacagc cgagccaatg aagacgaagt 120
ggctgctgcc gaggatggga gtctcactag agcacgcggc gctggacaac tcatcgactt 180
gtacgcttcc ggtagcttag cccattcagc tccactgacg acagagacgg agctggccac 240
tgccatctcg acgcagcggg acaaggagca gcttcggggc ccgtatgcat cactcgaaga 300
gaaccaggaa gcagccggaa gcaggaggcg ctgcacggta caggcacttt cggcgcttca 360
gcggatccat cgggcgcgac ccgtacgtca ccttcttgcg caagaacatc caggacgtcg 420
aatcgggtcg cgaccggaat gccatcttct actcgctctt ccaggaccgg gcgaaagcac 480
attgatgaca tgcagtgcct gcgcatgttt gtngcggcgc tacctggtgc acacgagcga 540
nggcaacaaa cccgcgcccc ggtgccgctc tatgcattcc tgttctgtcc ggggtgtgcat 600
ggcccgatg tggaacct 618

```

<210> 248  
<211> 622  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(622)  
<223> n=A,T,C or G

<400> 248  
gcacgagagc ggatccgtgt gtgctgtgtg caacggatgc cgccggcagc ttggcgcccg 60  
cgggtgccgt gacaaccgtg gcggagctag aaactgccga agtgccgcgac ggggatgtga 120  
gggagtgtgt gtcggaggcc gctcgacttt tgttgccgga gggcgccggt ggtgccggtt 180  
ctgtgagccg cggtttgcaa gtcagggcct ttcggcgctt cagcggatcc atcggggccga 240  
tcccgtagct gaccttcttg cgcaagagca tccacnacgt cgaatttggt cgcgaaccga 300  
acgccatctt ctactcgctc ttccagaacc cggcgaagca cattgacaac atgcnntgcc 360  
tgcgcatggt tgtgcggcgc tncctgntgc acacgaccga gggtagcaac ccgcgccagg 420  
ntgcncctct acgcattcct gtctgcccgg tgtgcgtggc cnggatgtgg accntgagcn 480  
ggngantccg ctggtgcntg aagacnttgc cgctctcgtc aaggccnacc gcccntcgcg 540  
gcggaaaaag gancaaaanc ccccgccaa gaaccggcnc tgcaccgttn tcgcgccct 600  
gctgggctct tctcctttac gg 622

<210> 249  
<211> 517  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(517)  
<223> n=A,T,C or G

<400> 249  
cattcgagct cggtagccgg gatccgattg gtaaaggggg tgcggaacag ccagctggtg 60  
ttttcgggtg ggcgggggca gccacatcg ctgtggtcgt tggcgtagtg gatgcgatgt 120  
gccgggacaa acgcgttttc caccacgatg tcatgactgc ctgtgccgag caggccagc 180  
acatcccagt tgcctcaat gcggtagtcc gccttgggca ccagaaaagt cacatgctcc 240  
aggccaggcg tgccatcacg cttgggcagc agaccgccta gaaacagcca gtcgcaatgc 300  
ttggagcccg tggaaaagct ccagcgaccg ttgaacctga atccgccttc cacgggctcg 360  
gccttgccag taggcatata ggtcgaggcg atgcgcacgc cgttatcctt gcccacaca 420  
tctgctggg cctggtcggg gaaaaancgc cagctgccaa ggggtgaacg ccgaccaccc 480  
cgtaaatcca ggccgtggac atgcagccct ttaccaa 517

<210> 250  
<211> 215  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(215)  
<223> n=A,T,C or G

&lt;400&gt; 250

mntncattgg gccgacgtcg catgctcccg gccgccatgg ccgcggggatt accgcttgtg 60  
accgcttgtg accgcttgtg accgcttgtg accgcttgtg accgcttgtg accgcttgtg 120  
accgcttgtg accgcttgtg accgcttgtg accgcttgtg accgcttgtg accgcttgtg 180  
accgcttgtg acnggggggtg tctggggggac tatga 215

&lt;210&gt; 251

&lt;211&gt; 231

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (1)...(231)

&lt;223&gt; n=A,T,C or G

&lt;400&gt; 251

ngcgccccacc tngtgattga tgggtcgttta ctatcaagta tgtacatctt gctctagaca 60  
actccnattc agtgggaagaa attgggaaag tatcccggt aagtaatagg nattaggtct 120  
nccttantgc ttggtgggat attccncaac tgnccngat cggatcagnc tcgtgtcngn 180  
gaatgtgctc gatcgtnatt ctactnctga gcttctatcc nnaegtggcc t 231

&lt;210&gt; 252

&lt;211&gt; 389

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (1)...(389)

&lt;223&gt; n=A,T,C or G

&lt;400&gt; 252

atgtatcanc nctgttggtg ttncatcttt tgcagtcngt tctaagggen gataantatc 60  
agagatgcta atgcatnttc tgccaggcca ncattgggtg cctatgcgta ctcttcttat 120  
cttctgaag agtcattctt ggnggatgtg tccccccctc tccacagtgt ttgcaagcgt 180  
taccacagcn tgcgnggcc gggaaggctn ncacatccgg gnagacttcc ccncgtntga 240  
atcgtntctn gaatctccgg cgtctccct naacctctg actnggacaa ngnccegtnt 300  
tccccntgt gaactngtan ccgccccctc tccccccctc agcctaancg ggaangaaga 360  
cngggtcnat ctngggcncc acaagaant 389

&lt;210&gt; 253

&lt;211&gt; 289

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (1)...(289)

&lt;223&gt; n=A,T,C or G

&lt;400&gt; 253

nggggcenna tgagcgcgcg taatacnatc actatngggc gaattgggta cgggcccccc 60  
tcnagcgcc gccctttntt nttttttnt tntttttnt caaaacaccc tccncntgg 120  
atgganacgt nacctttctc taaccanac ttcacaatnc nantctcagg cagccgctc 180

aaanccgatg tcangttggn atntcaantn caatcttatt ttgngaatta anctganatt 240  
gtggatggtn naccaatcan atacttgga tccgttgaac ccctgtgga 289

<210> 254  
<211> 410  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(410)  
<223> n=A,T,C or G

<400> 254  
attgtgttgg gaactttag acagctatat caattgcagt gctatttctc tgaggatttg 60  
aatctcantt attataattt tgaaatccaa ttggcttggga cttcattatt ttccaactaa 120  
aaagatgatt gaaggattta ttgaaatgt gtaaagagta atatagattt tatgcttatg 180  
tttccttgaa aaaagtaggt aaaattcttc tggaaagtgt actcctaaaa tacaaatgaa 240  
catgtcaaga attacataaa ttctttaaac tatccttaan aannaatggc tctatgtann 300  
gagngaccct tacagactat taagaattaa cttgcatggc anagactcat ttanattcat 360  
gaaatggnct tcactttctt ggtaagatct ggcttggacg tttttggtaa 410

<210> 255  
<211> 668  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(668)  
<223> n=A,T,C or G

<400> 255  
tttttttttt ttttctgtg ccaggcacta taccactgtg ctagggtgect tctttgcatt 60  
acttcatttc ctcataagct ttctgaggan acagaaagct tgagggttcac gtagctagca 120  
tctacataaa ttagtgtgcta aaaacataca atacgtcttc cggcaggctg tcattagtaa 180  
ctgatactac tagttgataa tctcataaac ctagcanaan ctaccattta agctgaaaca 240  
actgtcaata tcactaanta aaacttaaat ccataaatca actatattct aaaatctgac 300  
ttcagttcaa ttaaaaaatc actagttgtt acctacctec ttctgaaagc cagtacaagt 360  
taaatagaaca actcccaggt ttaacaaaca agtggcatct aaaaaaaga tttaaaaaat 420  
aatccactta catatattta aaatggcatt aataaaacaa aatttatcca ataacnaant 480  
ggcaaggaa ggtgtccaat tattacatgt tataaatctt taaattaaac ttttcttngg 540  
tttttctcc ctanaataaa tacaancctt tccccgccna accagaaaaa agcaaaaaac 600  
aaaacccaaa aactcccagc ncngcttaaa aaacncaaaa aaaataaaan ctctatttaa 660  
tgcccnaa 668

<210> 256  
<211> 487  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(487)  
<223> n=A,T,C or G

<400> 256  
cgnaaccgtn cntttttnat gtgcgcccgc cncagnacca gngccgctac aggcgaaggc 60  
cggaagcacg ggagaggntt nggaaaaaaa agagtgccta caaagagcat attcgagag 120  
ttgggatgag tgaaggggac cagaaggngc agcggtaggg acgcgtgaaa ggangcngcg 180  
gagaaatgac agcaagaagg gganaagcac acgaaaaggc agtatcctcc tcccccttt 240  
tcgaggactg ccgcattctt gttttctgcc cattccagtc accgaanaag atcccaana 300  
aagaagaaaa gaancagagg tgcacttcgc ttcattttc nctcgtttc ttttctgnet 360  
tcacnagttc tgcaggattg cccttgtcct ctcccgagca catctacgca cgnatgaggc 420  
tcggcaggtc aagccnacaa aacnctcgca ctccctcttt tctttgcngg tctgngtggt 480  
angngg 487

<210> 257  
<211> 502  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(502)  
<223> n=A,T,C or G

<400> 257  
ccttgaaag nccngctnaa ttcngnganc cccngatca gcaccaggga gctacaacna 60  
aggcgggaag caggggattt ngccggaaaa aaaagagtgc ttacaaagag nttatccnca 120  
nagatgggat gagtgaaggg gacgagaagg tgcagcggta gggacgcgtg aaaggaggca 180  
gcggagaaat gacagcaaga aggggagaag cacacgaaaa ggcagtatcc tctccccc 240  
tttctgagga ctgccgcac tttgttttct gccattcca gtcaccgaaa aagatcccaa 300  
agaaagaaga aaagaaacag aggtgcactt cgcttcatat ttcgctcgtt ttcttttctg 360  
tcttcacaag tctgcaggat tgccttctgc ctcttccgag cacatctacg cacgtatgag 420  
gctcggaggc caagccaaaa aaacgcttgc actcctcttt ttctttgctg gtctgtgtgt 480  
atgtggaatt ccgcggcncc gc 502

<210> 258  
<211> 510  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(510)  
<223> n=A,T,C or G

<400> 258  
actcgnact cgatncanta caagagnnta tgnattcgaa ngtgccccg catcagcacc 60  
aggagctac aacgaaggcc ggaagcaggg gagaggccg gaaaaaaaag agtgcttaca 120  
aagagcatat ccgcagagtt gggatgagtg aaggggacga gaagggtcag cggtagggac 180  
gcgtgaaaag aggcagcggg gaaatgacag caagaagggg agaagcacac gaaaaggcag 240  
tatectctc ccccttttct gaggactgcc gcatctttgt tttctgcca ttccagtcac 300  
cgaaaaagat cccaaagaaa gaanaaaaga aacagagggtg cacttcgctt catatttcgc 360  
tcgcttttct tttctgtctc caagtctgca ggattgccct tgcctcttcc cgagcacatc 420  
tacgcacgta tgaagctcgg aggtcnnngc aaaaaaacgc ttgcactcct ctttttcttt 480  
gcnagtctgt gtgcatgnng gaaatnctna 510

<210> 259

<211> 292  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(292)  
<223> n=A,T,C or G

<400> 259  
gannngagtc acgaaaaggc agtatcctcc tcccccttt tcgaggactg ccgcattcttt 60  
gttttctgcc cattccagtc accgaaaaag atcccaaaga aagaagaaaa gaaacagagg 120  
tgcacttcgc ttcatttttc gctcgttttc ttttctgtct tcacaagtct gcaggattgc 180  
ccttgctctc ttccgagcac atctacgcac gtatgaggct cggagggtcaa gccaaaaaaa 240  
cgcttgcaact cctctttttc tttgcgtgct tgtgtgtatg tggaattcct tg 292

<210> 260  
<211> 582  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(582)  
<223> n=A,T,C or G

<400> 260  
gcacgagggt ggggtgtact gtgtataata actccagatc cttgaccaag tttggagagt 60  
cacttatggc catttgaaac caaatgaagg atcaaaggac taattatttt gaatacctct 120  
gagtgttttc cccaagcttg agaagagttt cattcagcta taaaatgctc attgtgcaaa 180  
tgagtggttt ccatgctgta taattaaagc attgccttta ataatacttt attaccttta 240  
gcttgtcttt ttaatttgag gaaaatccaa acaattttaa gtaaaacgtg ataaagacag 300  
tttttcngga gananaaggg nagatcgcta tgtttattcc acttaatatc tatatcaaat 360  
atltgtatca aaagcagact ctcaacttta aaatattctt ctaatggcna gaatcttttn 420  
cctagattga gagtcagagc tcacatagna tnactgctgg taaatagaca cttagactat 480  
agagctnagc tnaagttcca actanccaac tgcatttctg aatatgcttt ttattnaaag 540  
gccagnnctt ttgccttttt nccnccctaa tnccttctat tg 582

<210> 261  
<211> 783  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(783)  
<223> n=A,T,C or G

<400> 261  
gcacgaggca aaatacagag ggtattttac catggacagg caaccatttt ttccaggaca 60  
actctttgca gcagagagct attctctttc ttttgcctta cactctcaac ctcaactctc 120  
gagtgtctgc atcctanttt tccatggcca taagataagg aaccatgagt gttactctag 180  
atgaggctgt ttcattgttg gagctcatcc aggatccaag gtagattcat cagaagggta 240  
agtataggag tgggaaccca aatctctact tttattttga ggccttctct cctcaatttt 300  
aaattgtaaa atcaaaactta aaactgggta tctgatggcc agttaaaga ctgggtatct 360  
gattgccagt taagagatgg tcatttatgc tcaccacat tctcaagacg cagggtgagg 420  
gacangcttg ctgggggaatg ctgancgaat cccccaatgc cttcaggatt ctgggaatgg 480

```
tggtctctgnt ttaaactggn tgactttttac aaagagccta cccgtcatgg ggggactggg 540
aagaaaaccc anangcagnt tctggcccan ggttacaccc ccanggn tac cttgaaggnt 600
ttttggacat acctnttnc cccctnttac tgnttcatta gggcntcnn cccccaant 660
tccaagttnt ggcccttcna aaanttttt ntttccntt tccanggacc cccctggnt 720
cctggnnccc cctttttata nccaaccttg ccnggnattt tttcncttn aaagggaaat 780
aat 783
```

```
<210> 262
<211> 741
<212> DNA
<213> Homo sapiens
```

```
<220>
<221> misc_feature
<222> (1)...(741)
<223> n=A,T,C or G
```

```
<400> 262
tgaaccctan tgggcccggc cccctcgagt cgacgggtatc gataagcttg atatcgaatt 60
cggcacgagt gtatattctg ttattatacc ccagattnaa gtgtatattc ttaggcagta 120
gttctgggta acatccttac tacataaaat ccacttacta tttaagtatt attctaacag 180
gaggtagaat agctgcctta aaaaatgtag tgatcgaatg gcagtttttc tgctgaatgg 240
aaattactga cacaaaattt gggtttggga gacattttcc tccttggtgt tgagttttcc 300
cattcacgga tagggcataa agcttggttt atagttgagg ggtgcaaaag ggggaatagga 360
ttgggaaaat acagtgttcc agcaaaggtc tgacaaggta catcttggag aggattccta 420
ttctgctang tggcactgta ngcttgaaa tactgtgtac tttccagaca aaggatagag 480
aaaaagacct tcactgggtg ggggagaaga aaacccttgt tcctagaaaa atcacaaaaa 540
aggcaccctt tancctatat tcccagnttt actgngcat ttgcttgatg tgactgacnc 600
ngattatttc ctttnactgg naaaaattcc tgcnccttg gatatnaang ggggnaccng 660
gaaaatnggg ggcnttgggg aaggaaanaa aaaaaattgg agggaccnaa ctttggaaaa 720
tggngtgctt nangccttaa g 741
```

```
<210> 263
<211> 437
<212> DNA
<213> Homo sapiens
```

```
<220>
<221> misc_feature
<222> (1)...(437)
<223> n=A,T,C or G
```

```
<400> 263
ggcacgagag aatgtgttca cagacactat tttatannta tctgatgtgt actgtgtctg 60
gtggatgtga aagccatact tcttaaactc gatttgaaaa gcaaatctga ttatcacagc 120
cataattaaa tttggccagc ctctcttcc cctccctcc ttcaacttcc tccttccctc 180
cgctcgtgc cgaattcggc acgagcctga cctcactacc aaaaaaaaaa aaattcaaag 240
tgctgaggt ttccaggcat tcttagctct atttacttac ttcccacctc aaatggcctt 300
agaattcaaa ttctgnanaa aatggattgc catanataat ccaatgaaaa tgggtcatat 360
tttgccatta atagaatcac agtcnacaag ggactaatag aattagtcac ttangtatcn 420
ttagatttgg gagacnn 437
```

```
<210> 264
<211> 706
<212> DNA
```



<213> Homo sapiens

<220>

<221> misc\_feature

<222> (1)...(706)

<223> n=A,T,C or G

<400> 264

```
gcacgagcac cccaagggtt taggacaaaa tgggatgagt gaattcatgg cttgacagac 60
tgaacagaaa aatgagggtc cgtgctccat attcatgtgc atctgcccct catggtgaca 120
tgctaattgg ttggccggtg cacaagacaa ggaagtgcag gtttctgtt gctcacacag 180
tgcttcctgt ctgctgtggc aggagccggg aggaaggag cgagccaaga ggggtgctgc 240
ccaccggaac cgatggcgcg aggcgcgaga gctaaatggg ggccctctca gggagtgtgc 300
tgttcacggc tccatcgctg ttagtaagta tcttgtgatt tcggaattta aatgaggttg 360
tgtttaacct gcataacatc tggcttttaa aatctgactt tattttcctt ttatttctgt 420
gcatcgggtc aggcacactt agtgggtggt taggtgttga agtcaggtta ccaaacagca 480
cgccctctct ttattctcag gctgcgtgtt tcattgattc tgaaggtcag atggctgtgt 540
tcaagttctg ttagtatatt ggtgtcagaa atgaaaagat gatgtaacct ttataaactt 600
cttaaaggct catatcatgt caggaaatta acctgtacga gttatggaca aatgcccatac 660
ctgatgatgt tcanccatga aaatgaatna aagggganaa gggcca 706
```

<210> 265

<211> 717

<212> DNA

<213> Homo sapiens

<400> 265

```
ggcacgagca gcattacggt ttatacacat gtccacaact cagcattgct ttcaaaatag 60
gaacacttta ttagtaaaga ggaagaaatt gcctaaacag actcagtgtc ttcccataa 120
caatcatctg ccaagccgca ggcctaacca ggaaatccca tttccttttg gcgttggtgc 180
ctccaccaac agatacaacc ctgatgccaa atgttgatg gttttaggt gttgtgagcc 240
aatgagggca tgcctaggc caaaggctgc cttttggaat gagggcaagg tcgtagactc 300
catcaaaaca caaatgcac ctcctccaaa atcaaatgct caacacatgc agcctttcgt 360
atgcccatct cccctttact ctttttcatg gctgaaaatc atcaggatgg gcatttgtcc 420
ataactccta cagggttaatt tcctgacatg atatgagcct ttaagaagtt ataaagggtt 480
acatcatctt ttcatttctg acaccaatat actaacagaa cttgaacaca gccatctgac 540
cttcagaatc aatgaaacac gcagcctgag aataaagaga gggcgtgctg tttggttaacc 600
tgacttcaac acctaaagca ccactaagtg tgcttgagcc gatgcacaga aataaaagga 660
aaataaagtc agattttaaa aagccagatg ttatgcaggg taaacacaac ctcat 717
```

<210> 266

<211> 362

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<222> (1)...(362)

<223> n=A,T,C or G

<400> 266

```
ggcacgaggt tagatttaac ttccacagat gactcagcag aggataacta ctaatcagag 60
tacaacatca aaactgtaac cagtataatc actggattat gagcaactca aaatagctcc 120
agtttccaaa gggccataaa ctgcacatat cagtactatg tgcaattaac acataattta 180
ttatgaaaat gtggacatgc caggtaagta aggggattta ggttgacttt ttataatact 240
ttaaatttga aatgccattt ctgtggattg gatgacatct tccagggtgct ntaatnctgg 300
```

gntacctnct gatanatcct gananaaaga ggtancacca gcgtctatca nacctcaata 360  
ca 362

<210> 267  
<211> 692  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(692)  
<223> n=A,T,C or G

<400> 267  
ggcacgaggt tagatttaac ttccacagat gactcagcag aggataacta ctaatcagag 60  
tacaacatca aaactgtaac cagtataatc actggattat gagcaactca aaatagctcc 120  
agtttccaaa gggccataac tggccctttt aanactttnn gcaattaaca cataatttat 180  
tatgaaaatg tggacatgcc aggtaagtaa ggggatttag gttgactttt tataataactt 240  
taaatttgaa atgccatttc tgtggattgg atgacatctt ccagggtgctt taatttggtt 300  
tacctcctga tagatcctga cagaaagagg naggaccagc gtctatcaaa cctcaatata 360  
gngtgtgaaa cacangagag cctgcttttg tcnacacggg gaaacacatt gttatcacia 420  
cacacaaaag gcaanctncc aatggggnan ncttacctgn cctctcatat tgggggcaan 480  
gaaaangggg ccccanatg gctgagtana tccccaaaaa cncactan tggtcagnnt 540  
gcttcccan acagccagat gactgaattt agcccaagct gcagtctcaa aaccagcttt 600  
ctgacaatca gtaacaagaa catactgggc tgttgcagtg agctcaagtg ttgggtgttc 660  
agtcaaaanc catggatgcc aatcatctcc ca 692

<210> 268  
<211> 605  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(605)  
<223> n=A,T,C or G

<400> 268  
cgtgccgaat tcggcacgag ngcacatc agtactatgt gcaattaaca cataatttat 60  
tatgaaaatg tggacatgcc aggtaagtaa ggggatttan gttgactttt tataataactt 120  
taaatttgaa atgccatttc tgtggattgg atgacatctt ccagggtgctt taatttggtt 180  
tacctcctga tagatcctga cagaaagagg tagcaccagc gtctatcaaa cctcaatata 240  
gttgtaaaac acagagagcc tgcttgccca cacatggaga aacattgtta tcacaagaca 300  
cagaaggcaa acttccaate tggcatactt ncctgtcttc tcatatttgg ggcaatgaga 360  
atgggtggacc agatggcttg antagatgcc aaagaacacc canactgggc agcatgcttn 420  
cccagacagc cngaagactg aaatttante ccagctgcag ncttaaacc tttttttgac 480  
nttccgtaac cagaccatac tttttttct gatgcttttc ttaacttcat cttttccaat 540  
taaattcatt agtnnaacc taaanggggc ccgttttccg aaaaatttcc nttntntntt 600  
cccn 605

<210> 269  
<211> 535  
<212> DNA  
<213> Homo sapiens

<220>  
 <221> misc\_feature  
 <222> (1)...(535)  
 <223> n=A,T,C or G

<400> 269  
 gcacgaggng caaccccagg gtgggtctct tgggatgaac ctggagacct gagcttgcac 60  
 agcttccttg gtaaattgag gaggcattgga ccacaagatt gccaaagctcc ttctatcca 120  
 aacttgatat tggtagattc catgatccag ttcacacagg ttgatggctg aatctcatgc 180  
 actanaaaaa ggtaatatata aaganaaaaa tanaangatn ttcaagttag tataaanacc 240  
 tttaatctca ntctttctag ttcaaagaga cggaacaatg agagatgctg gttcatanag 300  
 ctgntanatt taacttccac agatgactca ncagaggata actactaatc anagtacaac 360  
 atcaaaaactg taaccagtat aatcactgga ttatgagcaa ctcaaaatag ctccagtttc 420  
 caaagggcca taaactgcca tatcaantac tatgtgccat taaccataa tttattatga 480  
 aaatgtggac atgccangtn agtaagggga tttagggtga ctttttatna tactt 535

<210> 270  
 <211> 803  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <222> (1)...(803)  
 <223> n=A,T,C or G

<400> 270  
 gcacgagggc aaccccaggg tgggtctctt gggatgaacc tggagacctg agcttgacac 60  
 gcttccttgg taaattgagg aggcattggac cacaagattg ccaagctcct ttctatccaa 120  
 acttgatatt gtttagattcc atgatccagt tcatcacggg tgatggctga atctcatgca 180  
 ctagaaaaag gtaatatataa agaaaaaaat aaaaagatat tcaagttagt ataaagacct 240  
 ttaatctcag tctttctagt tcaaagagac ggaacaatga gagatgctgg ttcatagagc 300  
 tggtagattt aacttccaca gatgactcag cagaggataa ctactaatca gagtacaaca 360  
 tcaaaaactgt aaccagtata atcactggat tatgagcaac tcaaaatagc tccagtttcc 420  
 aaagggccat aaactgcaca tatcagtact atgtgcaatt aacacataat ttattatgaa 480  
 aatgtggaca tgccaggtaa gtaaggggat ttaggttgac tttttataat actttaaatt 540  
 tgaatgcca tttctgtgga ttggatgaca tcttccaggg gctttaattt ggtttacctc 600  
 ctgatatgac ctgacagaaa gaggtagcac cagcgtctat caaacctcaa tacagttgta 660  
 aaacacagag agcctgnttt gcctacncat ggagaacatt gttatcacia gacacagaag 720  
 ggaacttcca tctggctact tacctggctt tatttttggg gcaatganaa tngggggacc 780  
 aatggntgan tanatgcaa aaa 803

<210> 271  
 <211> 836  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <222> (1)...(836)  
 <223> n=A,T,C or G

<400> 271  
 gcacgagggc aaccccaggg tgggtctctt gggatgaacc tggagacctg agcttgacac 60  
 gcttccttgg taaattgagg aggcattggac cacaagattg ccaagctcct ttctatccaa 120  
 acttgatatt gtttagattcc atgatccagt tcatcacggg tgatggctga atctcatgca 180

```

ctagaaaaag gtaatatataa agaaaaaaat aaaaagatat tcaagtgagt ataaagacct 240
ttaatctcag tcttttctagt tcaaagagac ggaacaatga gagatgctgg ttcataagagc 300
tgttagattt aacttccaca gatgactcag cagaggataa ctactaatca gagtacaaca 360
tcaaaactgt aaccagtata atcactggat tatgagcaac tcaaaatagc tccagtttcc 420
aaagggccat aaactgcaca tatcagtact atgtgcaatt aacacataat ttattatgaa 480
aatgtggaca tgccaggtaa gtaaggggat ttaggttgac tttttataat actttaaatt 540
tgaaatgcc a tttctgtgga ttggatgaca ttttccaggt gctttaattt ggtttacctc 600
ctgatagatc ctgacagaaa gangtagcac cagcgtctat caaacctcaa tacagttgta 660
aaacacagag agcctgcttt gnctacacat ggagaaacat tgtatcaca gacacagnaa 720
ggcaacttcc atctgggata ctacctgtct ctctatttgg ggcattganat ggggacaatg 780
ntgananatg caanacacca atgngagctg ntccnacag cnatatgatt ntccat 836

```

<210> 272

<211> 203

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<222> (1)...(203)

<223> n=A,T,C or G

<400> 272

```

ggagaattgg gcccgtcang ggtgcattct gcatcacctg antttnaaat ctngatcaat 60
cnnctacta atantatcaa catnatttna acctgatctc cactgcttng tnattttcnn 120
ttcactgncc ctntcactng aacntctntt cacacagcca cccccatta tctggntggc 180
acctcncca aatncncct naa 203

```

<210> 273

<211> 594

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<222> (1)...(594)

<223> n=A,T,C or G

<400> 273

```

attcgggccn ctggatnctg gctcgagcgg ccgccgctgt gatggatatt tgcanaattc 60
ggcttctgga gagagctttn tttttgatgg ttgcangtac tctcgatgga gttggtgggt 120
gtggttatct ctctctgggt gtctttctgt ataaantctt tgcnctgact ncctanctcn 180
cctccccctg gtccttccct tagngtaaca nctggtaatc cctntcttct ttgctctcct 240
tncttctcct gancgatttc ctctntttgt ccactctcag gnanaaccct gntggtcagt 300
gttcattgact tcnngaagnt cgacccgcna aatagggncn cacggatnat gttgaancng 360
ggaagggagn gtccaanttc tctggttccan aggcctnagcc tagaganaat gatgggagan 420
gggtttactga gatcatngnn tcttctcgaa gatatnnttt aggggtggtcc cccataagng 480
aatttctcan cttcaaatct tctaatacat tactgaacan ctgncatttg ttacgccaca 540
nattgnaatt ctccatntct ttttagaaac nattncaagg tcatttattt ccct 594

```

<210> 274

<211> 229

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature  
<222> (1)...(229)  
<223> n=A,T,C or G

<400> 274  
ctactcactg tccggccatt tggncctctg natgcatnct caagcagcnc gccantatga 60  
tnnatatctg cacanttcag cttctngaga aaactatggt ttaaacagtt gcntanactt 120  
anaatanaaa tcgagtaagg tntagatnan tctctaacga tngaattatt ntacanaggg 180  
gtanncgatn accaggagta nctaganttg ancancancc taggtcnga 229

<210> 275  
<211> 651  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(651)  
<223> n=A,T,C or G

<400> 275  
atatctgntg aatacggntt cctgnaaaaa ggtntnattt agatggttga gtccgactca 60  
gcatgacgac ttggtgggtg tggtcantct cttatggttg agattgttca tgatatcatg 120  
ccctgagatg cctggactnn cctcaccgga gatcctagac ggtgntancc cctgagagtc 180  
tctctcctcc tgctctccta acttctccta atgatccctc cnattgtcta ctgtccnatt 240  
gaacccttct tgcttatgta tncaatcctt nacggtgtcc ctgctnantt tttganacga 300  
ngctcataat ggacngggga aggatagtnt gaataatntc ctgtataccc acgcnacnt 360  
ctacnctntg atctgacacg gtatactgat ttgtgctggt cnettcacca ttccantttc 420  
taccttccgc tcatatgctc tgtangctac accctctgtg actgctttct cagttacgtg 480  
caacaaggtn ttcatatctn gaactcttac accattctag anggatcncc cctcganaaa 540  
antttggaan aacaagcaag ancanaatnc ctctctngtg ntacacnanc cggettncgt 600  
atcctcgtnn aaggaattcc ccgctttcct gggctttaan tctcctaaac t 651

<210> 276  
<211> 392  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(392)  
<223> n=A,T,C or G

<400> 276  
accccccccg aattacgntg gccnatntaa aagtnatca ngcctccang caacntatcn 60  
tttcattacc acccacactc ctgttnnggg anggangtgg naatccttca ccatnetaat 120  
gtatgtgggt ctctcatgen ggtacgtata atctanncgt cccctnaaat cggatgcttc 180  
tgtaatcnnc agtcacnaaa ccacanggan caactgaaac angatttggc taacagccaa 240  
tgtctgggcc ctcncaatc cctnnaatat ctctacacc tgtagtanna atnaactacn 300  
ctacnctatt nnacacacgn ttaggttgt annaccaagc cntattgag tgaaatogtt 360  
tntatngtat naaatgccaa aagntgcggt aa 392

<210> 277  
<211> 212  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(212)  
<223> n=A,T,C or G

<400> 277  
ggtttgcggg natgaanttt gnaanaatna actttagnga taaccacccc accaatncct 60  
nctnagtatt tgncaacctn aaaactacag ctctctccag atagactntn ccttnctgat 120  
ttcaactctc cttggactgg tcagcctgaa ggggtgtaat gactcaccaa cgctactaat 180  
nccttnttna ctgtgccttn attttttcgc ct 212

<210> 278  
<211> 269  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(269)  
<223> n=A,T,C or G

<400> 278  
nnntccatcc taataccact cactatcggg ctccaancgg ccgcccgggc acgtntcttn 60  
tgngacagga tctgaatnaa ggggtggttg taacttnact naaaattctg aaatgatcct 120  
gcatcagaca ggggttctccg tntanaatan agtttcctg ttagttatcn agcctgggca 180  
ggggangana gattcgagga cntntgaaat gaaggnatta tttaggatgg gtgactcatt 240  
ccnaccnttc ncgctnacca gnccganga 269

<210> 279  
<211> 266  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(266)  
<223> n=A,T,C or G

<400> 279  
gttggtgant cngtttgng tcttctcgtt gntnggtggt tgggtgtgtg nnttggtgtn 60  
gggtngtntt tntggagaga gttgtagttc gtgagggttg cagtgtactt actatggagc 120  
ctaaggangt gngctaactt anantgatna ctttgctcat actgccctgc cctnaatgcc 180  
nngcttgctt caccctgggt ccnaaccnna tcgaacacct aacagtctag taggcttctt 240  
gctntancag actnctcttg aggate 266

<210> 280  
<211> 317  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(317)  
<223> n=A,T,C or G

<400> 280  
acactgttnag gtgnttgga ntgntgtagg catagncttt ntggcacaga gttggagccg 60  
tgaggcatag cntgtactta ctatggagcc taaggangga gctaacttat antnatnact 120  
ttgtcctaac tgccctgctc tnaatgccta ngcttgctc accctgntgc cttacnnnat 180  
cgaacaccta cgcggtctat aggtctcttg ctctatcagg actnctcttc nagcttcttc 240  
gcctcanttg actcactgtg ctcggtcgtt ctactngat ccagncgctc atnaacctna 300  
cttnggacgc aggtcat 317

<210> 281  
<211> 174  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(174)  
<223> n=A,T,C or G

<400> 281  
gnggtcatat tatacatcta aggcattggc aactccacgc cattatnaat tccatcgtag 60  
tgtccgcagt cactacttat aacctagatt aatagtgcct ggccccggac ngctctgtgca 120  
atctnccgcc ataccaattn cgatccnca accnecatna cactctctct tact 174

<210> 282  
<211> 169  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(169)  
<223> n=A,T,C or G

<400> 282  
atcgagctt gtacgatcgt catataacgc gcatgtgcgg atcgcttcag cgccgcccga 60  
ctgtcagaag gangagatct tttttatcac ttgtttgttt gactatanat aanancgact 120  
acagcattga tgtgtgtcct caaganttgt ctgggtctga naaagctga 169

<210> 283  
<211> 157  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(157)  
<223> n=A,T,C or G

<400> 283  
ggntntctaa gatcgagctt gtacgatcgt catatnacgc gcatgtgcgn atcgcttcac 60  
gtcgccnggc tgtccaggan atgcatntca acataatgtg cactctatat gggtattgat 120  
taatacgagn tangagcana tatcngatcac aacacaa 157

<210> 284  
<211> 133  
<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<222> (1)...(133)

<223> n=A,T,C or G

<400> 284

```
ggngtggtgt nagatacgca ngctgggacg aatcgnntca tagtacggcg catgtgttga 60
tcaattctga aaatccatcc cggcgcgctc ancatgcact anagggcaat cgcctatatg 120
antcgtatta caa 133
```

<210> 285

<211> 194

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<222> (1)...(194)

<223> n=A,T,C or G

<400> 285

```
ntntgngtga tgatacccaa gctggntacc nactngantc caattaccg ctcantntgc 60
tngaacaacgc ttogatngnc tcttgcatg tacttgaaac aggntanata tctaataagnn 120
tacngtgnn ttttcatca tacagnttnt atattncact nctnccatt cntttctant 180
ctctctctcc ntat 194
```

<210> 286

<211> 134

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<222> (1)...(134)

<223> n=A,T,C or G

<400> 286

```
gagggnttat gataccaagc tggtagcanc ccgtcactat nacggcccag tgtgtggatc 60
cgctanctgg tcncgcgatg tctacncaca cngaaactgc ctctcgnaa gatctcctct 120
cctctccnaa gaga 134
```

<210> 287

<211> 119

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<222> (1)...(119)

<223> n=A,T,C or G

<400> 287

```
tngggtatat ccagttgtac actggncata tacgcgcatt atgatcgttt cagccccgga 60
gtacggcatc attacganat ggnctcattc gtttaccttt ntcgctggac acaagcgtc 119
```

<210> 288



<211> 170  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(170)  
<223> n=A,T,C or G

<400> 288  
gggntgagat acncaagttg gtacgagtcg gatcatatna cggncgccat tttctggaat 60  
ccgcttacgt ggtcccggcg aagtactttt tcatgccttg caaaatngcg ttactgcact 120  
ancttgctta acctatgagt ggggtctttc ataccCNTc tntcatggaa 170

<210> 289  
<211> 126  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(126)  
<223> n=A,T,C or G

<400> 289  
ggccaattgg ggcctctana tgcntgctcg aacgggcgcc aatttnatgg atatctccaa 60  
aatteggctt accntggteg cggncnaagt acttaactca atccatctnt cactcaggat 120  
naatgc 126

<210> 290  
<211> 126  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<222> (1)...(126)  
<223> n=A,T,C or G

<400> 290  
ggccaattgg ggcctctana tgcntgctcg aacgggcgcc aatttnatgg atatctccaa 60  
aatteggctt accntggteg cggncnaagt acttaactca atccatctnt cactcaggat 120  
naatgc 126